

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Red
- Wrist Deviation: Red
- Pelvic Tilt: Yellow

Workstation Analysis

Pain & Risk Summary

Average Pain Level: 7

Main Pain Region(s): Neck, Elbows/Forearms, Wrists/Hands (VAS 8)

Posture & Ergonomic Corrections

- Reduce neck flexion to under 5°: Sit upright with your head aligned over your shoulders, keeping your chin parallel to the floor. Adjust your posture so your neck bends no more than 5° forward.
- Correct elbow angle to 90–100°: Adjust your arm position so your elbows are bent at a 90–100° angle, keeping forearms parallel to the floor and close to your body.
- Straighten wrists to neutral (0–10°): Keep your wrists straight and aligned with your forearms, avoiding any upward, downward, or sideways bending.
- Reduce pelvic tilt to under 5°: Sit with your hips level and your pelvis in a neutral position, ensuring your lower back maintains its natural curve.
- Raise monitor by 10 cm: Elevate your monitor so the top edge is at or just below eye level, reducing neck flexion to under 5°.
- Move keyboard 8 cm closer: Position your keyboard so your elbows remain at a 90–100° angle and your wrists stay straight while typing.
- Lower desk height to 72 cm: Adjust your desk so your forearms are parallel to the floor and your wrists remain neutral when typing.

Final Advice

Your assessment shows high ergonomic risk, especially for neck, wrist, and elbow regions, with significant ISO 9241-5 violations. Immediate workstation adjustments and posture corrections are critical. Perform the following exercises:

1. Neck Retraction: Sit upright, gently pull your chin back to align ears over shoulders. Hold for 5 seconds, repeat

10 times, 3 times daily.

2. Wrist Flexor Stretch: Extend arm, palm up, gently pull fingers back with opposite hand. Hold 15 seconds, repeat 5 times, 2 times daily.

3. Shoulder Rolls: Roll shoulders backward in a circular motion. Do 10 repetitions, 3 times daily.

4. Seated Pelvic Tilt: Sit with feet flat, gently tilt pelvis forward and back. Hold each position 3 seconds, repeat 10 times, 2 times daily.

5. Upper Back Extension: Interlace fingers behind head, gently arch upper back. Hold 10 seconds, repeat 5 times, 2 times daily.

Safety note: Perform all exercises slowly and stop if you feel pain or increased tingling. If symptoms worsen, consult a healthcare professional.



Annotated posture analysis from your uploaded photo