

Ergonomics Evaluation Report

Posture Analysis

- Neck Flexion: Red
- Shoulder Elevation: Green
- Elbow Angle: Red
- Wrist Deviation: Yellow
- Pelvic Tilt: Red

Workstation Analysis

- Chair: SeatDepth: Green

Pain & Risk Summary

Average Pain Level: 7


Main Pain Region(s): Neck, Elbows/Forearms, Hips/Glutes (VAS 8)

Posture & Ergonomic Corrections

- Raise monitor by 18 cm: Elevate your monitor so the top edge is at or just below eye level. This will reduce neck flexion from 65° to under 20°, in line with ISO 9241-5 recommendations.
- Bend elbows to 90-100 degrees: Adjust your arm position so your elbows are bent at 90-100°, not extended at 155°. This reduces strain on wrists and shoulders per ISO 9241-5.
- Reduce pelvic tilt to under 15 degrees: Sit with your pelvis more upright, using lumbar support if needed, to decrease pelvic tilt from 63° to below 15°, supporting your lower back and hips.
- Keep wrists straight, deviation under 5 degrees: Position your wrists so they remain neutral, with less than 5° deviation, to minimize strain and risk of tingling or carpal tunnel symptoms.
- Raise laptop screen by 18 cm: Use a laptop stand or stack books to lift your laptop screen, aligning it with your eye level to reduce neck and upper back strain.
- Use external keyboard at 5 cm from edge: Place an external keyboard 5 cm from the desk edge to allow elbows to rest at 90-100°, reducing wrist extension and upper limb discomfort.
- Add lumbar cushion, 5 cm thick: Place a 5 cm thick lumbar cushion at your lower back to support your pelvis and maintain a neutral spinal curve.

Final Advice

Your wrist and upper back pain are severe and associated with tingling. Please consult a healthcare professional if symptoms persist or worsen. Always perform posture exercises slowly and stop if you feel pain.



Annotated posture analysis from your uploaded photo