

CS 170: Algorithms

CS 170: Algorithms

Hello and ...

H . . . H . S . H H

CS 170: Algorithms

Hello and ...

H . . . H . . . H H
S

CS 170: Algorithms

Hello and ...

H . . . H . . . H
S H

CS 170: Algorithms

Hello and ...

H H
S H H

CS 170: Algorithms

Hello and ...

H
S H H H

CS 170: Algorithms

Hello and ...

S H H H H



CS 170: Algorithms

Hello and ...

S H H H H .



CS 170: Algorithms

Hello and ...

S H H H H . .



CS 170: Algorithms

Hello and ...

S H H H H . . .



CS 170: Algorithms

Hello and ...

S H H H H . . .

CS 170: Algorithms

Hello and ...

S H H H H

CS 170: Algorithms

Hello and ...

S H H H H

Please,

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides),

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story:

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay **still** until the end of class.

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay **still** until the end of class. Distracting!

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay **still** until the end of class. Distracting!

Thank you

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! ! ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! ! ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! ! ! ! ! !

CS 170: Algorithms

Hello and ...

S H H H H

Please, no laptops (unless lecture draft slides), ...

Story: “Shut”

If you must leave early, please sit by exit.

Please stay still until the end of class. Distracting!

Thank you ! ! ! ! ! ! ! !

Today.

...Complex numbers, polynomials today. FFT.

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

(A) 6

(B) 5

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

(A) 6

(B) 5

(A) 6

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

(A) 6

(B) 5

(A) 6 of course!

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

(A) 6

(B) 5

(A) 6 of course!

Coefficient of x^2 in result?

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Coefficient of x^4 in result?

(A) 6

(B) 5

(A) 6 of course!

Coefficient of x^2 in result?

Uh oh...

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4))$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4)) \quad = 4$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{r} x^0 \\ x^1 \end{array} ((1)(4)) = 4$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4))$$

$$x^1 \quad ((1)(3))$$

$$= 4$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{ll} x^0 & ((1)(4)) \\ x^1 & ((1)(3) + (2)(4)) \end{array} = 4$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4)) \quad = 4$$

$$x^1 \quad ((1)(3) + (2)(4)) \quad = 11$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2)) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3)) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4))) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2)) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & \end{array}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

x^0	$((1)(4))$	$= 4$
x^1	$((1)(3) + (2)(4))$	$= 11$
x^2	$((1)(2) + (2)(3) + (3)(4))$	$= 20$
x^3	$((2)(2) + (3)(3))$	$= 13$
x^4	$((3)(2))$	$= 6$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4)) \quad = 4$$

$$x^1 \quad ((1)(3) + (2)(4)) \quad = 11$$

$$x^2 \quad ((1)(2) + (2)(3) + (3)(4))) \quad = 20$$

$$x^3 \quad ((2)(2) + (3)(3)) \quad = 13$$

$$x^4 \quad ((3)(2)) \quad = 6$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$x^0 \quad ((1)(4)) \quad = 4$$

$$x^1 \quad ((1)(3) + (2)(4)) \quad = 11$$

$$x^2 \quad ((1)(2) + (2)(3) + (3)(4))) \quad = 20$$

$$x^3 \quad ((2)(2) + (3)(3)) \quad = 13$$

$$x^4 \quad ((3)(2)) \quad = 6$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d \quad \text{In example: } a_0 = 1, a_1 = 2, a_2 = 3$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$\begin{array}{l} a_0 + a_1x + \cdots a_dx^d \\ b_0 + b_1x + \cdots b_dx^d \end{array}$$

In example: $a_0 = 1, a_1 = 2, a_2 = 3$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d \quad \text{In example: } a_0 = 1, a_1 = 2, a_2 = 3$$

$$b_0 + b_1x + \cdots b_dx^d \quad \text{In example: } b_0 = 4, b_1 = 3, b_2 = 2$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d \quad \text{In example: } a_0 = 1, a_1 = 2, a_2 = 3$$

$$b_0 + b_1x + \cdots b_dx^d \quad \text{In example: } b_0 = 4, b_1 = 3, b_2 = 2$$

$$\text{Product: } c_0 + c_1x + \cdots c_{2d}x^{2d}$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d \quad \text{In example: } a_0 = 1, a_1 = 2, a_2 = 3$$

$$b_0 + b_1x + \cdots b_dx^d \quad \text{In example: } b_0 = 4, b_1 = 3, b_2 = 2$$

$$\text{Product: } c_0 + c_1x + \cdots c_{2d}x^{2d}$$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

Multiplying polynomials.

$$(1 + 2x + 3x^2)(4 + 3x + 2x^2)$$

$$\begin{array}{lll} x^0 & ((1)(4)) & = 4 \\ x^1 & ((1)(3) + (2)(4)) & = 11 \\ x^2 & ((1)(2) + (2)(3) + (3)(4)) & = 20 \\ x^3 & ((2)(2) + (3)(3)) & = 13 \\ x^4 & ((3)(2)) & = 6 \end{array}$$

$$4 + 11x + 20x^2 + 13x^3 + 6x^4$$

Given:

$$a_0 + a_1x + \cdots a_dx^d \quad \text{In example: } a_0 = 1, a_1 = 2, a_2 = 3$$

$$b_0 + b_1x + \cdots b_dx^d \quad \text{In example: } b_0 = 4, b_1 = 3, b_2 = 2$$

$$\text{Product: } c_0 + c_1x + \cdots c_{2d}x^{2d}$$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

$$\text{E.g.: } c_2 = a_2b_0 + a_1b_1 + a_0b_2.$$

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2.$

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

- (A) $O(d)$
- (B) $O(d \log d)$
- (C) $O(n^2)$
- (D) $O(d^2)$

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

- (A) $O(d)$
- (B) $O(d \log d)$
- (C) $O(n^2)$
- (D) $O(d^2)$

Time: $O(k)$ multiplications for each k up to $k = 2d$.

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

(A) $O(d)$

(B) $O(d \log d)$

(C) $O(n^2)$

(D) $O(d^2)$

Time: $O(k)$ multiplications for each k up to $k = 2d$.

$\implies O(d^2)$.

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

(A) $O(d)$

(B) $O(d \log d)$

(C) $O(n^2)$

(D) $O(d^2)$

Time: $O(k)$ multiplications for each k up to $k = 2d$.

$$\implies O(d^2).$$

or (D)

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

(A) $O(d)$

(B) $O(d \log d)$

(C) $O(n^2)$

(D) $O(d^2)$

Time: $O(k)$ multiplications for each k up to $k = 2d$.

$\implies O(d^2)$.

or (D) ..will use n as parameter shortly.

Multiplying polynomials.

Multiply: $(1 + 2x + 3x^2)(4 + 3x + 2x^2)$

Given:

$a_0 + a_1x + \dots a_dx^d$ In example: $a_0 = 1, a_1 = 2, a_2 = 3$

$b_0 + b_1x + \dots b_dx^d$ In example: $b_0 = 4, b_1 = 3, b_2 = 2$

Product: $c_0 + c_1x + \dots c_{2d}x^{2d}$

$$c_k = \sum_{0 \leq i \leq k} a_i * b_{k-i}.$$

E.g.: $c_2 = a_2b_0 + a_1b_1 + a_0b_2$.

Runtime?

(A) $O(d)$

(B) $O(d \log d)$

(C) $O(n^2)$

(D) $O(d^2)$

Time: $O(k)$ multiplications for each k up to $k = 2d$.

$\implies O(d^2)$.

or (D) ..will use n as parameter shortly. so (C) also.

Hmmm...

$O(d^2)$ time!

Hmmm...

$O(d^2)$ time!
Quadratic Time!

Hmmm...

$O(d^2)$ time!

Quadratic Time!

Can we do better?

Hmmm...

$O(d^2)$ time!

Quadratic Time!

Can we do better?

Yes?

Hmmm...

$O(d^2)$ time!

Quadratic Time!

Can we do better?

Yes? No?

Hmmm...

$O(d^2)$ time!

Quadratic Time!

Can we do better?

Yes? No?

How?

Hmmm...

$O(d^2)$ time!

Quadratic Time!

Can we do better?

Yes? No?

How?

Use different representation.

Another representation.

Represent a line?

Another representation.

Represent a line?

Slope and intercept!

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...and evaluate.

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...and evaluate.

How to find “line” from points?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...and evaluate.

How to find “line” from points?

Solve two variable system of equations!

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...and evaluate.

How to find “line” from points?

Solve two variable system of equations!

How to find polynomial from points?

Another representation.

Represent a line?

Slope and intercept! a_0, a_1

How many points determine a line? 2

Represent line as two points on line instead of coefficients!

How many points determine a parabola (a quadratic polynomial)? 3

How many points determine a a degree d polynomial?

$d + 1$

How to find points on function?

plug in x -values...and **evaluate**.

How to find “line” from points?

Solve two variable system of equations!

How to find polynomial from points?

Solve $d + 1$ variable system of equations!

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Given: a_0, \dots, a_d and b_0, \dots, b_d .

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Given: a_0, \dots, a_d and b_0, \dots, b_d .

Evaluate: $A(x), B(x)$ on $2d + 1$ points: x_0, \dots, x_{2d} .

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Given: a_0, \dots, a_d and b_0, \dots, b_d .

Evaluate: $A(x), B(x)$ on $2d + 1$ points: x_0, \dots, x_{2d} .

Recall(from CS70): unique representation of polynomial.

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Given: a_0, \dots, a_d and b_0, \dots, b_d .

Evaluate: $A(x), B(x)$ on $2d + 1$ points: x_0, \dots, x_{2d} .

Recall(from CS70): unique representation of polynomial.

Multiply: $A(x)B(x)$ on points to get points for $C(x)$.

Point-value representation.

$A(x_0), \dots, A(x_{2d})$

$B(x_0), \dots, B(x_{2d})$

Product: $C(x_0), \dots, C(x_{2d})$

$$C(x_i) = A(x_i)B(x_i)$$

$O(d)$ multiplications!

Given: a_0, \dots, a_d and b_0, \dots, b_d .

Evaluate: $A(x), B(x)$ on $2d + 1$ points: x_0, \dots, x_{2d} .

Recall(from CS70): unique representation of polynomial.

Multiply: $A(x)B(x)$ on points to get points for $C(x)$.

Interpolate: find $c_0 + c_1x + c_2x^2 + \dots c_{2d}x^{2d}$.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13,$

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$,

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

n multiplications/additions to evaluate one point.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

n multiplications/additions to evaluate one point.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

n multiplications/additions to evaluate one point.

Evaluate on n points

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

n multiplications/additions to evaluate one point.

Evaluate on n points ——— $O(n^2)$ time.

Polynomial Evaluation.

Evaluate $A(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ on n points: x_0, \cdots, x_{n-1} .

On one point at a time:

Example: $4 + 3x + 5x^2 + 4x^3$ on 2.

Horners Rule: $4 + x(3 + x(5 + 4x))$

$5 + 4x = 13$, then $3 + 2(13) = 29$, then $4 + 2(29) = 62$.

In general: $a_0 + x(a_1 + x(a_2 + x(\dots)))$.

n multiplications/additions to evaluate one point.

Evaluate on n points ——— $O(n^2)$ time.

Could have just multiplied polynomials!

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2....$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$A(x) = 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \end{aligned}$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

$$A_e(x) = 4 + 20x + 6x^2$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

$$A_e(x) = 4 + 20x + 6x^2$$

$$A_o(x) = 12 + 13x + 7x^2$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

$$A_e(x) = 4 + 20x + 6x^2$$

$$A_o(x) = 12 + 13x + 7x^2$$

$$A(x) = A_e(x^2) + xA_o(x^2)$$

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

$$A_e(x) = 4 + 20x + 6x^2$$

$$A_o(x) = 12 + 13x + 7x^2$$

$$A(x) = A_e(x^2) + xA_o(x^2)$$

Plug in x^2 into A_e and A_o

Evaluation of polynomials: Recursive.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Example:

$$\begin{aligned} A(x) &= 4 + 12x + 20x^2 + 13x^3 + 6x^4 + 7x^5 \\ &= (4 + 20x^2 + 6x^4) + (12x + 13x^3 + 7x^5) \\ &= (4 + 20x^2 + 6x^4) + x(12 + 13x^2 + 7x^4) \end{aligned}$$

$$A_e(x) = 4 + 20x + 6x^2$$

$$A_o(x) = 12 + 13x + 7x^2$$

$$A(x) = A_e(x^2) + xA_o(x^2)$$

Plug in x^2 into A_e and A_o use results to find $A(x)$.

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1$$

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1 = O(n).$$

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1 = O(n).$$

$O(n)$ for 1 point!

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1 = O(n).$$

$O(n)$ for 1 point!

n points – $O(n^2)$ time to evaluate on n points.

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1 = O(n).$$

$O(n)$ for 1 point!

n points – $O(n^2)$ time to evaluate on n points.

No better than polynomial multiplication!

Recursive Evaluation.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

where

Even coefficient polynomial.

$$A_e(x) = a_0 + a_2x + a_4x^2 \dots$$

Odd coefficient polynomial.

$$A_o(x) = a_1 + a_3x + a_5x^2 \dots$$

Evaluate recursively:

For a point x :

Compute $A_e(x^2)$ and $A_o(x^2)$.

$$T(n) = 2T(n/2) + 1 = O(n).$$

$O(n)$ for 1 point!

n points – $O(n^2)$ time to evaluate on n points.

No better than polynomial multiplication! **Bummer.**

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating 2 $\frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating $2 \frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n)$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating $2 \frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n) = O(n \log n)$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating $2 \frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n) = O(n \log n) !!!!$$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating $2 \frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n) = O(n \log n) !!!!$$

From $O(n^2)$ to $O(n \log n)$

Recursive on more than one point.

$$A(x) = A_e(x^2) + x(A_o(x^2))$$

Reuse computations.

n points: $\pm x_0, \pm x_1, \dots, \pm x_{(n-1)/2}$. Also $n = d + 1$: number of coefficients.

Two points: $+x_0$ and $-x_0$ One square: $(+x_0)^2 = (-x_0)^2 = x_0^2$.

$$A(x_0) = A_e(x_0^2) + x_0 A_o(x_0^2)$$

$$A(-x_0) = A_e((-x_0)^2) + (-x_0) A_o((-x_0)^2)$$

$$A(-x_0) = A_e(x_0^2) - x_0 A_o(x_0^2)$$

From $A_e(x_0^2)$ and $A_o(x_0^2)$ compute both $A(-x_0)$ and $A(x_0)$?

From $A_e(x_i^2)$ and $A_o(x_i^2)$ compute both $A(-x_i)$ and $A(x_i)$?

Evaluate n coefficient polynomial on n points by

Evaluating $2 \frac{n}{2}$ coefficient polynomials on $\frac{n}{2}$ points.

$$T(n) = 2T\left(\frac{n}{2}\right) + O(n) = O(n \log n) !!!!$$

From $O(n^2)$ to $O(n \log n)$!!!

Explore recursion.

Recursive Condition:

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,
 $\pm x_1$ both have x_1^2 as square.

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

But all our $\frac{n}{2}$ points are squares

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

But all our $\frac{n}{2}$ points are squares ...and positive!

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

But all our $\frac{n}{2}$ points are squares ...and positive!

Need the $\frac{n}{2}$ points to come in Positive/Negatives pairs!

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

But all our $\frac{n}{2}$ points are squares ...and positive!

Need the $\frac{n}{2}$ points to come in Positive/Negatives pairs!

How can squares be negative?

Explore recursion.

Recursive Condition:

n points: $\frac{n}{2}$ pairs of distinct numbers with common squares.

E.g., $\pm x_0$ both have x_0^2 as square,

$\pm x_1$ both have x_1^2 as square.

.....

Next step:

$\frac{n}{2}$ points: squares should only be $n/4$ distinct numbers

But all our $\frac{n}{2}$ points are squares ...and positive!

Need the $\frac{n}{2}$ points to come in Positive/Negatives pairs!

How can squares be negative?

Complex numbers!

Pairs with common squares.

Want n numbers:

Pairs with common squares.

Want n numbers:

$$x_0, \dots, x_{n-1}$$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where
 $|\{x_0^2, \dots, x_{n-1}^2\}|$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where
 $|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}|$$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}|$$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: $1, -1$.

Take square roots: $1, -1, i, -i$.

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Take square roots: 1, -1 , i , $-i$.

Uh oh.

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Take square roots: 1, -1 , i , $-i$.

Uh oh.

Actually: $\pm 1, \pm i, \pm \frac{1}{\sqrt{2}}(1 + i), \pm \frac{1}{\sqrt{2}}(-1 + i),$

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Take square roots: 1, -1 , i , $-i$.

Uh oh.

Actually: $\pm 1, \pm i, \pm \frac{1}{\sqrt{2}}(1 + i), \pm \frac{1}{\sqrt{2}}(-1 + i)$,

Complex numbers!

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Take square roots: 1, -1 , i , $-i$.

Uh oh.

Actually: $\pm 1, \pm i, \pm \frac{1}{\sqrt{2}}(1 + i), \pm \frac{1}{\sqrt{2}}(-1 + i)$,

Complex numbers!

Uh oh?

Pairs with common squares.

Want n numbers:

x_0, \dots, x_{n-1} where

$$|\{x_0^2, \dots, x_{n-1}^2\}| = \frac{n}{2},$$

and

$$|\{x_0^4, \dots, x_{n-1}^4\}| = \frac{n}{4},$$

...and ...

$$|\{x_0^{\log n}, \dots, x_{n-1}^{\log n}\}| = 1.$$

Each recursive level evaluates:

polynomials of half the degree on half as many points.

n represents both degree and number of points.

In reverse: start with a number 1

Take square roots: 1, -1 .

Take square roots: 1, -1 , i , $-i$.

Uh oh.


Actually: $\pm 1, \pm i, \pm \frac{1}{\sqrt{2}}(1 + i), \pm \frac{1}{\sqrt{2}}(-1 + i)$,

Complex numbers!


Uh oh? Can we get a pattern?

Complex plane


$$z = a + bi$$



Complex plane



Complex plane




$$z = a + bi$$

$$r = \sqrt{a^2 + b^2}$$

$$\theta = \tan^{-1}\left(\frac{b}{a}\right)$$

Complex plane




$$z = a + bi$$

$$r = \sqrt{a^2 + b^2}$$


$$\theta = \tan^{-1}\left(\frac{b}{a}\right)$$

Polar coordinate: $r(\cos \theta + i \sin \theta) = re^{i\theta}$ or (r, θ)


Multiplying Complex Numbers




Multiplying Complex Numbers



Multiplying Complex Numbers




Multiplying Complex Numbers




$$\begin{aligned}(r_1, \theta_1) \times (r_2, \theta_2) \\&= r_1 e^{i\theta_1} \times r_2 e^{i\theta_2} \\&= (r_1 r_2) e^{i(\theta_1 + \theta_2)}\end{aligned}$$

Multiplying Complex Numbers




$$\begin{aligned}(r_1, \theta_1) \times (r_2, \theta_2) &= r_1 e^{i\theta_1} \times r_2 e^{i\theta_2} \\ &= (r_1 r_2) e^{i(\theta_1 + \theta_2)} \\ &= (r_1 r_2, \theta_1 + \theta_2)\end{aligned}$$


Multiplying Complex Numbers



Multiplying Complex Numbers



Multiplying Complex Numbers




$$\begin{aligned}(r_1, \theta_1) \times (r_2, \theta_2) &= r_1 e^{i\theta_1} \times r_2 e^{i\theta_2} \\ &= (r_1 r_2) e^{i(\theta_1 + \theta_2)} \\ &= (r_1 r_2, \theta_1 + \theta_2)\end{aligned}$$

$$\begin{aligned}-1 \times (r_1, \theta_1) &= (r_1, \theta_1 + \pi) \\ &= r_1 e^{r_1(\theta_1 + \pi)i}\end{aligned}$$

The n th complex roots of unity.


Solutions to $z^n = 1$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2\pi}{n}\right)^n = \left(1, \frac{2\pi}{n} \times n\right) = (1, 2\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2\pi}{n}\right)^n = \left(1, \frac{2\pi}{n} \times n\right) = (1, 2\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{4\pi}{n}\right)^n = \left(1, \frac{4\pi}{n} \times n\right) = (1, 4\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2k\pi}{n}\right)^n = \left(1, \frac{2k\pi}{n} \times n\right) = (1, 2k\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



The n th complex roots of unity.

Solutions to $z^n = 1$

$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



Squares: $\frac{n}{2}$ th roots.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}}$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}} = \omega^j \omega^{\frac{n}{2}}$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}} = \omega^j \omega^{\frac{n}{2}} = -\omega^j$.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}} = \omega^j \omega^{\frac{n}{2}} = -\omega^j$. Common square!

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \cdots a_{n-1}x^{n-1}$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \cdots a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \cdots a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}} = \omega^j \omega^{\frac{n}{2}} = -\omega^j$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+\frac{n}{2}} = \omega^j \omega^{\frac{n}{2}} = -\omega^j$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:

$\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+\frac{n}{2}}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+\frac{n}{2}}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+\frac{n}{2}}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

A_e and A_o are degree $\frac{n}{2}$, $\frac{n}{2}$ points in recursion.

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+\frac{n}{2}}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

A_e and A_o are degree $\frac{n}{2}$, $\frac{n}{2}$ points in recursion.

$$T(n) = 2T(\frac{n}{2}) + O(n)$$

The FFT!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+\frac{n}{2}} = \omega^i \omega^{\frac{n}{2}} = -\omega^i$. Common square!

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on points $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq \frac{n}{2}$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+\frac{n}{2}}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

A_e and A_o are degree $\frac{n}{2}$, $\frac{n}{2}$ points in recursion.

$$T(n) = 2T(\frac{n}{2}) + O(n) = O(n \log n)!$$

More FFT ..

...on Friday..