

CS 170: Algorithms

S

CS 170: Algorithms

S H

CS 170: Algorithms

S
H
H

CS 170: Algorithms

S
H H
H

CS 170: Algorithms

S H H H

CS 170: Algorithms

S H H H
H H
H
.

CS 170: Algorithms

S H H H
H H
H
.
.

CS 170: Algorithms

S H H H H . .


CS 170: Algorithms

S H H H .
.
.

CS 170: Algorithms


S H H H .
H H .
H .
.
.

CS 170: Algorithms



No laptops please.


CS 170: Algorithms



No laptops please.

Thank you


CS 170: Algorithms



No laptops please.

Thank you !


CS 170: Algorithms



No laptops please.

Thank you ! !


CS 170: Algorithms



No laptops please.

Thank you ! ! !


CS 170: Algorithms



No laptops please.

Thank you ! ! ! !


CS 170: Algorithms



No laptops please.

Thank you ! ! ! ! !


CS 170: Algorithms



No laptops please.

Thank you ! ! ! ! !


CS 170: Algorithms



No laptops please.

Thank you ! ! ! ! ! ! !

CS 170: Algorithms



No laptops please.

Thank you ! ! ! ! ! ! ! !

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

At time i :

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

At time i :

$$b_0 a_i + b_1 a_{i-1} + \dots + b_i a_0$$

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

At time i :

$$b_0 a_i + b_1 a_{i-1} + \dots + b_i a_0$$

Response: c_0, \dots, c_{2n-1}

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

At time i :

$$b_0 a_i + b_1 a_{i-1} + \dots + b_i a_0$$

Response: c_0, \dots, c_{2n-1}

$$c_i = \sum_j b_j a_{i-j}$$

Convolution = polynomial multiplication.

System: a_0, \dots, a_{n-1}

Signal: b_0, \dots, b_{n-1}

At time 0:

$$b_0 a_0$$

At time 1:

$$b_0 a_1 + b_1 a_0$$

At time i :

$$b_0 a_i + b_1 a_{i-1} + \dots + b_i a_0$$


Response: c_0, \dots, c_{2n-1}

$$c_i = \sum_j b_j a_{i-j}$$

Same as multiplying polynomials $A(\cdot)$ and $B(\cdot)$!

The n th complex roots of unity.


Solutions to $z^n = 1$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2\pi}{n}\right)^n = \left(1, \frac{2\pi}{n} \times n\right) = (1, 2\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2\pi}{n}\right)^n = \left(1, \frac{2\pi}{n} \times n\right) = (1, 2\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{4\pi}{n}\right)^n = \left(1, \frac{4\pi}{n} \times n\right) = (1, 4\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$\left(1, \frac{2k\pi}{n}\right)^n = \left(1, \frac{2k\pi}{n} \times n\right) = (1, 2k\pi) = 1!$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



The n th complex roots of unity.

Solutions to $z^n = 1$


$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



The n th complex roots of unity.

Solutions to $z^n = 1$

$$(1, \theta + \pi)^2 = (1, 2\theta + 2\pi) = \boxed{(1, 2\theta)} = (1, \theta)^2.$$



Squares: $\frac{n}{2}$ th roots.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2}$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2} = \omega^j \omega^{n/2}$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2} = \omega^j \omega^{n/2} = -\omega^j$.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2} = \omega^j \omega^{n/2} = -\omega^j$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \cdots a_{n-1}x^{n-1}$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2} = \omega^j \omega^{n/2} = -\omega^j$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \cdots a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^j and $\omega^{j+n/2} = \omega^j \omega^{n/2} = -\omega^j$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+n/2}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+n/2}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+n/2}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

$$T(n) = 2T(n/2) + O(n)$$

The Fast Fourier Transform!

Defn: $\omega = (1, \frac{2\pi}{n}) = e^{\frac{2\pi}{n}i}$, n th root of unity.

Pairs: ω^i and $\omega^{i+n/2} = \omega^i \omega^{n/2} = -\omega^i$. Common square.

Common Squares: are $\frac{n}{2}$ root of unity.

Fast Fourier Transform:

Evaluate $A(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1}$
on $\omega^0, \omega, \omega^2, \dots, \omega^{n-1}$.

Procedure:

Recursively compute A_e and A_o on $\frac{n}{2}$ roots of unity:
 $\omega^2, \omega^4, \omega^6, \dots, \omega^n$.

For each $i \leq n/2$.

$$A(\omega^i) = A_e(\omega^{2i}) + \omega^i A_o(\omega^{2i})$$

$$A(\omega^{i+n/2}) = A_e(\omega^{2i}) - \omega^i A_o(\omega^{2i})$$

Runtime Recurrence:

$$T(n) = 2T(n/2) + O(n) = O(n \log n)!$$

Multiplying polynomials?

Coefficient representation:


$$a_0, a_1, \dots, a_{n-1}$$

Value representation:

$$A(x_0), A(x_1), \dots, A(x_{n-1}).$$


Multiplying polynomials?

Evaluation: $O(n \log n)$ if choose $1, \omega, \omega^2, \dots, \omega^{n-1}$.



Multiplying polynomials?


Evaluation: $O(n \log n)$ if choose $1, \omega, \omega^2, \dots, \omega^{n-1}$.



Interpolation: From points $A(x_0), \dots, A(x_{n-1})$ to “function”.

Multiplying polynomials?

Evaluation: $O(n \log n)$ if choose $1, \omega, \omega^2, \dots, \omega^{n-1}$.



Interpolation: From points $A(x_0), \dots, A(x_{n-1})$ to “function”.
How?

Polynomial Evaluation and Matrices

Compute $A(\cdot)$ from a_i 's:

$$\begin{bmatrix} A(x_0) \\ A(x_1) \\ \vdots \\ A(x_{n-1}) \end{bmatrix} = \begin{bmatrix} 1 & x_0 & x_0^2 & \cdots x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots x_1^{n-1} \\ & & \vdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots x_{n-1}^{n-1} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Compute inverse of matrix above.

Polynomial Evaluation and Matrices

Compute $A(\cdot)$ from a_i 's:

$$\begin{bmatrix} A(x_0) \\ A(x_1) \\ \vdots \\ A(x_{n-1}) \end{bmatrix} = \begin{bmatrix} 1 & x_0 & x_0^2 & \cdots x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots x_1^{n-1} \\ & & \vdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots x_{n-1}^{n-1} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Compute inverse of matrix above.
Multiply.

Polynomial Evaluation and Matrices

Compute $A(\cdot)$ from a_i 's:

$$\begin{bmatrix} A(x_0) \\ A(x_1) \\ \vdots \\ A(x_{n-1}) \end{bmatrix} = \begin{bmatrix} 1 & x_0 & x_0^2 & \cdots x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots x_1^{n-1} \\ & & \vdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots x_{n-1}^{n-1} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Compute inverse of matrix above.

Multiply. $O(n^2)$!

Polynomial Evaluation and Matrices

Compute $A(\cdot)$ from a_i 's:

$$\begin{bmatrix} A(x_0) \\ A(x_1) \\ \vdots \\ A(x_{n-1}) \end{bmatrix} = \begin{bmatrix} 1 & x_0 & x_0^2 & \cdots x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots x_1^{n-1} \\ & & \vdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots x_{n-1}^{n-1} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Compute inverse of matrix above.

Multiply. $O(n^2)$!

Doh!!

Polynomial Evaluation and Matrices

Compute $A(\cdot)$ from a_i 's:

$$\begin{bmatrix} A(x_0) \\ A(x_1) \\ \vdots \\ A(x_{n-1}) \end{bmatrix} = \begin{bmatrix} 1 & x_0 & x_0^2 & \cdots x_0^{n-1} \\ 1 & x_1 & x_1^2 & \cdots x_1^{n-1} \\ & & \vdots & \cdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots x_{n-1}^{n-1} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Compute inverse of matrix above.

Multiply. $O(n^2)$!

Doh!!

Also, computing inverse not even easy.

Getting done?

FFT: ω is complex n th root of unity

Getting done?

FFT: ω is complex n th root of unity
and matrix is ...

$$M_n(\omega) = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{(n-1)} \end{bmatrix}$$

Getting done?

FFT: ω is complex n th root of unity
and matrix is ...

$$M_n(\omega) = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{(n-1)} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \end{bmatrix}$$

Getting done?

FFT: ω is complex n th root of unity
and matrix is ...

$$M_n(\omega) = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{(n-1)} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \\ \vdots & & \vdots & & \\ 1 & \omega^j & \omega^{2j} & \dots & \omega^{j(n-1)} \end{bmatrix}$$

Getting done?

FFT: ω is complex n th root of unity
and matrix is ...

$$M_n(\omega) = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{(n-1)} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \\ \vdots & & \vdots & & \\ 1 & \omega^j & \omega^{2j} & \dots & \omega^{j(n-1)} \\ \vdots & & \vdots & & \\ 1 & \omega^{(n-1)} & \omega^{2(n-1)} & \dots & \omega^{(n-1)(n-1)} \end{bmatrix}$$

Getting done?

FFT: ω is complex n th root of unity
and matrix is ...

$$M_n(\omega) = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{(n-1)} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \\ \vdots & & \vdots & & \\ 1 & \omega^j & \omega^{2j} & \dots & \omega^{j(n-1)} \\ \vdots & & \vdots & & \\ 1 & \omega^{(n-1)} & \omega^{2(n-1)} & \dots & \omega^{(n-1)(n-1)} \end{bmatrix}$$

Compute inverse of $M_n(\omega)$?

Geometry and FFT.

Each row is orthogonal.

Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)


Rigid Rotation (and scaling.)!

Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Rigid Rotation (and scaling.)!




Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Rigid Rotation (and scaling.)!




Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Rigid Rotation (and scaling.)!




Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Rigid Rotation (and scaling.)!




Reverse Rotation is inverse operation.

Geometry and FFT.

Each row is orthogonal.

Multiply by $M_n(\omega)$: project point onto each row (and scaled.)

Rigid Rotation (and scaling.)!



Reverse Rotation is inverse operation.

Scaling: for rotation, axis should have length 1, FFT length n .


Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})?$$

i  j

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline j \\ \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline (i,j) \\ \hline \end{array}$$

Recall: $\omega = e^{2\pi/n}$.

$$C_{ij} = \sum_k \omega^{ik} \omega^{-kj}$$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$C = M_n(\omega) \times M_n(\omega^{-1})?$

Recall: $\omega = e^{2\pi/n}$.

$$C_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)}$$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \end{array} = \begin{array}{|c|} \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \text{ } \\ \hline \end{array}$$

The diagram illustrates the element-wise multiplication of two matrices. The first matrix has a horizontal blue band at row i , labeled i to its left. The second matrix has a vertical blue band at column j , labeled j above it. The result is a matrix with a single blue square at the intersection of row i and column j , labeled (i, j) below it.

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$C = M_n(\omega) \times M_n(\omega^{-1})?$

The diagram shows the multiplication of two matrices. The first matrix has a horizontal blue band labeled i on the left. The second matrix has a vertical blue band labeled j on top. The result is a matrix with a small blue square at the intersection labeled (i,j) .

Recall: $\omega = e^{2\pi/n}$.

$$C_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$C = M_n(\omega) \times M_n(\omega^{-1})?$

The diagram shows the multiplication of two matrices. The first matrix has a horizontal blue band labeled i on the left. The second matrix has a vertical blue band labeled j on top. The result is a matrix with a single blue square at the intersection labeled (i,j) .

Recall: $\omega = e^{2\pi/n}$.

$$C_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1}$$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline j \\ \hline \end{array} = \begin{array}{|c|} \hline (i,j) \\ \hline \end{array}$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array} j = \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array} (i,j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$:

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array}^j = \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array}^{(i,j)}$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array}^j = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array}^j \quad (i,j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} (i,j)$$

The diagram illustrates the element-wise multiplication of two $n \times n$ matrices. The first matrix has a horizontal blue bar at row i , labeled with i to its left. The second matrix has a vertical blue bar at column j , labeled with j above it. The result is a matrix with a single blue square at the intersection of row i and column j , labeled with (i,j) below it.

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$:

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array} \quad j \quad = \quad \begin{array}{|c|} \hline \\ \hline \text{---} \\ \hline \end{array} \quad (i,j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} (i, j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n = (\omega^n)^{(i-j)} = 1^n$

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} (i, j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n = (\omega^n)^{(i-j)} = 1^n \implies c_{ij} = 0$.

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} (i, j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n = (\omega^n)^{(i-j)} = 1^n \implies c_{ij} = 0$.

For C – diagonals are n and the off-diagonals are 0.

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} (i, j)$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n = (\omega^n)^{(i-j)} = 1^n \implies c_{ij} = 0$.

For C – diagonals are n and the off-diagonals are 0.

Divide by n get identity!

Algebraically.

Inversion formula: $(M_n(\omega))^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

$$C = M_n(\omega) \times M_n(\omega^{-1})? \quad i \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \times \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} = \begin{array}{|c|} \hline (i,j) \\ \hline \end{array}$$

Recall: $\omega = e^{2\pi/n}$.

$$c_{ij} = \sum_k \omega^{ik} \omega^{-kj} = \sum_k \omega^{(ik-kj)} = \sum_k \omega^{k(i-j)}$$

If $i \neq j$, let $r = \omega^{(i-j)}$ (note: $r^n = 1$)

$$c_{ij} = 1 + r + r^2 + \dots + r^{n-1} = \frac{1-r^n}{1-r}$$

Case $i = j$: $r = \omega^0 = 1$ and $c_{ij} = n$.

Case $i \neq j$: $r^n = (\omega^{(i-j)})^n = (\omega^n)^{(i-j)} = 1^n \implies c_{ij} = 0$.

For C – diagonals are n and the off-diagonals are 0.

Divide by n get identity!

Inversion formula: $M_n(\omega)^{-1} = \frac{1}{n} M_n(\omega^{-1})$.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}
 $1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}$.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1})$.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1})$.

$\implies O(n \log n)$ time for multiplying degree n polynomials.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}
 $1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega).$

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1}).$

$\implies O(n \log n)$ time for multiplying degree n polynomials.

Fast convolution!

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}
 $1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}$.

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1})$.

$\implies O(n \log n)$ time for multiplying degree n polynomials.

Fast convolution! Digital signal processing.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1})$.

$\implies O(n \log n)$ time for multiplying degree n polynomials.

Fast convolution! Digital signal processing. Many other things.

Computing inverse.

FFT works with points with basic root of unity: ω or ω^{-1}

$$1, \omega^{-1}, \omega^{-2}, \dots, \omega^{-(n-1)}.$$

ω^{-1} is a primitive n th root of unity!

Evaluation: $\text{FFT}(A, \omega)$.

Interpolation: $\frac{1}{n} \text{FFT}(A, \omega^{-1})$.

$\implies O(n \log n)$ time for multiplying degree n polynomials.

Fast convolution! Digital signal processing. Many other things.

Cool!!