

# CS 170: Algorithms

Account forms now or after class!

# CS 170: Algorithms

Account forms now or after class!

Static Course Webpage. ([inst.cs.berkeley.edu/~cs170](http://inst.cs.berkeley.edu/~cs170))

# CS 170: Algorithms

Account forms now or after class!

Static Course Webpage. ([inst.cs.berkeley.edu/~cs170](http://inst.cs.berkeley.edu/~cs170))

Will mostly use piazza. Should have/get invitation soon.

# CS 170: Algorithms

Account forms now or after class!

Static Course Webpage. ([inst.cs.berkeley.edu/~cs170](http://inst.cs.berkeley.edu/~cs170))

Will mostly use piazza. Should have/get invitation soon.

Did you find a scanner, yet?

Today.

Modular arithmetic.

# Today.

Modular arithmetic.

...up to ...

# Today.

Modular arithmetic.

...up to ...

# Arithmetic.

Addition:  $O(n)$



# Arithmetic.

Addition:  $O(n)$

	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1

---

# Arithmetic.

Addition:  $O(n)$

								1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									0

# Arithmetic.

Addition:  $O(n)$

							1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>								8	0

# Arithmetic.

Addition:  $O(n)$

[illegible]

# Arithmetic.

Addition:  $O(n)$

					1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>						4	4	8	0

# Arithmetic.

Addition:  $O(n)$

				0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>					9	4	4	8	0

# Arithmetic.

Addition:  $O(n)$

		0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
			4	6	9	4	4	8	0

# Arithmetic.

Addition:  $O(n)$

	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
		4	4	6	9	4	4	8	0



# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
	0	4	4	6	9	4	4	8	0

# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
1	0	4	4	6	9	4	4	8	0

# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
1	0	4	4	6	9	4	4	8	0

Time:  $O(n)$

# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
1	0	4	4	6	9	4	4	8	0

Time:  $O(n)$

Can we do better?

# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
1	0	4	4	6	9	4	4	8	0

Time:  $O(n)$

Can we do better?

Need to look at the numbers to add them...

# Arithmetic.

Addition:  $O(n)$

1	0	0	0	0	1	1	1	1	
	1	2	3	4	5	6	7	8	9
+	9	2	1	2	3	7	6	9	1
<hr/>									
1	0	4	4	6	9	4	4	8	0

Time:  $O(n)$

Can we do better?

Need to look at the numbers to add them... optimal.

## More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

## More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

	1	2	3	4	5	6	7	8	9
×	9	2	1	2	3	7	6	9	1

---



## More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

	1	2	3	4	5	6	7	8	9
×	9	2	1	2	3	7	6	9	1
<hr/>									
	1	2	3	4	5	6	7	8	9



## More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

[illegible]

## More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

[illegible]

# More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

			1	2	3	4	5	6	7	8	9
		×	9	2	1	2	3	7	6	9	1
<hr/>											
			1	2	3	4	5	6	7	8	9
	9	2	2	2	2	2	2	2	2	1	
	.	.	.	.	.	.	.	.	.	.	.
	.	.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.	.	.

$\updownarrow$   
 $n$

# More Al Khwarizmi's: algorithms.

Addition:  $O(n)$

Multiplication:

			1	2	3	4	5	6	7	8	9
		×	9	2	1	2	3	7	6	9	1
<hr/>											
			1	2	3	4	5	6	7	8	9
	9	2	2	2	2	2	2	2	2	1	
	.	.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.	.	.

$\updownarrow$   
 $n$

Time:  $O(n^2)$

# Multiplication

Multiplication:  $O(n^2)$ .

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?



# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

(a) Yes.

(b) No.

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

(a) Yes.

(b) No.

No.

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

(a) Yes.

(b) No.

No.

What ?!?!?

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

(a) Yes.

(b) No.

No.

What ?!?!

Really!

# Multiplication

Multiplication:  $O(n^2)$ .

Is the best possible?

Every digit in  $x$  must multiply every digit in  $y$  at least once!

$\Theta(n^2)$  such pairs.

Is this the best possible?

(a) Yes.

(b) No.

No.

What !?!?

Really!

Later.



# Exponentiation.

Compute  $x^y$ ?

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

Number of bits is  $\log_2 x^y$

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

Number of bits is  $\log_2 x^y = y \log x$



# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

Number of bits is  $\log_2 x^y = y \log x \leq 2^n \times n$ .

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

Number of bits is  $\log_2 x^y = y \log x \leq 2^n \times n$ .

Does it make sense to do this?

# Exponentiation.

Compute  $x^y$ ?

If  $x$  and  $y$  are  $n$ -bit numbers,  
tightest valid upper bound on the number of bits for  $x^y$ ...

(a)  $2n$

(b)  $n^2$

(c)  $2^n$

(d)  $n2^n$

Number of bits is  $\log_2 x^y = y \log x \leq 2^n \times n$ .

Does it make sense to do this?

Seems better just to keep  $x$  and  $y$  around.

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .



# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!



# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!  $y$  is  $n$  bits so

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!  $y$  is  $n$  bits so  $y - 1 = \Theta(2^n)$ !

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!  $y$  is  $n$  bits so  $y - 1 = \Theta(2^n)$ !

Total is  $O(n^2 2^n)$  time.

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!  $y$  is  $n$  bits so  $y - 1 = \Theta(2^n)$ !

Total is  $O(n^2 2^n)$  time. Output is  $n$ -bits.

# Modular exponentiation.

Compute  $x^y \pmod{z}$ .

$x, y, z$  are  $n$ -bit numbers.

Then the result is an  $n$ -bit number!

Recall: Modular multiplication  $a \cdot b \pmod{z}$ .

Multiply  $a$  and  $b$  to get  $c$ .  $O(n^2)$ .

Divide  $c$  by  $z$ , and output remainder.  $O(n^2)$ .

Addition:  $(a + b) \pmod{z}$  - add  $a, b$  maybe subtract  $z$ .  $O(n)$

Compute  $x^y$ , with  $y - 1$  modular multiplications with  $O(n^2)$

Still exponential!  $y$  is  $n$  bits so  $y - 1 = \Theta(2^n)$ !

Total is  $O(n^2 2^n)$  time. Output is  $n$ -bits.

Can we do better?

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$



## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$213 \pmod{900}$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

$$213^{32} \cdot 213^{32} = 213^{64} \pmod{900}$$

# Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

$$213^{32} \cdot 213^{32} = 213^{64} \pmod{900}$$

Only 6 ( $< 7$ ) modular multiplications to compute the powers.

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

$$213^{32} \cdot 213^{32} = 213^{64} \pmod{900}$$

Only 6 ( $< 7$ ) modular multiplications to compute the powers.

At most 6 more modular multiplications to compute the result.

## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

$$213^{32} \cdot 213^{32} = 213^{64} \pmod{900}$$

Only 6 ( $< 7$ ) modular multiplications to compute the powers.

At most 6 more modular multiplications to compute the result.



## Repeated squaring

Compute  $213^{87} \pmod{900}$ ?

Notice:

$$213^{87} = 213^{1+2+4+16+64} \pmod{900}.$$

$$213^{87} = 213^1 \times 213^2 \times 213^4 \times 213^{16} \times 213^{64} \pmod{900}.$$

87 in binary?

$87 \equiv 1010111$  in binary. 7 bits.

$$213 \pmod{900}$$

$$213 \cdot 213 = 213^2 \pmod{900}$$

$$213^2 \cdot 213^2 = 213^4 \pmod{900}$$

$\vdots$

$$213^{32} \cdot 213^{32} = 213^{64} \pmod{900}$$

Only 6 ( $< 7$ ) modular multiplications to compute the powers.

At most 6 more modular multiplications to compute the result.

$O(\text{Number of bits})$  modular multiplications.

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

# Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

Multiply (at most)  $n$  products according to “set” bit positions in  $y$ .

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

Multiply (at most)  $n$  products according to “set” bit positions in  $y$ .

If  $y$  is  $n$ -bit number,

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

Multiply (at most)  $n$  products according to “set” bit positions in  $y$ .

If  $y$  is  $n$ -bit number,

..  $O(n)$  multiplications of  $n$  bit numbers each in time  $O(n^2)$ .



## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

Multiply (at most)  $n$  products according to “set” bit positions in  $y$ .

If  $y$  is  $n$ -bit number,

..  $O(n)$  multiplications of  $n$  bit numbers each in time  $O(n^2)$ .

Total time is  $O(n^3)$ .

## Repeated squaring.

General:  $x, y, z$ . ( $y$  is nonzero)

Compute  $x^y \pmod{z}$ .

Write  $y$  in binary.

Compute  $x^1, x^2, x^4, \dots, x^{2^n} \pmod{z}$  using  $n$  modular  $\times$ 's.

Multiply (at most)  $n$  products according to “set” bit positions in  $y$ .

If  $y$  is  $n$ -bit number,

..  $O(n)$  multiplications of  $n$  bit numbers each in time  $O(n^2)$ .

Total time is  $O(n^3)$ .

Coding it up.

```
def exp(x,y,z):  
    if (y==1):
```

Coding it up.

```
def exp(x,y,z):  
    if (y==1):  
        return x
```

Coding it up.

```
def exp(x,y,z):  
    if (y==1):  
        return x  
    else:
```

## Coding it up.

$$r = x^{y/2} \pmod{z}$$

```
def exp(x,y,z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x,y/2,z)
```

## Coding it up.

$$r = x^{y/2} \pmod{z}$$

$$v = (x^{y/2})^2 = x^{\lfloor y/2 \rfloor \cdot 2} \pmod{z}$$

```
def exp(x,y,z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x,y/2,z)  
        v = (r*r) %z
```

## Coding it up.

$$r = x^{y/2} \pmod{z}$$

$$v = (x^{y/2})^2 = x^{\lfloor y/2 \rfloor \cdot 2} \pmod{z}$$

Deal with  $\lfloor y/2 \rfloor \cdot 2 \neq y$ !

```
def exp(x, y, z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x, y/2, z)  
        v = (r*r) %z
```



## Coding it up.

$$r = x^{y/2} \pmod{z}$$

$$v = (x^{y/2})^2 = x^{\lfloor y/2 \rfloor \cdot 2} \pmod{z}$$

Deal with  $\lfloor y/2 \rfloor \cdot 2 \neq y$ !

odd  $y$ ,  $y = 1 + \lfloor y/2 \rfloor \cdot 2$ .

$$x^y = x^{1 + \lfloor y/2 \rfloor \cdot 2} = x \cdot x^{\lfloor y/2 \rfloor \cdot 2} = x \cdot v$$

```
def exp(x, y, z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x, y/2, z)  
        v = (r*r) %z  
        if (odd(y)):
```

## Coding it up.

$$r = x^{y/2} \pmod{z}$$

$$v = (x^{y/2})^2 = x^{\lfloor y/2 \rfloor \cdot 2} \pmod{z}$$

Deal with  $\lfloor y/2 \rfloor \cdot 2 \neq y$ !

odd  $y$ ,  $y = 1 + \lfloor y/2 \rfloor \cdot 2$ .

$$x^y = x^{1 + \lfloor y/2 \rfloor \cdot 2} = x \cdot x^{\lfloor y/2 \rfloor \cdot 2} = x \cdot v$$

```
def exp(x, y, z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x, y/2, z)  
        v = (r*r) %z  
        if (odd(y)):  
            return (x * v) %z  
        else:
```

## Coding it up.

$$r = x^{y/2} \pmod{z}$$

$$v = (x^{y/2})^2 = x^{\lfloor y/2 \rfloor \cdot 2} \pmod{z}$$

Deal with  $\lfloor y/2 \rfloor \cdot 2 \neq y$ !

odd  $y$ ,  $y = 1 + \lfloor y/2 \rfloor \cdot 2$ .

$$x^y = x^{1 + \lfloor y/2 \rfloor \cdot 2} = x \cdot x^{\lfloor y/2 \rfloor \cdot 2} = x \cdot v$$

```
def exp(x, y, z):  
    if (y==1):  
        return x  
    else:  
        r = exp(x, y/2, z)  
        v = (r*r) %z  
        if (odd(y)):  
            return (x * v) %z  
        else:  
            return v
```

# Modular Division.

What is division?

# Modular Division.

What is division?

How do I divide by 5?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.



# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11}?$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x?$$



# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{ ? What is } x? \quad 9 \times 5$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45,$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, \quad 45 = 1 \pmod{11}.$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

What is  $y$ ?

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad 9(5y) = 9(7)$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad 9(5y) = 9(7) = 63$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad 9(5y) = 9(7) = 63 = 8 \pmod{11}.$$



# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad y = 9(5y) = 9(7) = 63 = 8 \pmod{11}.$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad y = 9(5y) = 9(7) = 63 = 8 \pmod{11}.$$

$$\text{Check: } 5 \times 8$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad y = 9(5y) = 9(7) = 63 = 8 \pmod{11}.$$

$$\text{Check: } 5 \times 8 = 40$$

# Modular Division.

What is division?

How do I divide by 5?

Multiply by  $1/5$ ?

Multiplicative inverse of 5:

The number you multiply 5 by to get 1.

1 is the multiplicative identity.

Division  $\equiv$  multiply by inverse.

Why divide?

$$5y = 7 \pmod{11}.$$

Find  $y$ ?

Multiply both sides by multiplicative inverse of 5?

$$5x = 1 \pmod{11} \text{? What is } x? \quad 9 \times 5 = 45, 45 = 1 \pmod{11}. \quad x = 9.$$

$$\text{What is } y? \quad y = 9(5y) = 9(7) = 63 = 8 \pmod{11}.$$

$$\text{Check: } 5 \times 8 = 40 = 7 \pmod{11}.$$

# Is there an inverse?

Inverse of 4 (mod 6)?

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.



# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  “ $\equiv$ ”

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .  $x = id, y = jd$  since  $d$  divides both.



# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .  $x = id, y = jd$  since  $d$  divides both.

$a(id) - b(jd) = 1 \rightarrow d(ia - jb) = 1$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .  $x = id, y = jd$  since  $d$  divides both.

$a(id) - b(jd) = 1 \rightarrow d(ia - jb) = 1$ .

$d$  must be a factor of 1.

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .  $x = id, y = jd$  since  $d$  divides both.

$a(id) - b(jd) = 1 \rightarrow d(ia - jb) = 1$ .

$d$  must be a factor of 1. That is,  $d = 1$ .

# Is there an inverse?

Inverse of 4 (mod 6)?

No!

$4j$  is at least 2 away from  $6k$  for any  $j, k$ .

They have a common divisor that is greater than 1.

$\gcd(x, y)$  - greatest common divisor of  $x$  and  $y$ .

$\gcd(x, y) \neq 1$  implies no inverse.

Theorem:

$\gcd(x, y) = d, d \geq 1 \rightarrow x$  has no multiplicative inverse modulo  $y$ .

Proof:  $ax = 1 \pmod{y}$  " $\equiv$ "  $ax = 1 + by$  for integer  $b$

$ax - by = 1$ .  $x = id, y = jd$  since  $d$  divides both.

$a(id) - b(jd) = 1 \rightarrow d(ia - jb) = 1$ .

$d$  must be a factor of 1. That is,  $d = 1$ .



Review: extended euclid's algorithm and inverses.

Extended GCD:

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

Find inverse of  $x$  modulo  $N$ , if  $\gcd(x, N) = 1$ ?



## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

Find inverse of  $x$  modulo  $N$ , if  $\gcd(x, N) = 1$ ?

- (A) Run Euclid on  $x, N$ , output  $a$ .
- (B) Run Euclid on  $x, N$ , output  $b$ .

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

Find inverse of  $x$  modulo  $N$ , if  $\gcd(x, N) = 1$ ?

(A) Run Euclid on  $x, N$ , output  $a$ .

(B) Run Euclid on  $x, N$ , output  $b$ .

A.  $1 = ax + bN$

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

Find inverse of  $x$  modulo  $N$ , if  $\gcd(x, N) = 1$ ?

(A) Run Euclid on  $x, N$ , output  $a$ .

(B) Run Euclid on  $x, N$ , output  $b$ .

A.  $1 = ax + bN = ax \pmod{N}$ ,

## Review: extended euclid's algorithm and inverses.

Extended GCD:

Given  $x, y$ .

Returns:  $(d, a, b)$  where  $ax + by = d$ , and  $d = \gcd(x, y)$

Find inverse of  $x$  modulo  $N$ , if  $\gcd(x, N) = 1$ ?

(A) Run Euclid on  $x, N$ , output  $a$ .

(B) Run Euclid on  $x, N$ , output  $b$ .

A.  $1 = ax + bN = ax \pmod{N}$ ,  
so  $a$  is multiplicative inverse of  $x$  modulo  $N$ .

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ 's:

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

$$x = \underline{\hspace{2cm}}$$

$$y = \underline{\hspace{10cm}}$$

Get “close” to  $y$  with  $x$ 's:

$$kx = \underline{\hspace{10cm}}$$

$$y = \underline{\hspace{10cm}}$$



## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

$$x = \underline{\hspace{2cm}}$$

$$y = \underline{\hspace{10cm}}$$

Get “close” to  $y$  with  $x$ ’s:

$$kx = \underline{\hspace{10cm}}$$

$$y = \underline{\hspace{10cm}}$$

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b); d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ):  $(d, a, b)$ ;  $d = ax + by$ .

$$x = \underline{\hspace{2cm}}$$

$$y = \underline{\hspace{15cm}}$$

Get “close” to  $y$  with  $x$ ’s:

$$kx = \underline{\hspace{15cm}}$$

$$y = \underline{\hspace{15cm}}$$

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides both  $x$  and  $y$ , divides  $(y - kx)$

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides both  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$$x = \underline{\hspace{2cm}}$$

$$y = \underline{\hspace{10cm}}$$

Get “close” to  $y$  with  $x$ ’s:

$$kx = \underline{\hspace{10cm}}$$

$$y = \underline{\hspace{10cm}}$$

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides **both**  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$$d|x \text{ and } d|(y - kx)$$

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$$x = \underline{\hspace{2cm}}$$

$$y = \underline{\hspace{10cm}}$$

Get “close” to  $y$  with  $x$ ’s:

$$kx = \underline{\hspace{10cm}}$$

$$y = \underline{\hspace{10cm}}$$

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides **both**  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx)$

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides **both**  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides **both**  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

$\rightarrow \gcd(x, y) = \gcd(x, y - kx)$ .



## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides both  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

$\rightarrow \gcd(x, y) = \gcd(x, y - kx)$ .

Get  $(d, a', b')$  where  $d = a'(y - kx) + b'x$

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides **both**  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

$\rightarrow \gcd(x, y) = \gcd(x, y - kx)$ .

Get  $(d, a', b')$  where  $d = a'(y - kx) + b'x = (b' - ka')x + a'y$ .

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides both  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

$\rightarrow \gcd(x, y) = \gcd(x, y - kx)$ .

Get  $(d, a', b')$  where  $d = a'(y - kx) + b'x = (b' - ka')x + a'y$ .

Return  $(d, b' - ka', a')$ .

## Extended Euclid/Correctness.

Ext-gcd( $x, y$ ): ( $d, a, b$ );  $d = ax + by$ .

$x =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

Get “close” to  $y$  with  $x$ ’s:

$kx =$  \_\_\_\_\_

$y =$  \_\_\_\_\_

$k = \lfloor y/x \rfloor$  (Use long division.) (Time:  $O(n^2)$  time.)

$(y - kx)$  preserves common divisor!

Anything that divides both  $x$  and  $y$ , divides  $(y - kx)$

Recurse for  $y - kx$  and  $x$

$d|x$  and  $d|(y - kx)$  Also  $d'|x$  and  $d'|(y - kx) \implies d'|y$ .

$\rightarrow \gcd(x, y) = \gcd(x, y - kx)$ .

Get  $(d, a', b')$  where  $d = a'(y - kx) + b'x = (b' - ka')x + a'y$ .

Return  $(d, b' - ka', a')$ .

Time for one recursive call:  $O(n^2)$ .

See you Friday..

....