



## Lecture 2: Exploratory Data Analysis and Summary Statistics

EDA



creative adventures with your data

# Announcements and reminders

---

- **Canvas:** make sure you have looked over the syllabus and schedule  
<https://canvas.colorado.edu/courses/24706>
- **Piazza:** be on it, because no more emails, and I don't like Canvas very much!  
<https://piazza.com/colorado/spring2019/csci3022/>
- Get **Jupyter notebook / Anaconda Python** -- make sure you have a working install and check out the Numpy/Pandas tutorial (github/notebooks)

<https://www.anaconda.com/downloads>



# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**



# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**

**Definition:** A population is a collection of units (people, songs, tweets, marmots)

**Definition:** A sample is a subset of the population

**Definition:** A characteristic/variable of interest (VOI) is something we want to measure for each unit.



# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**

**Example:** S'pose the city of Boulder wants to estimate its per-household income via a phone survey. They call every 50th number on a list of Boulder phone numbers between 6 PM and 8 PM. In this case, we have:

Population: *Boulder residents*

Sample: *every 50th residents*

Variable of Interest: *household income*



# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**

**Example:** S'pose the city of Boulder wants to estimate its per-household income via a phone survey. They call every 50th number on a list of Boulder phone numbers between 6 PM and 8 PM. In this case, we have:

Population: Boulder residents

Sample: every 50th person w/ phone who answers

Variable of Interest: household income

**Definition:** the **sample frame** is the source material or device from which sample is drawn

## Sample types

---

- **Simple random sample:** randomly select people from sample frame
- **Systematic sample:** order the sample frame. Choose integer  $k$ . Sample every  $k$ th unit in the sample frame.
- **Census sample:** sample literally *everyone/everything* in the population
- **Stratified sample:** if you have a heterogeneous population that can be broken up into homogeneous groups, randomly sample from each group proportionate to their prevalence in the population

# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**

So how do we make the jump from studying a sample to drawing meaningful conclusions about the characteristic of the population?





# Populations and samples

---

Data scientists hope to learn about some **characteristic/variable** of a **population**

But, we usually can't actually see/study the whole population → so we study a **sample**

So how do we make the jump from studying a sample to drawing meaningful conclusions about the characteristic of the population?

... **inference!**



# Exploratory data analysis (EDA)

---

Before we learn about **inference** though, we first need to learn how to **explore** the data

Useful for summarizing, recognizing patterns, etc. in the data

There are two main types of data exploration: **numerical** and **graphical**




# Numerical summaries

---

The calculation and interpretation of certain summarizing numbers can help us gain a better understanding of the data

These sample numerical summaries are called **sample statistics**



# Measures of centrality


---

Summarizing the “center” -- or better yet -- “**central tendency**” of the sample data is a popular and important characteristic of a set of numbers

**Goal:** capture something about the “typical” unit in the sample with respect to the VOI

3 main measures:

- 1) Mean
- 2) Median
- 3) Mode



## Sample mean

---

**Definition:** For a given set of numbers  $x_1, x_2, \dots, x_n$ , the sample mean is given by:

Also called the **arithmetic average**

**Example:** Compute the sample mean of the data 2, 4, 3, 5, 6, 4

$$\frac{24}{6} = 4$$

## Sample mean

---

**Definition:** For a given set of numbers  $x_1, x_2, \dots, x_n$ , the **sample mean** is given by:

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k$$

Also called the **arithmetic average**

**Example:** Compute the sample mean of the data 2, 4, 3, 5, 6, 4

$$\begin{aligned}\bar{x} &= \frac{1}{6}(2 + 4 + 3 + 5 + 6 + 4) \\ &= \frac{1}{6} \cdot 24 \\ &= 4\end{aligned}$$

# Sample mean

---

**Definition:** For a given set of numbers  $x_1, x_2, \dots, x_n$ , the **sample mean** is given by:

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k$$

Also called the **arithmetic average**

**Advantages:**

**Disadvantages:**

# Sample mean

---

**Definition:** For a given set of numbers  $x_1, x_2, \dots, x_n$ , the **sample mean** is given by:

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k$$

Also called the **arithmetic average**

**Advantages:** easy to calculate

**Disadvantages:** outliers can make interpretation misleading



# Sample median

---

**Definition:** For a given set of numbers the sample median is the “middle” value when the observations are ordered from smallest to largest.

## Calculation:

- Order the  $n$  observations from smallest to largest
- Include multiple instances of repeated values
- If  $n$  is odd, then  $\tilde{x} = \left(\frac{n+1}{2}\right)^{th}$  ordered value
- If  $n$  is even, then  $\tilde{x} =$  average of  $\left(\frac{n}{2}\right)^{th}$  and  $\left(\frac{n}{2} + 1\right)^{th}$  ordered values

## Sample median

---

**Definition:** For a given set of numbers the sample median is the “middle” value when the observations are ordered from smallest to largest.

**Example:** Calculate the sample median of the data 36, 15, 39, 41, 40, 42, 47, 49, 7, 6

6, 7, 15, 36, 39, 40, 41, 42, 47, 49

$$39 + 40 = 79 \begin{array}{r|l} 2 & \\ \hline 19 & 39.5 \\ \hline 50 & \\ 0 & \end{array}$$

## Sample median

---

**Definition:** For a given set of numbers the sample median is the “middle” value when the observations are ordered from smallest to largest.

**Example:** Calculate the sample median of the data 36, 15, 39, 41, 40, 42, 47, 49, 7, 6

**Solution:**  $n = 10$  is even so it's the average of the middle 2 numbers when sorted:


6, 7, 15, 36, **39, 40**, 41, 42, 47, 49

→ **39.5**

# Sample mode

---

**Definition:** The sample mode is the value that occurs the most often in the sample.




## Mean vs median

The population mean and median will generally not be equal.


If the population distribution is positively or negatively skewed ...



negative skew  
(left-skew)



symmetric



positive skew  
(right skew)


## Mean vs median

The population mean and median will generally not be equal.


If the population distribution is positively or negatively skewed ...



negative skew  
(left-skew)



symmetric



positive skew  
(right skew)

→ Which measure of central tendency is most important?

## Other sample measures

---

**Quartiles:** Divide the data into 4 equal parts

- Lower quartile ( $Q_1$  or  $P_{25}$ ) splits the lowest 25% of the data from the other 75%
- Middle quartile ( $Q_2$  or  $P_{50}$ ) splits the data in half (i.e., the **median**)
- Upper quartile ( $Q_3$  or  $P_{75}$ ) splits the highest 25% of the data from the lowest 75%

## Other sample measures

---

**Quartiles:** Divide the data into 4 equal parts

- Lower quartile ( $Q_1$  or  $P_{25}$ ) splits the lowest 25% of the data from the other 75%
- Middle quartile ( $Q_2$  or  $P_{50}$ ) splits the data in half (i.e., the **median**)
- Upper quartile ( $Q_3$  or  $P_{75}$ ) splits the highest 25% of the data from the lowest 75%

**Computation:**

- 1) Use the median to divide the ordered data set into 2 halves
  - If  $n$  is odd, include the median in both halves
  - If  $n$  is even, split the data exactly in half
- 2) The lower quartile is the median of the lower half
- 3) The upper quartile is the median of the upper half



## Other sample measures

---

**Quartiles:** Divide the data into 4 equal parts

- Lower quartile ( $Q_1$  or  $P_{25}$ ) splits the lowest 25% of the data from the other 75%
- Middle quartile ( $Q_2$  or  $P_{50}$ ) splits the data in half (i.e., the **median**)
- Upper quartile ( $Q_3$  or  $P_{75}$ ) splits the highest 25% of the data from the lowest 75%

**Example:** Compute the quartiles of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

$$Q_2 = 40$$

$$Q_1 = \frac{51}{2} = 25.5$$

$$Q_3 = \frac{85}{2} = 42.5$$

## Other sample measures

---

**Quartiles:** Divide the data into 4 equal parts

- Lower quartile ( $Q_1$  or  $P_{25}$ ) splits the lowest 25% of the data from the other 75%
- Middle quartile ( $Q_2$  or  $P_{50}$ ) splits the data in half (i.e., the **median**)
- Upper quartile ( $Q_3$  or  $P_{75}$ ) splits the highest 25% of the data from the lowest 75%

**Example:** Compute the quartiles of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

**Solution:**

- 1) Data are already sorted
- 2) Compute median  $\rightarrow n=11$  is odd, so middle value is median,  $Q_2 = 40$
- 3) Compute  $Q_1$  and  $Q_3$  from first and second halves of data:  
 $Q_1 = \text{median of first half (6, 7, 15, 36, 39, 40)} = (15+36)/2 = 25.5$   
 $Q_3 = \text{median of second half (40, 41, 42, 43, 47, 49)} = (42+43)/2 = 42.5$

## Other sample measures

---

**Quartiles:** Divide the data into 4 equal parts

- Lower quartile ( $Q_1$  or  $P_{25}$ ) splits the lowest 25% of the data from the other 75%
- Middle quartile ( $Q_2$  or  $P_{50}$ ) splits the data in half (i.e., the **median**)
- Upper quartile ( $Q_3$  or  $P_{75}$ ) splits the highest 25% of the data from the lowest 75%

**Percentiles:**

- Generalization of quartiles
- $Q_1$  is the 25th percentile,  $P_{25}$
- Can also calculate general percentiles:


e.g., the 16th percentile ( $P_{16}$ ) splits off the lower 16% of the data.

# Variability

---

So far, we have learned about measuring the **central tendency** of data


But what about the **spread**?



# Variability

So far, we have learned about measuring the **central tendency** of data


But what about the **spread**?



# Variability

The simplest measure of variability is the range

**Definition:** The range of a sample is the difference between the max and min values



# Variability

---

What if we combined the deviations into a single quantity by finding the average deviation?

A more robust measure of variation takes into account deviations from the mean

$$x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x}$$

# Variability

---

What if we combined the deviations into a single quantity by finding the average deviation?

A more robust measure of variation takes into account deviations from the mean

$$x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x}$$

So... what do we do with these things?



# Variability

---

What if we combined the deviations into a single quantity by finding the average deviation?

A more robust measure of variation takes into account deviations from the mean

$$x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x}$$

So... what do we do with these things?

... add them?

$$\frac{1}{n} [(x_1 - \bar{x}) + (x_2 - \bar{x}) + \dots + (x_n - \bar{x})]$$

# Variability

---

What if we combined the deviations into a single quantity by finding the average deviation?

A more robust measure of variation takes into account deviations from the mean

$$x_1 - \bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x}$$

So... what do we do with these things?

... add them?

$$\frac{1}{n} [(x_1 - \bar{x}) + (x_2 - \bar{x}) + \dots + (x_n - \bar{x})]$$

... square, **then** add them?

$$\frac{1}{n} [(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \dots + (x_n - \bar{x})^2]$$

# Variability

---

**Definition:** The **sample variance**, denoted by  $s^2$ , is given by

$$s^2 = \frac{1}{n-1} \sum_{k=1}^n (x_k - \bar{x})^2$$

**Definition:** The **sample standard deviation**, denoted by  $s$ , is given by the (positive) square root of the variance:

$$s = \sqrt{s^2}$$

**NB:**

- The variance and SD are both **nonnegative** ( $\geq 0$ )
- The units for SD are the same as for the data

# Variability

---

**Example:** Compute the SD of the data: 2, 4, 3, 5, 6, 4

# Variability

---

**Example:** Compute the SD of the data: 2, 4, 3, 5, 6, 4

**Solution:**

1) Need  $\bar{x} \dots = (2+4+3+5+6+4) / 6 = 24 / 6 = 4$

2) Calculate  $s^2 \dots$

$$\begin{aligned}s^2 &= \frac{1}{6-1} [(2-4)^2 + (4-4)^2 + (3-4)^2 + (5-4)^2 + (6-4)^2 + (4-4)^2] \\&= \frac{1}{5} [4 + 0 + 1 + 1 + 4 + 0] \\&= \frac{1}{5} \cdot 10 = 2\end{aligned}$$

3)  $s = \sqrt{s^2} = \sqrt{2}$


# Interquartile range

**Definition:** The interquartile range is defined to be the difference between the upper and lower quartiles:

$$\text{IQR} = Q_3 - Q_1$$

→ IQR gives the spread of 50% of the data

**Examples:**



## Interquartile range

---

**Example:** Compute the IQR of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

## Interquartile range

---

**Example:** Compute the IQR of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

**Solution:**

- 1) Data are already sorted
- 2) Compute median  $\rightarrow n=11$  is odd, so middle value is median,  $Q_2 = 40$
- 3) Compute  $Q_1$  and  $Q_3$  from first and second halves of data:

$$Q_1 = \text{median of first half (6, 7, 15, 36, 39, 40)} = (15+36)/2 = 25.5$$

$$Q_3 = \text{median of second half (40, 41, 42, 43, 47, 49)} = (42+43)/2 = 42.5$$

- 4)  $IQR = Q_3 - Q_1 = 42.5 - 25.5 = 17$



## Tukey 5-number summary

---

John Tukey, father of modern EDA, advocated summarizing data sets with 5 values:

- 1) Min value
- 2) Lower quartile
- 3) Median
- 4) Upper quartile
- 5) Max value

**Example:** Find the 5-number summary of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

# Tukey 5-number summary

---

John Tukey, father of modern EDA, advocated summarizing data sets with 5 values:

- 1) Min value
- 2) Lower quartile
- 3) Median
- 4) Upper quartile
- 5) Max value

**Example:** Find the 5-number summary of the data 6, 7, 15, 36, 39, 40, 41, 42, 43, 47, 49

## Advantages:

- gives the center of the data
- gives the spread of the data (range in IQR)
- gives an idea of skewness

# Tukey 5-number summary

---

## Advantages:

- gives the center of the data
- gives the spread of the data (range in IQR)
- gives an idea of skewness
  - E.g., if  $Q_2$  is closer to  $Q_1$  than to  $Q_3$ , then you know the median is “leaning left” (so, distribution is right-skewed)

## Next time...

- We'll see how to visualize this!  
(histograms and box-whisker plots)

### BOX & WHISKER PLOT



### BOX & BEARD PLOT



