

NetExam

Sri Lanka Institute of Information Technology

Let,
$$A = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix}$$
, $B = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 2 \\ 1 & -2 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 4 \\ 5 & -6 \end{bmatrix}$

Find A+ 2B - 3C + D.

$$Answer = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{matrix} 1 & -3 \\ 0 & 2 \end{matrix}$$



ed

NetExam

Sri Lanka Institute of Information Technology

Let $A = \begin{bmatrix} 8 & 3 \\ 4 & -6 \end{bmatrix}$ A.A = $\begin{bmatrix} 76 & 6 \\ 8 & 48 \end{bmatrix}$

Find $A^2 - 3I + 2A$.

Answer =

89 12

16 33



Sri Lanka Institute of Information Technology

on 9 It arthurs not be been be

ing question

Let $A = \begin{bmatrix} 3 & 7 \\ -4 & 5 \end{bmatrix}$

Find $A^2 - 3I + 2A$.

Answer =

-16 70 -40 4 A.A = -19 56 -32 -3



Sri Lanka Institute of Information Technology

Question 3

Not yet answered

Marked out of 4.00

P Flag question

$$A = \begin{bmatrix} 21 & 13 & -4 \\ 15 & 14 & 0 \end{bmatrix} \cdot \begin{bmatrix} -7 & 11 \\ 4 & -4 \\ -12 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

NetExauni

Sri Lanka Institute of Information Technology

Question 3


Not yet answered

Marked out of 4.00

P Flag question

$$A = \begin{bmatrix} 21 & 13 & -4 \\ 15 & 14 & 0 \end{bmatrix} \cdot \begin{bmatrix} -7 & 11 \\ 4 & -4 \\ -12 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$





NetExam

Sri Lanka Institute of Information Technology


Let,
$$A = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix}$$
, $B = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 2 \\ 1 & -2 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 4 \\ 5 & -6 \end{bmatrix}$

Find A+ 2B - 3C + D.

$$Answer = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Answer
$$- \mid_{c} d$$

$$a = 1$$
 $b = 1$ 1 -3 $c = 1$ $d = 1$ 0 2





Sri Lanka Institute of Information Technology

Question 2

Not yet answered

Marked out of 4.00

P Flag question

Let
$$A = \begin{bmatrix} 3 & 7 \\ -4 & 5 \end{bmatrix}$$

Find
$$A^2 - 3I + 2A$$
.

Moodle

oodle ×

F



NetExam

Sri Lanka Institute of Information Technology

Question 3

Not yet answered Marked out of 4.00

P Hag question

$$A = \begin{bmatrix} 21 & 13 & -4 \\ 15 & 14 & 0 \end{bmatrix} \cdot \begin{bmatrix} -7 & 11 \\ 4 & -4 \\ -12 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$



Sri Lanka Institute of Information Technology

Question 1

Not yet answered

Marked out of 4.00

P Flag question

$$A = \begin{bmatrix} 12 & 10 & -7 \\ 15 & 11 & 0 \end{bmatrix} \cdot \begin{bmatrix} -3 & 10 \\ 7 & -2 \\ -11 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$



Sri Lanka Institute of Information Technology

Question 1

Not yet answered

Marked out of 3.00

P Flag question

Assume the degree sequence of a graph is 5, 4, 3, 2, 2, 1,1. Determine whether, it is possible to draw the graph with the above degree sequence.



ONO

If this graph exists, does it has an Euler Path?

Oyes



XNO

If this graph exists, does it has an Euler Circuit?

Dyes



Question 10


Not yet answered


Marked out of 4.00

Flag question

$$A = \begin{bmatrix} 12 & 10 & -7 \\ 15 & 11 & 0 \end{bmatrix} \cdot \begin{bmatrix} -3 & 10 \\ 7 & -2 \\ -11 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$






Question 1

Not yet answered Marked out of 4.00

P Flag question

$$A = \begin{bmatrix} 12 & 10 & -7 \\ 15 & 11 & 0 \end{bmatrix} \cdot \begin{bmatrix} -3 & 10 \\ 7 & -2 \\ -11 & 8 \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$





Sri Lanka Institute of Information Technology

tion 5

ret answered

red out of


lag question


Let
$$A = \begin{bmatrix} 4 & -3 \\ 6 & 5 \end{bmatrix}$$

Find
$$A^2 - 3I + 2A$$
.

Answer =

3 -33 66 14





ta Institute of Information Technology

Let,
$$A = \begin{bmatrix} 11 & -13 \\ -12 & 21 \end{bmatrix}$$
, $B = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 2 \\ 1 & -2 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 4 \\ 5 & -6 \end{bmatrix}$

Find A+ 2B - 3C + D.


Let,

$$Answer = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

a = : b = : 11 -13 -12 21

c=: d=:

11 -13 -12 21




3

answered

out of

question

Consider the following graph.



Determine whether the above graph has the following.

Hamilton Path =

- O Yes
- ONO

Hamilton Circuit =

- O Yes
- ONO

Euler Path =

- O Yes
- ONO

Euler Circuit =

O Yes

0

Sri Lanka Institute of Information Technology

Question 3

Not yet answered Marked out of

8.00


P Flag question

Let,
$$A = \begin{bmatrix} 11 & -13 \\ -12 & 21 \end{bmatrix}$$
, $B = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 2 \\ 1 & -2 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 4 \\ 5 & -6 \end{bmatrix}$

Find A+ 2B - 3C + D

Let,

$$Answer = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$



Assume the degree sequence of a graph is 6, 4, 4, 2, 2, 2, 0, 0. Question 1 Not yet answered Determine whether, it is possible to draw the graph with the Marked out of above degree sequence. 3.00 P Flag question **Ves** ONO If this graph exists, does it have an Euler Path? O Yes If this graph exists, does it have an Euler Circuit? ONO

ka Institute of Information Technology

ion 1

et answered

ted out of

lag question

$$\operatorname{Let} A = \begin{bmatrix} 2 & 9 \\ 7 & -6 \end{bmatrix}$$

Find
$$A^2 - 3I + 2A$$
.

Answer =

19

A.A = 67

28

-31

99



Sri Lanka Institute of Information Technology

Let,
$$A = \begin{bmatrix} 31 & -15 \\ 13 & 20 \end{bmatrix}$$
, $B = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix}$, $C = \begin{bmatrix} 3 & 2 \\ 1 & -2 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 4 \\ 5 & -6 \end{bmatrix}$

Let,

$$Answer = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$