

Calculus I, II, & III

A Problem-Based Approach with Early Transcendentals

Mahavier < Allen, Browning, Daniel, & So

Lamar University

Contents

To the Student	v
How this Class Works	v
A Common Pitfall	v
Boardwork	vi
How to Study each Day	vi
What is Calculus?	vii
1 Functions, Limits, Velocity, and Tangents	1
1.1 Average Speed and Velocity	1
1.2 Limits and Continuity	3
1.3 Instantaneous Velocity and Growth Rate	7
1.4 Tangent Lines	10
2 Derivatives	11
2.1 Derivatives of Polynomial, Root, and Rational Functions	12
2.2 Derivatives of the Trigonometric Functions	16
2.3 The Chain Rule	17
2.4 Derivatives of the Inverse Trigonometric Functions	19
2.5 Derivatives of the Exponential and Logarithmic Functions	21
2.6 Logarithmic and Implicit Differentiation	23
3 Applications of Derivatives	27
3.1 Linear Approximations	27
3.2 Limits involving Infinity and Asymptotes	28
3.3 Graphing	30
3.4 The Theorems of Calculus	33
3.5 Maxima and Minima	37
3.6 Related Rates	39
4 Integrals	41
4.1 Anti-differentiation and the Indefinite Integral	41
4.2 Riemann Sums and Definite Integrals	43
4.3 The Fundamental Theorem of Calculus	47

4.4	Area, Volume, Arc length and Center of Mass	50
5	Techniques of Integration	56
5.1	Change of Variable Method	56
5.2	Partial Fractions	57
5.3	Integration by Parts	58
5.4	Integrals of Trigonometric Functions	60
5.5	Integration by Trigonometric Substitution	61
5.6	The Hyperbolic Functions	62
5.7	Improper Integrals	63
6	Sequences & Series	65
6.1	Sequences	65
6.2	Series with Positive Constant Terms	68
6.3	Alternating Series	75
6.4	Power Series	79
7	Conic Sections, Parametric Equations, and Polar Coordinates	84
7.1	Conic Sections	84
7.2	Parametric Equations	86
7.3	Polar Coordinates	90
8	Vectors and Lines	93
9	Cross Product and Planes	99
10	Limits and Derivatives	102
11	Optimization and Lagrange Multipliers	113
12	Integration	118
13	Line Integrals, Flux, Divergence, Gauss' and Green's Theorem	127
14	Practice Problems	135
14.1	Functions, Limits, Velocity, and Tangents Drill	135
14.2	Differentiation Drill	139
14.3	Derivative Applications Drill	143
14.4	Integration Drill	148
14.5	Integration Techniques Drill	156
14.6	Sequences and Series Drill	162
14.7	Conic Sections, Parametric Equations, and Polar Coordinates Drill	171
14.8	Vectors, Lines, and Functions Drill	175
14.9	Cross Products, Planes, and Limits Drill	177

14.10	Domains, Graphing, and Derivatives Drill	179
14.11	Optimization and Lagrange Multipliers Drill	184
14.12	Integration Drill	186
14.13	Line Integrals, Flux, Divergence, and Gauss' Theorem Drill	191
15	Appendices	194
15.1	Additional Materials on Limits	194
15.2	The Extended Reals	195
15.3	Exponential and Logarithmic Functions	196
15.4	Trigonometry	200
15.5	Summation Formulas	201

To the Student

“The men who try to do something and fail are infinitely better than those who try to do nothing and succeed.” - Martin Lloyd Jones

How this Class Works

This class will be taught in a way that is likely to be quite different from mathematics classes you have encountered in the past. Much of the class will be devoted to students working problems at the board and much of your grade will be determined by the amount of mathematics that you produce in this class. I use the word produce because it is my belief that the best way to learn mathematics is by doing mathematics.

Therefore, just as I learned to ride a bike by getting on and falling off, I expect that you will learn mathematics by attempting it and occasionally falling off! You will have a set of notes, provided by me, that you will turn into a book by working through the problems. If you are interested in watching someone else put mathematics at the board, working ten problems like it for homework, and then regurgitating this material on tests, then you are not in the correct class. Still, I urge you to seriously consider the value of becoming an independent thinker who tackles doing mathematics, and everything else in life, on your own rather than waiting for someone else to show you how to do things.

A Common Pitfall

There are two ways in which students often approach my classes. The first is to say, “I’ll wait and see how this works and then see if I like it and put some problems up later in the semester after I catch on.” Think of the course as a forty-yard dash. Do you really want to wait and see how fast the other runners are? If you try every night to do the problems then either you will get a problem (Yay!) and be able to put it on the board with pride and satisfaction or you will struggle with the problem, learn a lot in your struggle, and then watch someone else put it on the board. When this person puts it up you will be able to ask questions and help yourself and others understand it, as you say to yourself, “Ahhhh, now I see where I went wrong and now I can do this one and a few more for next class.” If you do not try problems each night, then you will watch the student put the problem on the board, but perhaps will not quite catch all the details and then when you study for the tests or try the next problems you will have only a loose idea of how to tackle such problems. Basically, you have seen it only once in this case. The first student saw it once when s/he tackled it on her/his own, again when either s/he put it on the board or another student presented it, and then a third time when s/he studies for the next test or quiz. Hence the difference between these two approaches is the difference between participating and watching a movie. Movies are filled with successfully married couples and yet something like 60% of marriages end in divorce. Watching successful marriages doesn’t teach one to be successfully married any more than watching mathematics go by

will teach you to do mathematics. I hope that each of you will tackle this course with an attitude that you will learn this material and thus will both enjoy and benefit from the class.

Board Work

Because the board work constitutes a reasonable amount of your grade, let's put your mind at ease regarding this part of the class. First, by coming to class everyday you have a 60% on board work. Every problem you present pushes that grade a little higher. You may come see me any time for an indication of what I think your current level of participation will earn you by the end of the semester for this portion of the grade. Here are some rules and guidelines associated with the board work. I will call for volunteers every day and will pick the person with the least presentations to present a given problem. You may inform me that you have a problem in advance (which I appreciate), but the problem still goes to the person with the least presentations on the day I call for a solution. Ties are broken randomly at the beginning of the course. Once the first test has been returned, ties are broken by giving precedence to the student with the lower test average. A student who has not gone to the board on a given day will be given precedence over a student who has gone to the board that day. To "present" a problem at the board means to have written the problem statement up, to have written a correct solution using complete mathematical sentences, and to have answered all students' questions regarding the problem. Since you will be communicating with other students on a regular basis, here are several guidelines that will help you.

Most importantly, remember that the whole class is on your side and wants to see you succeed, so questions are intended to help everyone, not to criticize you.

When you speak, don't use the words "obvious," "stupid," or "trivial." Don't attack anyone personally or try to intimidate anyone. Don't get mad or upset at anyone. If you do, try to get over it quickly. Don't be upset when you make a mistake – brush it off and learn from it. Don't let anything go on the board that you don't fully understand. Don't say to yourself, "I'll figure this out at home." Don't use concepts we have not defined. Don't use or get examples or solutions from other books. Don't work together without acknowledging it at the board. Don't try to put up a problem you have not written up.

Do prepare arguments in advance. Do be polite and respectful. Do learn from your mistakes. Do ask questions such as, "Can you tell me how you got the third line?" Do let people answer when they are asked a question. Do refer to earlier results and definitions by number when possible.

How to Study each Day

1. Read over your notes from class that day.
2. Make a list of questions to ask me at the beginning of the next class. (I love these!)
3. Review the recent problems.
4. Work on *several* new problems.
5. Write up as many solutions as you can so that you can copy these onto the board the next day.

What is Calculus?

The first semester of Calculus consists of four main concepts: *limits*, *continuity*, *differentiation* and *integration*. Limits are required for defining each of continuity, differentiation, and integration. All four concepts are central to an understanding of applications in fields including biology, business, chemistry, economics, engineering, finance, and physics. The second semester extends your study of *integration techniques* and adds *sequences* and *series*. The third semester of calculus is a repeat of these concepts in higher dimensions with a few other topics tossed in. All of these topics depend on a deep understanding of what might be called *limiting* processes, which is why limits are one of the topics that we will spend a lot of time with in the beginning and will recur throughout the entire course.

In addition to mastering these concepts, I hope to impart in you the essence of the way a mathematician thinks of the world, an axiomatic way of viewing the world. And I hope to help you master the important skill of solving some difficult problems on your own, communicating these solutions to your peers, posing questions respectfully, and responding to questions your peers have. These skills transcend calculus and will help you in all aspects of your life.

Chapter 1

Functions, Limits, Velocity, and Tangents

“The harder I work, the luckier I get.” - Samuel Goldwyn

1.1 Average Speed and Velocity

The **average speed** of an object that has traveled a distance d in time t is $v = d/t$. The terms velocity and speed are interchangeable except that velocity may be negative to indicate direction. For example, if a rock is thrown in the air, we would typically assign a positive velocity on the way up and a negative velocity on the way down.

Problem 1. *A car goes from Houston, Texas to Saint Louis, Missouri (a distance of 600 miles). The trip takes 9 hours and 28 minutes. What was the average speed of the trip? What is one true statement that can you make about the time(s) (if any) at which the car moved at this speed?*

Problem 2. *Robbie the robot is walking down a road in a straight line, away from a pothole. Table 1.1 gives the distance between Robbie and the pothole at different times, where the time is measured in seconds and distance is measured in feet. Write a formula that expresses Robbie’s distance from the pothole as a function of time. Use R to represent distance and t to represent time in your formula. This function will be called Robbie’s position function.*

Table 1.1: Position Table

Time (sec)	Position (feet)
0	2
1	3
2	10
3	29
4	66
5	127

Definition 1. A **relation** is a collection of points in the plane.

Definition 2. A **function** is a collection of points in the plane with the property that no two points have the same first coordinates. The set of all first coordinates of these points is referred to as the **domain** of the function and the set of all second coordinates of these points is referred to as the **range** of the function.

Example 1. Functions may be expressed in many different forms.

The function

$$f = \{(x, y) : y = 6x + 2\} = \{\dots, (-1, -4), (0, 2), (1, 8), (\pi, 6\pi + 2), \dots\}$$

might be expressed via the short-hand,

$$y = 6x + 2 \quad \text{or} \quad f(x) = 6x + 2 \quad \text{or} \quad f(z) = 6z + 2.$$

The function is not the written equation, rather it is the set of points in the plane that satisfy the equation. While every function is a relation, every relation is not necessarily a function. Examples of relations that are not functions would be the vertical line,

$$\{(x, y) : x = 6\} \quad \text{written in short-hand as } x = 6,$$

and the circle,

$$\{(x, y) : x^2 + y^2 = 9\} \quad \text{written in short-hand as } x^2 + y^2 = 9.$$

Problem 3. Express Robbie's position function in the three different forms just discussed. Graph Robbie's position function, labeling the axes and several points.

Interval and Set Notation. We use $\mathbb{R}$ to denote the set of all real numbers and $x \in S$ to mean that x is a member of the set S . If $a \in \mathbb{R}$ and $b \in \mathbb{R}$ and $a < b$ then the **open interval** (a, b) denotes the set of all numbers between a and b . For example,

1. $(a, b) = \{x \in \mathbb{R} : a < x < b\}$ – an open interval
2. $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ – a closed interval
3. $(a, b] = \{x \in \mathbb{R} : a < x \leq b\}$ – a half-open interval

When convenient, we'll use the notation $(-\infty, b)$ to mean all numbers less than b , even though $-\infty$ is not a real number. We won't define ∞ or $-\infty$ in this class, although you can read a bit about them in Appendix 15.2.

Problem 4. Sketch the graph of each of the following and state the domain and range of each.

1. $f(x) = \frac{1}{x-1}$
2. $g(x) = \sqrt{4-x}$
3. $x = y^2 + 1$
4. $i(x) = -(x-2)^3 + 4$

The next several problems deal with Robbie's motion where $R(t)$ represents Robbie's distance from the pothole in feet at time t in seconds.

Problem 5. Assume c is a positive number and compute $R(3)$ and $R(c)$. Write a sentence that says what $R(3)$ means. What does $R(c)$ mean?

Problem 6. Compute the value of $R(5) - R(3)$. What is the physical (real world, useful) meaning of this number? Assume u and v are positive numbers and compute $R(u) - R(v)$. What is the physical meaning of $R(u) - R(v)$?

Problem 7. Evaluate each of the following ratios and explain in detail their physical meaning. Assume that u and v are positive numbers and $u > v$.

1. $\frac{R(5) - R(3)}{5 - 3}$
2. $\frac{R(u) - R(v)}{u - v}$

Problem 8. Assuming that the function you found is valid for all times $t > 4$, where is Robbie when the time is $t = 5$ seconds? $t = 6$ seconds? $t = 5.5$ seconds? When is Robbie 106 feet from the pothole?

Problem 9. Recall from Problem 1 that the distance from Houston, Texas to Saint Louis, Missouri is 600 miles. A car goes from Houston to Saint Louis traveling at 65 mph for the first 22 minutes, accelerates to 95 mph, and remains at that speed for the rest of the trip. Estimate the average speed of the trip. Explain whether your answer is exact or an approximation.

1.2 Limits and Continuity

Definition 3. If f is a function and each of a and L are numbers, then we say that the **limit of f as x approaches a is L** if the values for $f(x)$ get close to the number L as the values for x get close to the number a . We write this as

$$\lim_{x \rightarrow a} f(x) = L \quad \text{or} \quad f(x) \rightarrow L \text{ as } x \rightarrow a.$$

Example 2. What number is the function $f(x) = \frac{x^5 - 32}{x - 2}$, approaching as x approaches 2?

Table 1.2: First Limit Table

x	$f(x)$
1.9	72.39
1.99	79.20
1.999	79.92
1.9999	79.992
2.1	88.41
2.01	80.80
2.001	80.08
2.0001	80.008

Table 1.2 shows that as x approaches 2 from either the left or the right, $f(x)$ approaches 80. From this table, we assume that the limit as x approaches 2 of the function $f(x) = \frac{x^5 - 32}{x - 2}$ is 80. Our notation for this will be:

$$\lim_{x \rightarrow 2} \frac{x^5 - 32}{x - 2} = 80.$$

This is an intuitive approach to the notion of a limit because we did not give specific meaning to the phrases “ x approaches 2” or “ $f(x)$ approaches 80.” To give a mathematically accurate definition for the word “limit” took mathematicians almost 200 years, yet our intuitive approach will serve us well for this course. If you are interested in a more precise definition of the limit, read Appendix 15.1 or take *Real Analysis*.

Problem 10. Compute $\lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2}$ by constructing a limit table like Table 1.2. What is $\lim_{x \rightarrow -2} \frac{x^3 - 8}{x - 2}$?

Problem 11. Arguments for trigonometric functions in calculus are in radians unless stated otherwise, so be sure your calculator is in radian mode. Compute $\lim_{x \rightarrow 0} \frac{\sin(x)}{2x}$ by constructing a limit table.

Example 3. Computing limits using algebra.

A reasonable attempt to find the limit of the function f as x approaches a would be to simply plug a in for x in the function. As illustrated in the last two problems, *this does not always work*. There is an important distinction between the value $f(a)$ and the limit of f as x approaches a . In addition to tables and graphing, we may also use algebra to evaluate some limits.

$$\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3} = \lim_{x \rightarrow 3} \frac{(x - 3)(x + 3)}{x - 3} = \lim_{x \rightarrow 3} (x + 3) = 6$$

Problem 12. Graph the two functions $f(x) = x + 3$ and $g(x) = \frac{x^2 - 9}{x - 3}$, and list their domains. Are f and g the same function?

Problem 13. Compute the following limit: $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$.

Problem 14. Compute these limits and make a note that you will need them in a problem later!

1. $\lim_{t \rightarrow 0} \frac{\sin(t)}{t}$.

2. $\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t}$

Problem 15. Let x be a real number and compute $\lim_{h \rightarrow 0} \frac{hx^2 + h}{h}$. Your answer will be a number written in terms of the number x .

Problem 16. Compute the following limits.

1. $\lim_{x \rightarrow 3} \frac{x^3 - 27}{x - 3}$

2. $\lim_{x \rightarrow 1} \frac{\frac{1}{x} - 1}{x^2 - 1}$

3. $\lim_{x \rightarrow -2} \frac{x^3 + 3x^2 + 3x + 2}{x^2 + x - 2}$

There are rules (theorems) that will facilitate the computation of limits. The next problem will help you find these basic rules.

Problem 17. Read Problem 18 and then compute the following limits.

1. $\lim_{x \rightarrow 3} 5$

2. $\lim_{x \rightarrow 3} x$

3. $\lim_{x \rightarrow \frac{\pi}{2}} 6 \sin(x)$ and $6 \lim_{x \rightarrow \frac{\pi}{2}} \sin(x)$
4. $(\lim_{x \rightarrow 4} x^2 + \lim_{x \rightarrow 4} 4x^5)$ and $\lim_{x \rightarrow 4} (x^2 + 4x^5)$
5. $\left(\lim_{x \rightarrow -3} x^2 \right) \cdot \left(\lim_{x \rightarrow -3} \frac{1}{x} \right)$ and $\lim_{x \rightarrow -3} \left(x^2 \cdot \frac{1}{x} \right)$
6. $\lim_{x \rightarrow 5} \frac{x^2}{x-4}$ and $\frac{\lim_{x \rightarrow 5} x^2}{\lim_{x \rightarrow 5} x-4}$
7. $\lim_{x \rightarrow 2} \sqrt{x^2 + x + 3}$ and $\sqrt{\lim_{x \rightarrow 2} (x^2 + x + 3)}$
8. $\lim_{x \rightarrow -1} (4x + 1)^3$ and $\left[\lim_{x \rightarrow -1} (4x + 1) \right]^3$

Problem 18. Write five theorems (facts) about limits from your observations in the previous problems.

Problem 19. Graph and state the domain for the following functions.

1. $p(x) = 3x^2 - 6x$
2. $f(x) = \begin{cases} -2x - 4 & \text{if } x < 1 \\ x + 1 & \text{if } x > 1 \end{cases}$
3. $g(x) = \begin{cases} x^2 + 2 & \text{if } x < 1 \\ 0 & \text{if } x = 1 \\ 2x + 1 & \text{if } x > 1 \end{cases}$

Sometimes we are interested in the limit of a function f as x approaches a from the left or right side. To make clear which side we are discussing, we write $\lim_{x \rightarrow a^-} f(x)$ (left-hand limit) or $\lim_{x \rightarrow a^+} f(x)$ (right-hand limit). If both the left- and right-hand limits exist and are equal, then the limit exists. Otherwise, the limit does not exist. This can happen when either the left-hand limit or the right-limit does not exist, or when both exist but are not equal.

Theorem 1. Limit Existence Theorem. If f is a function, then the $\lim_{x \rightarrow a} f(x)$ exists if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist and are equal.

Problem 20. Using the functions from the previous problem, compute the following limits if they exist. If they do not exist, state why.

1. $\lim_{x \rightarrow -3^+} p(x)$, $\lim_{x \rightarrow -3^-} p(x)$, and $\lim_{x \rightarrow -3} p(x)$
2. $\lim_{x \rightarrow 1^-} f(x)$, $\lim_{x \rightarrow 1^+} f(x)$, and $\lim_{x \rightarrow 1} f(x)$
3. $\lim_{x \rightarrow 1^-} g(x)$, $\lim_{x \rightarrow 1^+} g(x)$, and $\lim_{x \rightarrow 1} g(x)$

Definition 4. If f is a function and a is in the domain of f then we say that f is **continuous at a** if $\lim_{x \rightarrow a} f(x) = f(a)$.

There are three subtle points in this definition. The definition requires that for f to be continuous at a , each of the following conditions must be met.

1. a must be in the domain of f (otherwise $f(a)$ doesn't exist),

2. $\lim_{x \rightarrow a} f(x)$ must exist, and
3. these two numbers must be equal.

The *absolute value* function is an example of what we call a *piecewise defined* function.

Definition 5. The **absolute value function** is defined by:

$$|x| = \begin{cases} +x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Problem 21. Determine whether the following functions are continuous at the indicated points.

1. $f(x) = \frac{1}{x+2}$ at $x = -2$.
2. $g(x) = \begin{cases} -x-1 & \text{if } x < 1 \\ x-3 & \text{if } x \geq 1 \end{cases}$ at $x = 1$.
3. $k(x) = \begin{cases} \frac{x^2-4}{x-2} & \text{if } x \neq 2 \\ 5 & \text{if } x = 2 \end{cases}$ at $x = 1$.
4. $f(x) = 4x^3 + x^2 - 6x - 7$ at $x = -2$
5. $C(x) = \cos(x)$ at $x = \frac{\pi}{3}$
6. $a(x) = \frac{|x-2|}{x-2}$ at $x = 2$.

Definition 6. We say that a function is **continuous on a set** (such as an open interval) if it is continuous at every point in the set.

Problem 22. For the following, list the open intervals on which the function is continuous.

1. $f(x) = x^2 + 5x + 3$
2. $g(x) = \frac{3x-1}{2x+5}$
3. $H(x) = \sin(x) + 5x + \sqrt{1-x}$
4. $T(z) = 2 + \tan(z)$

Theorem 2. Continuous Functions Theorem.

1. Every polynomial is continuous on $(-\infty, \infty)$.
2. Every rational function is continuous on its domain.
3. Every trigonometric function is continuous on its domain.
4. The sum, product, and difference of continuous functions is continuous.
5. The quotient $\frac{f}{g}$ of continuous functions f and g is continuous wherever g is non-zero.
6. If g is continuous at a and f is continuous at $g(a)$ then $f \circ g$ is continuous at a .

Problem 23. Compute the following limits:

1. $\lim_{h \rightarrow 0} \frac{9h^2 - 3h}{h}$
2. $\lim_{h \rightarrow 0} \frac{x^2h^2 - xh}{h}$, where x is some real number.
3. $\lim_{x \rightarrow 2} (2xt + 3x^2t^2)$, where t is some real number.

Problem 24. Let $f(x) = x^2 + x + 1$.

1. Compute and simplify $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$.
2. Compute and simplify $\lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x}$.

Example 4. Graph $f(x) = x^2$, $g(x) = -x^2$, and $b(x) = x^2 \sin\left(\frac{1}{x}\right)$. Compute $\lim_{x \rightarrow 0} f(x)$ and $\lim_{x \rightarrow 0} g(x)$.

What can you conclude about $\lim_{x \rightarrow 0} b(x)$?

Example 5. Show that $\lim_{h \rightarrow 0} \frac{\sin(h)}{h} = 1$ using the unit circle and starting with the fact that $\sin(h) \leq h \leq \tan(h)$.

The previous two examples illustrate the **Squeeze Theorem**. The importance of the second example will become clear later in the course when we take the derivative of the trigonometric function, sine. The notation $f \in C_{[a,b]}$ translates as “ f is an element of the set of all continuous function on the interval $[a,b]$,” which means that f is continuous at every number x in the interval $[a,b]$. When we write g is between f and h , on $[a,b]$ we mean that for every number $x \in [a,b]$ we have either $f(x) \leq g(x) \leq h(x)$ or $h(x) \leq g(x) \leq f(x)$.

Theorem 3. The Squeeze Theorem. If $f, g, h \in C_{[a,b]}$ and $c \in [a,b]$ and g is between f and h on $[a,b]$, and $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L$ for some number L , then $\lim_{x \rightarrow c} g(x) = L$.

Problem 25. Use the **Squeeze Theorem** to solve each of these problems.

1. Compute $\lim_{x \rightarrow 0} (x^2 \cos(x))$.
2. Compute $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right)$ using $f(x) = -x$, $g(x) = x \sin\left(\frac{1}{x}\right)$, and $h(x) = x$.
3. Compute $\lim_{x \rightarrow \infty} \frac{x}{x^2 + x + 1}$ using $f(x) = \frac{1}{x}$, $g(x) = \frac{x}{x^2 + x + 1}$, and $h(x) = 0$.

1.3 Instantaneous Velocity and Growth Rate

If you have been struggling, relax. This is where it all comes together.

Problem 26. Recall from Problem 2, that Robbie’s position function is $R(t) = t^3 + 2$.

1. What is Robbie’s average velocity from time $t = 3$ till time $t = 4$?

2. What is Robbie's average velocity from time $t = 3$ till time $t = 3.5$?
3. What is Robbie's average velocity from time $t = 3$ till time $t = 3.1$?
4. What is Robbie's instantaneous velocity at time $t=3$?

Problem 27. Fill in Table 1.3 of average velocities for Robbie. Write down an equation for a function with input time and output the average velocity of Robbie over the time interval $[3,t]$.

Table 1.3: Robbie's Average Velocity Table

Time	Time Interval	Average Velocity
4	$[3, 4]$	
3.5	$[3, 3.5]$	
3.1	$[3, 3.1]$	
3.01	$[3, 3.01]$	

Problem 28. A population P of robots (relatives of Robbie) is growing according to Table 1.4, where time is measured in days and population is measured in number of robots. Determine a population function P that represents the number of robots as a function of time and graph this function.

Table 1.4: Population Table

Time	Population
0	2
1	4
2	12
3	32
4	70
5	132

Problem 29. Using the function P from the previous problem:

1. Assume that c is a positive number and compute $P(3)$ and $P(c)$. What do these numbers represent about the robots?
2. Determine the value of $P(5) - P(3)$. What does this number say about the robot population?
3. Assume that u and v are positive numbers with $u > v$. Compute the value of $P(u) - P(v)$ and explain what this expression says about the robot population.

Problem 30. Using the function, P , from the previous problem:

1. Compute the value of $\frac{P(5) - P(3)}{5 - 3}$ and explain the physical meaning of this number.
2. Assume that u and v represent times. Simplify and explain the meaning of $\frac{P(u) - P(v)}{u - v}$.
3. What is the population of robots when time is equal to 5? At time equal to 5.5?

4. At what time is there a population of 106 robots?

Problem 31. A Maple leaf grows from a length of 5.62 cm to a length of 8.12 cm. in 1 year and 3 months. How fast did the leaf grow? Does your answer represent an instantaneous rate of growth or an average rate of growth?

Problem 32. Experimental data indicates that on January 1, the leaf is 8.37 centimeters long and it grows at the rate of 0.18 centimeters per month for 3.5 months. From that time on, the leaf grows at a rate of 0.12 centimeters per month until it reaches a length of 9.6 centimeters. Determine the average growth rate for the whole time period. At what time did the leaf grow at this rate?

Problem 33. Use the robot population function you found in Problem 28 to fill in Table 1.5 and determine how fast the robot population is growing at time $t=4$. This is called the instantaneous growth rate and is measured in robots per day.

Table 1.5: Population Growth Rate Table

Time	Time Interval	Average Growth Rate
5	[4, 5]	
4.5	[4, 4.5]	
4.1	[4, 4.1]	
4.01	[4, 4.01]	

Problem 34. Write a function with input time and output the average growth rate of the population over the time interval $[4, t]$.

Problem 35. State the limit that would produce the limit table from Problem 27 and hence establish the instantaneous velocity of Robbie at time $t=3$.

Problem 36. State the limit that would produce the limit table from Problem 33 and hence establish the instantaneous growth rate of the population at time $t=4$.

At this point, you know how to compute average velocity, average growth rate, instantaneous velocity, and instantaneous growth rate. You don't want to create a limit table every time you want the instantaneous velocity, so we want a function for Robbie's velocity. That is, we want a function that takes *time* as input and produces Robbie's *velocity* as output. Similarly, we want a function that takes *time* as input and produces the *robot population growth rate* as output.

Problem 37. Write a function for the instantaneous velocity of Robbie.

Problem 38. Write a function for the instantaneous growth rate of the robot population.

Roberta the Robot (Robbie's potential girlfriend) is walking along the same road as Robbie. Roberta's distance from the same pothole after t seconds is given by $f(t) = 2t^2 - 16$ feet. Recall from Problem 37, we know that if we have a position function then we can find the corresponding velocity function by taking a certain limit.

Problem 39. Find the velocity function for Roberta.

Problem 40. Determine Roberta's velocity at 10 seconds. At what time(s) is Roberta 2 feet from the pothole?

Problem 41. At what time(s) is Roberta stopped? At what time is Roberta traveling at a velocity of -4 feet per second? Do Robbie and Roberta ever meet? If so, when and where?

A colony of **androids** is threatening our robot population. On day t , there are $N(t) = 2t^2 + 4t + 6$ androids.

Problem 42. Find the growth rate function for the android population. What are the units? At what time(s) is the android population equal to 200?

Problem 43. When is the android population increasing at a rate of 10 androids per day? How fast is the androids population growing on day 10?

1.4 Tangent Lines

We are going to define **tangent line** from a physical point-of-view rather than give a rigorous mathematical definition. Sketch any function, f , and pick a point on it. Label your point $(a, f(a))$. Place your pencil on the graph at a point to the left of $(a, f(a))$. Imagine that your pencil is a car, the graph is a road, and you are driving toward the point $(a, f(a))$ from the left. Suppose that there is a perfectly frictionless patch of ice at the point $(a, f(a))$ and keep drawing in the direction the car would travel as it slides off the road. Now do the same from the right. If the two lines you drew form a line, then you have drawn the **tangent line to f at the point $(a, f(a))$** . If they don't form a line, then f does not have a tangent line at this point.

Definition 7. A **secant line** for the graph f is a line passing through two points on the graph of f .

Problem 44. Let $f(x) = x^2 + 1$ and graph f . Graph and compute the slope of the secant lines passing through:

1. $(2, f(2))$ and $(4, f(4))$
2. $(2, f(2))$ and $(3, f(3))$
3. $(2, f(2))$ and $(2.5, f(2.5))$
4. $(2, f(2))$ and $(2.1, f(2.1))$

What is the slope of the line tangent to f at the point $(2, 5)$?

Problem 45. Write down a limit that represents the work you did in the last problem.

Problem 46. Find the slope of the tangent line to the curve $f(x) = x^3 + 2$ at the point $(3, 29)$.

Problem 47. Find the equation of the tangent line in Problem 46.

Problem 48. Assume $a \in \mathbb{R}$. Find the equation of the line tangent to the graph of $f(x) = x^3 + 2$ at the point $(a, f(a))$.

Chapter 2

Derivatives

“...the secret of education lies in respecting the pupil...” - Ralph Waldo Emerson

Up to this point, we have solved several different problems and discovered that the same concept has generated a solution each time. Limits were used to find:

- Robbie’s velocity function given his position function,
- the instantaneous growth function for the robot population given the population function, and
- the function that tells us the slope of the line tangent to a function at any point.

The following definition is the result of all of our work to this point.

Definition 8. Given a function f , the **derivative** of f , denoted by f' , is the function so that for every point x in the domain of f where f has a tangent line, $f'(x)$ equals the slope of the tangent line to f at $(x, f(x))$. Either

$$f'(x) = \lim_{b \rightarrow x} \frac{f(b) - f(x)}{b - x} \quad \text{or} \quad f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

may be used to compute the derivative.

While both limits above will always yield the same result, there are times when one may be preferable to the other in terms of the algebraic computations required to obtain the result. Given a function f , the derivative of f is usually denoted by f' , $\frac{df}{dx}$ or $\dot{f}$. Regardless of how it is written, the derivative of f at x represents the slope of the line tangent to f at the point $(x, f(x))$ or the rate at which the quantity f is changing at x . The origins of calculus are generally credited to Leibnitz and Newton. The first notation, credited to Lagrange, is referred to as *prime* or *functional* notation. The second notation is due to Leibnitz. The third is due to Newton. Engineers sometimes use the dot to represent derivatives that are taken with respect to time.

If $f(x) = x^2$, then each of

$$f'(x) \text{ and } (x^2)' \text{ and } \frac{d}{dx}(x^2)$$

all mean “the derivative of f at x .”

2.1 Derivatives of Polynomial, Root, and Rational Functions

Problem 49. Compute the derivative of each function using Definition 8.

1. $f(x) = x^2 - 3x$

2. $g(x) = 3x - 7$

3. $h(t) = 4t^3$

4. $u(t) = \frac{1}{t}$

5. $v(x) = \frac{x-5}{x+2}$

Problem 50. Compute the following using the derivatives you computed in the previous problem. Graph both the original function and the tangent line at the point of interest.

1. $f'(x)$ at $x = 5$

2. $h'(t)$ at $t = e$ (Recall that Euler's constant, e , is approximately 2.71828...)

3. $v'(x)$ at $x = -2$

Algebra Reminder. An expression such as $\frac{\sqrt{a} - \sqrt{b}}{a - b}$, can be simplified by multiplying the numerator and the denominator of the fraction by $\sqrt{a} + \sqrt{b}$. This is called *rationalizing the numerator*.

Problem 51. If $f(x) = \sqrt{x+1}$, compute f' . Graph f and the tangent line to f at $(1, f(1))$ and state the slope of the tangent line.

Problem 52. If $f(x) = 2x^3 + 3x^2 - 7$, compute f' . Graph f and compute the equation of the tangent line to f at $(1, f(1))$.

Problem 53. If $f(x) = \frac{1}{x+1}$, compute f' .

Problem 54. If $f(x) = \sqrt{x+1}$, find the equation of the tangent line at $x = a$.

Problem 55. For each of the following, check the left- and right-hand limits to determine if the limit exists:

1. $\lim_{x \rightarrow 0} \frac{0}{x}$

2. $\lim_{x \rightarrow 0} \frac{|x|}{x}$

Definition 9. We say that a function, f , is **differentiable at a** if $f'(a)$ exists.

Definition 10. We say that a function, f , is **differentiable** if f is differentiable at every point in its domain.

Definition 11. We say that a function, f , is **left differentiable at a** if $\lim_{t \rightarrow a^-} \frac{f(t) - f(a)}{t - a}$ exists. If f is left differentiable at a , we denote the **left derivative** of f at a by $f'(a-)$. Right differentiability is defined similarly.

Definition 12. We say that f is **differentiable on (a,b)** if f is differentiable at t for all t in (a,b) .

Definition 13. We say that f is **differentiable on $[a,b]$** if f is differentiable on (a,b) and $f'(a+)$ and $f'(b-)$ both exist.

Problem 56. Let $f(x) = |x - 2|$.

1. Graph f .
2. Determine the equation of the line tangent to f at $(-1, 3)$.
3. Determine the equation of the line tangent to f at $(5, 3)$.
4. Compute $f'(2+)$ and $f'(2-)$.
5. Does $f'(2)$ exist? Why?
6. What is the derivative of f ?

Example 6. Illustrate ways in which a function might fail to be differentiable at a point c such as:

- the point c might not be in the domain of the function,
- the function might have a vertical tangent line at c , or
- the left and right-hand derivatives might not agree at c (i.e. $f'(c+) \neq f'(c-)$).

Problem 57. For each of the following, use a sketch of the graph to determine the set of those points at which the function is not differentiable.

1. $f(x) = \frac{3}{1-x^2}$
2. $g(x) = \sqrt{5+x}$
3. $i(x) = \sec(x)$

Theorem 4. Differentiable Functions Theorem.

1. Every polynomial is differentiable on $(-\infty, \infty)$.
2. Every rational function is differentiable on its domain.
3. Every trigonometric function is differentiable on its domain.
4. The sum, product, and difference of differentiable functions is differentiable.
5. The quotient $\frac{f}{g}$ of differentiable functions f and g is differentiable wherever g is not zero.
6. If g is differentiable at $x = a$ and f is differentiable at $x = g(a)$ then $f \circ g$ is differentiable at $x = a$.

We know how to compute derivatives of functions using limits. By proving theorems we will increase the ease with which we compute derivatives of more challenging functions.

Problem 58. Compute f' for each function listed, using the definition of the derivative.

1. $f(x) = a$, where a is a constant
2. $f(x) = x$

3. $f(x) = x^2$

4. $f(x) = x^3$

5. $f(x) = x^4$

At this point we have used the definition of the derivative (limits) to compute the derivatives of many different functions, including: a (constant function), x (the identity function), x^2 , x^3 , $4x^3$, x^4 , $\sqrt{x+1}$, $2x^3 + 3x^2 - 7$, $\frac{1}{x}$, $\frac{1}{x+1}$, $\frac{x-5}{x+2}$, $|x-2|$, and $x^3 + 2$. Based on the derivatives we have computed thus far, you can likely guess the answer to the next question.

Question. What is the derivative of $f(x) = x^{200}$?

Guess. $f'(x) = 200x^{199}$.

How do we *know* if our guess is correct? We don't. We must show that it is true using the definition of the derivative. Rather than work this one problem and still not know what the derivative of $g(x) = x^{201}$ is, we'll let n represent some unknown natural number and compute the derivative of $f(x) = x^n$. Because we will not choose a specific value for n , we will now *know* the derivative of an entire class of functions.

Problem 59. Power Rule. Let n be a natural number and use Definition 8 to show that the derivative of $f(x) = x^n$ is $f'(x) = nx^{n-1}$.

Problem 60. Constant Rule. Show that if k is a constant (i.e. any real number) and $h(x) = kf(x)$, then $h'(x) = kf'(x)$.

Note. The previous problem is not stated accurately. Here is a precise statement. Show that if k is a constant (i.e. any real number) and f is differentiable on (a, b) and $h(x) = kf(x)$ for all x in the interval (a, b) , then $h'(x) = kf'(x)$ for all x in (a, b) .

Problem 61. Sum Rule. Show that if each of f and g is a function and $S(x) = f(x) + g(x)$, then $S'(x) = f'(x) + g'(x)$.

Problem 62. Difference Rule. Show that if each of f and g is a function and $D(x) = f(x) - g(x)$, then $D'(x) = f'(x) - g'(x)$.

Problem 63. Use Problems 59, 60, 61, and 62 to compute f' when $f(x) = 6x^5 - 4x^4 + 7x^3$. Apply one rule per equal sign and be sure to explain which rules you are using.

The next theorem allows us to avoid using limits to determine if a function is continuous, since it states that all differentiable functions are also continuous functions.

Theorem 5. A Continuity Theorem. If x is a number and f is differentiable at x , then f is continuous at x .

Problem 64. Product Rule. Show that if $P(x) = f(x)g(x)$, then $P'(x) = f'(x)g(x) + f(x)g'(x)$.

Problem 65. Quotient Rule. Show that if $Q(x) = \frac{f(x)}{g(x)}$, then $Q'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$.

Problem 59 only proved the power rule for exponents that were positive integers. With the new tools, the next problem extends the power rule to include negative and zero exponents. A natural question would then be, "Is this true for exponents that are real numbers but not integers?" The answer is "yes" but we'll simply assume this and not prove it in this course.

Problem 66. Power Rule Revisited. Show that if n is any integer and $f(x) = x^n$, then $f'(x) = nx^{n-1}$. Before you start, ask yourself how this is different from Problem 59.

Problem 67. Suppose a, b, c , and d are constants such that $d \neq 0$, and find f' where $f(x) = \frac{ax^2 + bx + c}{d}$.

Problem 68. Find f' when f is the function defined by $f(x) = (x - \frac{1}{x})(2x^2 + 3x + 4)$.

Problem 69. Find f' when f is the function defined by $f(x) = \frac{x-1}{x+1} \cdot \frac{x-2}{x+2}$.

Problem 70. Find f' when f is the function defined by $f(r) = \frac{r^2s^2 - r^4}{s - m}$ and s and m are real numbers and $s \neq m$.

Since the derivative of a function is itself a function, there is no reason not to take the derivative of the derivative. When this is done, the new function is called the **second derivative** and is denoted by f'' . Similarly the **third derivative** is denoted by f''' .

Problem 71. Find f'' when $f(x) = \frac{2}{x} - \frac{x^4}{2}$.

You have solved *algebraic* equations such as

$$5x + 2 = 3x + 4 \quad \text{or} \quad 3x^2 - 2x + 1 = 0$$

where the goal is to find each value of x that solves the equations. Engineering applications often need to solve *differential* equations such as

$$f'(x) = xf(x) \quad \text{or} \quad f''(x) = x^3 \quad \text{or} \quad f''(x) + f(x) = 0$$

where the goal is to find each function f that satisfies the equation. In the real world, solutions to differential equations might represent the shape of an airplane wing or the temperature distribution of an engine block.

Problem 72. Show that if x is not 0, then the function $f(x) = \frac{2}{x}$ is a solution to the equation $x^3 f''(x) + x^2 f'(x) - xf(x) = 0$.

Problem 73. Find one function f that satisfies $f'(x) = 4x^2 - 2x$.

We have been very careful up to this point in the book to write the equations in *functional* notation. That is, we never wrote, $y = x^2$, but rather wrote $f(x) = x^2$ to *emphasize* that f is the function and it depends on the *independent* variable, x . As you will see in other courses, it is very common to write, $y = x^2$ and then follow this with $y' = 2x$. Differential equations are almost always written in this short-hand, omitting the independent variable where possible, and from this point forward, we too will use this short-hand on occasion.

Problem 74. Sketch the graph of $y = 2x^3 - 3x^2 - 12x + 2$ and locate (by using the derivative) the points on the graph that have a horizontal tangent line.

Problem 75. Find an equation for the line that is tangent to $y = \frac{2-x}{2+x}$ at $(1, \frac{1}{3})$.

Algebra Reminder. Any point where a function crosses the x -axis is called an **x -intercept** of the function, and any point where a function crosses the y -axis is called a **y -intercept** of the function. To find an x -intercept we set y equal to zero and solve for x . To find a y -intercept we set x equal to zero and solve for y .

Problem 76. Find numbers a , b , and c so that the function $f(x) = ax^2 + bx + c$ has an x -intercept of 2, a y -intercept of 2, and a tangent line with slope of 2 at $(2, 4a + 2b + c)$.

Problem 77. Sketch the function $f(x) = ax^2$, where $a > 0$. Let c and d be two (distinct) real numbers. Let x be the x -coordinate of the point where the tangent line to f at $(c, f(c))$ intersects the tangent line to f at $(d, f(d))$. Show that the distance between x and c equals the distance between x and d .

Problem 78. Let $f(x) = \frac{2}{x}$ and c be any real number. Find the area of the triangle ($A = \frac{1}{2}bh$) formed by the tangent line to the graph at $(c, f(c))$ and the coordinate axes.

Problem 79. Prove that if the graph of $h(x) = f(x) - g(x)$ has a horizontal tangent line at $(c, h(c))$, then the tangent line to f at $(c, f(c))$ is parallel to the tangent line to g at $(c, g(c))$.

Problem 80. Show that if the function f is differentiable at x and $g(x) = (f(x))^2$, then $g'(x) = 2f(x)f'(x)$.

Problem 81. Sketch a graph of

$$f(x) = \begin{cases} x^2 & \text{if } x \leq 1 \\ \sqrt{x} & \text{if } x > 1. \end{cases}$$

Use left and right hand derivatives at $x = 1$ to determine if f is differentiable at $x = 1$.

2.2 Derivatives of the Trigonometric Functions

When a mathematician says something is “true,” s/he means that s/he can prove it from a set of axioms and definitions. This is called *deductive* reasoning. We used deductive reasoning on Problem 64 (product rule) and on Problem 65 (quotient rule) by proving (deducing) them from the definition of the derivative. When a scientist says something is true, s/he means s/he believes it to be true based on data. This is called *inductive* reasoning because s/he *induces* the truth from observations. We are going to determine the derivative of $s(x) = \sin(x)$ in two different ways, inductively and deductively.

Problem 82. Inductive: Use the unit circle to fill in the second row of Table 2.1 and make a very accurate graph of the sine function using this data, labeling each of these points on the graph. Sketch the tangent line to the graph of sine at each of these points.

Table 2.1: The Derivative of $s(x) = \sin(x)$

x	0	$\pi/4$	$\pi/2$	$3\pi/4$	π	$5\pi/4$	$3\pi/2$	$7\pi/4$	2π
$s(x)$									
$s'(x)$									

Problem 83. Approximate the slope of each tangent line you graphed in the last problem and fill in the third row of Table 2.1 with this data. Sketch this new function, the derivative of sine, by plotting the points $(x, s'(x))$ from the table. Does the graph of s' look familiar?

Based on the last two problems, we have a guess for the derivative of sine. Now, we'll see if our guess is correct.

Trigonometry Reminder – The Sum and Difference Identities. Appendix 15.4 has most of the common trigonometric identities, in case you need them later.

$$1. \sin(x \pm y) = \sin(x)\cos(y) \pm \cos(x)\sin(y)$$

$$2. \cos(x \pm y) = \cos(x)\cos(y) \mp \sin(x)\sin(y)$$

Problem 84. Deductive: Compute the derivative of the sine function using Definition 8 and the Trigonometry Reminder above.

Problem 85. Compute the derivative of $f(x) = \cos(x)$.

Problem 86. Compute the derivatives of the tangent, cotangent, secant, and cosecant functions by using the derivatives of sine and cosine along with the quotient rule.

Problem 87. Find a function f with derivative $f'(x) = 3\cos(x) - 2\sin(x) + x + 2$.

Problem 88. If $y = \frac{\cos(x)}{x+3}$, compute y' .

Problem 89. If $y = \sin(x)\cos(x)$, compute y' .

Problem 90. Suppose that L is a line tangent to $y = \cos(x)$ at the point, $(c, f(c))$ and L passes through the point $(0,0)$. Show that c satisfies the equation $t = -\cot(t)$. That is show that $c = -\cot(c)$.

Problem 91. Find the equation of the line tangent to the curve $y = \tan(x) + \pi$ at the point on the curve where $x = \pi$. Repeat this exercise for $x = \pi/4$.

2.3 The Chain Rule

We can take the derivative of $f(x) = \sqrt{x}$ and $g(x) = 5x + 1$ easily, but to take the derivative of $h(x) = \sqrt{5x+1}$, we would need to return to the limit definition for the derivative. We can take the derivative of $f(x) = \sin(x)$ and $g(x) = x^2$ easily, but to take the derivative of $h(x) = \sin(x^2)$ we would again return to the limit definition. The *chain rule* remedies this problem by giving a rule for computing the derivatives of functions that are the composition of two functions.

Algebra Reminder. The function $h(x) = \sqrt{5x+1}$ is the composition of two functions $f(x) = \sqrt{x}$ and $g(x) = 5x + 1$ since we may write $f(g(x)) = f(5x + 1) = \sqrt{5x+1} = h(x)$. The function $h(x) = \sin(x^2)$ is the composition of $f(x) = \sin(x)$ and $g(x) = x^2$. If h equals f composed with g , we write $h = f \circ g$ and $h(x) = f(g(x))$ for all numbers x in the domain of h .

Theorem 6. The Chain Rule. If f and g are differentiable functions and $h(x) = f(g(x))$, then $h'(x) = f'(g(x)) \cdot g'(x)$.

A sort-of proof. Assume that each of f and g is a differentiable function and that $h(x) = f(g(x))$. From the limit definition of the derivative of h , we know that

$$h'(x) = \lim_{t \rightarrow x} \frac{h(t) - h(x)}{t - x}.$$

Since $h(x) = f(g(x))$,

$$h'(x) = \lim_{t \rightarrow x} \frac{f(g(t)) - f(g(x))}{t - x}.$$

Since

$$\frac{g(t) - g(x)}{g(t) - g(x)} = 1 \text{ as long as } g(t) \neq g(x),$$

by using the *Product Rule for Limits* we may write,

$$h'(x) = \lim_{t \rightarrow x} \frac{f(g(t)) - f(g(x))}{g(t) - g(x)} \cdot \frac{g(t) - g(x)}{t - x} = \lim_{t \rightarrow x} \frac{f(g(t)) - f(g(x))}{g(t) - g(x)} \cdot \lim_{t \rightarrow x} \frac{g(t) - g(x)}{t - x}.$$

Since g is differentiable, we know by Theorem 5 that g is continuous, so by the definition of continuity, we have $\lim_{t \rightarrow x} g(t) = g(x)$. Thus, in the first limit, we can replace $t \rightarrow x$ with $g(t) \rightarrow g(x)$.

$$h'(x) = \lim_{g(t) \rightarrow g(x)} \frac{f(g(x)) - f(g(t))}{g(x) - g(t)} \cdot \lim_{t \rightarrow x} \frac{g(x) - g(t)}{t - x}.$$

The right-hand side of this equation is the product of two limits. The second limit is the definition of $g'(x)$. Substituting $A = g(t)$ and $B = g(x)$ into the first limit, we may write this as,

$$\lim_{A \rightarrow B} \frac{f(B) - f(A)}{B - A}.$$

But this is just $f'(B)$, or $f'(g(x))$, so we have shown that

$$h'(x) = f'(g(x)) \cdot g'(x).$$

q.e.d.

Problem 92. Let $h(x) = \sqrt{7 + \sin(x)}$.

1. Find functions f and g such that $h = f \circ g$.
2. Compute f' .
3. Compute $f' \circ g$.
4. Compute g' .
5. Use the Chain Rule to compute h' .

Problem 93. Let $h(x) = \cos(x^2)$.

1. Find functions f and g such that $h = f \circ g$.
2. Compute f' .
3. Compute $f' \circ g$.
4. Compute g' .
5. Use the Chain Rule to compute h' .

Problem 94. Given that $f(x) = (x^{5/2} - 4x^{1/3} + 365)^{42}$, compute f' .

Problem 95. If $y = (\cos(x^2))^2$, compute y' .

Problem 96. Let $y = \left(\frac{1-x^2}{1+x^2}\right)^{10}$ and compute y' using the Chain Rule first and then the Quotient Rule. Check your answer by rewriting $y = \frac{(1-x^2)^{10}}{(1+x^2)^{10}}$ and computing y' using the Quotient Rule first and then the Chain Rule.

Problem 97. Find a function f with derivative $f'(x) = 5x + 3$.

Problem 98. Find the equation of the tangent line to the curve $y = (x + 1/x)^3$ at the point where $x = -1$. Graph the curve and the line.

Problem 99. Assume a, b, c and d are real numbers and $f(w) = a(\cos(wb))^2 + c(\sin(wd))^2$. Compute f' .

Problem 100. Find the real number m such that $y = m \cos(2t)$ satisfies the differential equation $y'' + 5y = 3 \cos(2t)$.

Problem 101. Given that $f'(x) = \sqrt{2x+3}$, $g(x) = x^2 + 2$, and $F(x) = f(g(x))$, compute F' .

Problem 102. Given that $f'(x) = \frac{x}{x^2-1}$ and $g(x) = \sqrt{2x-1}$, compute F' where $F(x) = f(g(x))$.

2.4 Derivatives of the Inverse Trigonometric Functions

Definition 14. Given a function f with domain D and range R , if there exists a function g with domain R and range D so that for all $x \in D$ and all $y \in R$ we have

$$g(f(x)) = x \text{ and } f(g(y)) = y,$$

then we call g the **inverse** of f .

Example 7. Discuss inverse functions and compute the inverse of $f(x) = \frac{1}{x-3}$.

Problem 103. Find the inverse of each of the following functions.

1. $f(x) = \frac{2}{3}x + 4$
2. $g(x) = x^3 - 1$
3. $h(x) = \sqrt[3]{x-4}$
4. $i(x) = 4 - x^5$

Problem 104. 1. Does $n(x) = x^2$ have an inverse? If so what is it? If not, why not?
2. Does $y(x) = \sqrt{x}$ have an inverse? If so what is it? If not, why not?

Notice in the next definition that we restrict the domain of sine to talk about the inverse sine. This is because if we simply reversed all the coordinates of sine, the resulting graph would not be a function. There are many ways we could have restricted the domain, so one simply must memorize the domain and range of all the inverse trigonometric functions.

Table 2.2: Inverse Trigonometric Functions

Inverse Function	Domain	Range
$\sin^{-1}$	$[-1, 1]$	$[-\pi/2, \pi/2]$
$\cos^{-1}$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1}$	$\mathbb{R}$	$(-\pi/2, \pi/2)$
$\csc^{-1}$	$ x \geq 1$	$(-\pi/2, \pi/2) \cup (\pi, 3\pi/2)$
$\sec^{-1}$	$ x \geq 1$	$[0, \pi/2) \cup [\pi, 3\pi/2)$
$\cot^{-1}$	$\mathbb{R}$	$(0, \pi)$

Definition 15. The **inverse sine** function, denoted by $\sin^{-1}$ or $\arcsin$ is the inverse of the function $f(x) = \sin(x)$ with the domain $[-\pi/2, \pi/2]$.

The domain of the inverse sine function is the range of the sine function and the range of the inverse sine function is the restricted domain, $[-\pi/2, \pi/2]$, of our sine function. We define all six inverse trigonometric functions in a similar manner and Table 2.2 lists these functions along with their respective domains and ranges.

Theorem 7. Inverse Sine Derivative Theorem. If $f(x) = \sin^{-1}(x)$, then $f'(x) = \frac{1}{\sqrt{1-x^2}}$.

Proof. Computing the derivative of the inverse sine function is an application of the Pythagorean Theorem and the Chain Rule. Let

$$f(x) = \sin^{-1}(x).$$

Since

$$\sin(f(x)) = \sin(\sin^{-1}(x)) = x,$$

we have,

$$\sin(f(x)) = x.$$

Taking the derivative of both sides yields,

$$\cos(f(x))f'(x) = 1,$$

and solving for $f'(x)$ yields

$$f'(x) = \frac{1}{\cos(f(x))}.$$

Because we don't want f' in terms of f , we apply the Pythagorean Theorem. Draw a right triangle with one angle, $f(x)$. Since

$$\sin(f(x)) = x = \frac{x}{1}$$

place x on the side opposite the angle $f(x)$ and 1 on the hypotenuse. Note that

$$\cos(f(x)) = \frac{\sqrt{1-x^2}}{1} = \sqrt{1-x^2}.$$

Thus

$$f'(x) = \frac{1}{\sqrt{1-x^2}}.$$

q.e.d.

Problem 105. Compute the derivatives of the remaining five inverse trigonometric functions and memorize them.

Problem 106. Compute the derivative of each of the following functions.

1. $y(x) = \sin^{-1}(2x)$
2. $H(t) = (t - \cos^{-1}(t))^3$
3. $F(x) = \csc^{-1}(x)\tan^{-1}(x)$
4. $G(z) = \frac{z^2 - 7z + 5}{\cot^{-1}(z)}$
5. $h(x) = \tan^{-1}(\sin^{-1}(x))$

2.5 Derivatives of the Exponential and Logarithmic Functions

We know how to take the derivative of $f(x) = x^2$, but what about $g(x) = 2^x$ or $h(x) = e^x$? We wish to be able to differentiate exponential and logarithmic functions. If you need a review of these functions, then work through Appendix 15.3.

Problem 107. Sketch graphs of $f(x) = 2^x$ and $g(x) = 3^x$ on the same coordinate axes labeling several points and being very careful to show which function is above the other at various points on the graph.

Example 8. Defining e and finding the derivative of the function e^x .

Consider finding the derivative of $f(x) = 2^x$. Using limits to compute the derivative, we have:

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad \text{by definition of derivative} \\
 &= \lim_{h \rightarrow 0} \frac{2^{x+h} - 2^x}{h} \quad \text{by definition of } f \\
 &= \lim_{h \rightarrow 0} \frac{2^x(2^h - 1)}{h} \quad \text{by algebra} \\
 &= \lim_{h \rightarrow 0} 2^x \frac{2^h - 1}{h} \quad \text{by algebra} \\
 &= 2^x \lim_{h \rightarrow 0} \frac{2^h - 1}{h} \quad \text{by the Constant Multiple Rule for limits} \\
 &= 2^x f'(0)
 \end{aligned}$$

We would now know the derivative of the function 2^x if only we could determine $f'(0) = \lim_{h \rightarrow 0} \frac{2^h - 1}{h}$.

We can approximate $f'(0) = \lim_{h \rightarrow 0} \frac{2^h - 1}{h} \approx .69$ by using a limit table. Similarly, if we went through

the same process for $g(x) = 3^x$ we would see that $g'(0) = \lim_{h \rightarrow 0} \frac{3^h - 1}{h} \approx 1.09$ by using a limit table.

Since the slope of f at 0 is less than 1 and the slope of g at 0 is greater than one, it makes sense that there should be an exponential function which has slope *exactly* 1 at 0. We define a number,

called e , to be the number (between 2 and 3) so that $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$. Now, if we let $j(x) = e^x$ then we have

$$\begin{aligned}
 j'(x) &= \lim_{h \rightarrow 0} \frac{j(x+h) - j(x)}{h} \quad \text{by definition of derivative} \\
 &= \lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} \quad \text{by definition of } f \\
 &= \lim_{h \rightarrow 0} \frac{e^x(e^h - 1)}{h} \quad \text{by algebra} \\
 &= \lim_{h \rightarrow 0} e^x \frac{e^h - 1}{h} \quad \text{by algebra} \\
 &= e^x \lim_{h \rightarrow 0} \frac{e^h - 1}{h} \quad \text{by the Constant Multiple Rule for limits} \\
 &= e^x f'(0) \\
 &= e^x.
 \end{aligned}$$

This work results in a definition and a theorem.

Definition 16. Let e be the number that satisfies $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$.

Theorem 8. Exponential Derivative Theorem. If $f(x) = e^x$, then $f'(x) = e^x$.

Definition 17. Let $g(x) = \ln(x)$ be the function that is the inverse of the function $f(x) = e^x$.

Algebra Reminders

1. $f(x) = e^x$ has domain all $x \in \mathbb{R}$ and range all $x > 0$
2. $g(x) = \ln(x)$ has domain all $x > 0$ and range all $x \in \mathbb{R}$
3. $e^{x+y} = e^x e^y$ and $(e^x)^y = e^{xy}$ for all $x, y \in \mathbb{R}$
4. $\ln(xy) = \ln(x) + \ln(y)$ for all $x > 0, y > 0$
5. $\ln(x^y) = y \ln(x)$ for all $y \in \mathbb{R}$ and $x > 0$
6. $e^{\ln(x)} = x$ for all $x > 0$
7. $\ln(e^x) = x$ for all $x \in \mathbb{R}$
8. $\log_b(x) = \ln(x)/\ln(b)$ for all $b \in \mathbb{R}$ (except $b=1$) and for all $x > 0$

Problem 108. Compute and simplify the derivatives of the following functions:

1. $g(x) = e^x + x^2 + 2x$
2. $f(x) = xe^x$
3. $h(x) = x^3/e^x$
4. $f(x) = e^{x^2}$
5. $f(x) = 3x^2e^{x^4}$

Problem 109. Let $g(x) = \ln(x)$. Here are two clever ways to compute the derivative of g ; do which ever you like better!

1. Since we know that $g(e^x) = x$, differentiate both sides of this equation using the Chain Rule and then solve for $g'(x)$.
2. Apply the exponential function to both sides of the equations $g(x) = \ln(x)$, differentiate, and then solve for $g'(x)$.

Problem 110. Compute and simplify the derivative of the following functions:

1. $g(x) = \ln(7x - 8)$
2. $f(x) = x^2 \ln(x)$
3. $h(x) = x^3 / \ln(3x^2 - x^{1/3})$

Problem 111. Let $h(x) = 2^x$ and compute h' first by taking the natural log of both sides and then taking the derivative of both sides.

Problem 112. Compute and simplify the derivative of the following functions:

1. $g(x) = 2^{7x-8}$
2. $f(x) = 2^x - x^2$
3. $h(x) = 3^{x^2}$

Problem 113. Use the change-of-base formula, $\log_b(x) = \ln(x)/\ln(b)$, to compute the derivative of $L(x) = \log_3(x^5 - 2x)$.

2.6 Logarithmic and Implicit Differentiation

We know how to take the derivative of $f(x) = x^2$ and $g(x) = 2^x$. But what about functions like $h(x) = x^x$ where the variable occurs in both the base and the exponent? Fortunately, the technique you used in Problem 111, called **logarithmic differentiation**, and can be applied to this function as well.

Example 9. Compute the derivative of $f(x) = x^{2x}$.

Define f .

$$f(x) = x^{2x}$$

Apply the natural logarithm to both sides.

$$\ln(f(x)) = \ln(x^{2x})$$

Apply a property of logarithms to the right-hand side.

$$\ln(f(x)) = 2x \ln(x)$$

Take the derivative of both sides.

$$\frac{f'(x)}{f(x)} = 2\ln(x) + (2x)\frac{1}{x} = 2\ln(x) + 2$$

Multiply through by $f(x)$.

$$f'(x) = f(x)(2\ln(x) + 2)$$

Substitute $f(x) = x^{2x}$ and simplify.

$$f'(x) = x^{2x}(2\ln(x) + 2) = 2x^{2x}(\ln(x) + 1)$$

Problem 114. Compute the derivative of $y(x) = (3x)^{x+1}$.

Problem 115. Compute $H'(x)$ for $H(x) = (\sin(x))^{\cos(2x)}$.

Example 10. Computing derivatives of relations that are not functions.

Implicit Differentiation. Thus far we have only taken the derivatives of functions, and the derivatives of these functions were also functions. Implicit differentiation will allow us to find an equation that tells us the slopes of the lines tangent to an equation that does not represent a function, for example, a circle. For a single value of x , a circle may have two values for y and thus two tangent lines. Therefore, our “derivative” will not be a function because for a single value of x there may be two lines tangent to the circle.

Let’s look at an example. The circle $x^2 + y^2 = 9$ is a relation (a set of ordered pairs), but *not* a function since $(1, 2\sqrt{2})$ and $(1, -2\sqrt{2})$ are points that satisfy the equation but have the same x -coordinate and distinct y -coordinates. How can we determine the slope of any tangent line to this equation, recalling that the derivative was, thus far, defined only for functions? For this problem, we can do it in two ways, straight differentiation and implicit differentiation.

Straight Differentiation. First, rewrite the equation,

$$x^2 + y^2 = 9,$$

as,

$$y = \pm \sqrt{9 - x^2}.$$

This may be written in what *looks* like functional notation as,

$$f(x) = \pm \sqrt{9 - x^2},$$

but f is *not* a function since for each value of $x \in (-3, 3)$, we have two values for $f(x)$. Still, taking the derivative of either

$$f(x) = +\sqrt{9 - x^2} \quad \text{or} \quad f(x) = -\sqrt{9 - x^2}$$

is exactly the same except for the sign, so we do both simultaneously:

$$f'(x) = \pm \frac{1}{2}(9 - x^2)^{-\frac{1}{2}} \cdot -2x = \pm \frac{x}{\sqrt{9 - x^2}}.$$

Implicit differentiation is an alternative way to differentiate such equations by first taking the derivative of both sides of the equation and then solving the resulting equation for the derivative instead of solving for y and then taking the derivative.

Implicit Differentiation. We again start with

$$x^2 + y^2 = 9.$$

Replacing y with $f(x)$ as we did in our first example yields,

$$x^2 + (f(x))^2 = 9.$$

Differentiating both sides using the Power, Chain, and Constant rules yields,

$$2x + 2f(x)f'(x) = 0.$$

Solving for $f'(x)$ we have,

$$f'(x) = \frac{-2x}{2f(x)} = \frac{-x}{f(x)}.$$

This does not look the same as the answer we computed using straight differentiation, but if we again solve

$$x^2 + (f(x))^2 = 9$$

for $f(x)$ to get

$$f(x) = \pm\sqrt{9-x^2},$$

and substitute this into our answer

$$f'(x) = \frac{-x}{f(x)}$$

then we have

$$f'(x) = \frac{-x}{\pm\sqrt{9-x^2}} = \pm\frac{x}{\sqrt{9-x^2}}.$$

This agrees with the result we obtained using straight differentiation.

Problem 116. Find y' where $x^3 + y^3 = 1$ in two ways. First, solve for y and differentiate. Second, differentiate and then solve for y' . Show that both answers are the same. Graph the set of points that satisfy $x^3 + y^3 = 1$. Is this a function?

Problem 117. Find y' for $x^3 + y^3 = 3xy$.

This problem is based on a true story. Only the type of invertebrate and the name of the snack food have been changed in order to protect their identities.

Problem 118. You sketch an x - y -coordinate plane on your notebook and then note an ant walking around the page nibbling on your Cheeto crumbs. The ant's position changes with time, so you define a function p so that the position of the ant at time t is given by the function $p(t) = (x(t), y(t))$. After considerable observation, you observe that the position function satisfies $2(x(t))^2 + 3(y(t))^2 = 100$, the speed at $t = 5$ in the x -direction is 2 (i.e. $x'(5) = 2$), and the x -position at time $t = 5$ is 6 (i.e. $x(5) = 6$). Compute the ant's y position at time $t = 5$ (i.e. $y(5)$) and the ant's speed in the y direction at time $t = 5$ (i.e. $y'(5)$).

Problem 119. Given the curve $x^2 - 4x + y^2 - 2y = 4$, find:

1. the two points on the curve where there are horizontal tangent lines, and
2. the points on the curve where the tangent lines are vertical.

If you feel lucky, try solving this problem without Calculus by completing the square.

Problem 120. Find the equations for the two lines that pass through the origin and are tangent to the curve defined by $x = y^2 + 1$. Illustrate with a graph of the curve and tangent lines.

We proved the Power Rule only for functions of the form $f(x) = x^n$ where n is an integer. However, we have been using it where n was any number, for example $n = 1/2$ in the case where $f(x) = \sqrt{x}$. We will now use the Chain Rule, Implicit and Logarithmic Differentiation to prove that the Power Rule is valid for *rational* exponents.

Problem 121. Use implicit differentiation to find y' if $y = x^{4/3}$.

Problem 122. Suppose $\frac{p}{q}$ is a rational number. Show that the derivative of the function $f(x) = x^{p/q}$ is $f'(x) = \frac{p}{q}x^{\frac{p}{q}-1}$.

Problem 123. Find the second derivative, y'' , for the curve $x^3 + y^3 = 1$.

Problem 124. Use implicit differentiation to find the equation of the line tangent to the curve $x^{2/3} - y^{2/3} - y = 4$ at the point $(8, 0)$.

Chapter 3

Applications of Derivatives

“I am always doing that which I can not do, in order that I may learn how to do it.” - Pablo Picasso

Now that we have a solid understanding of what derivatives are and how to compute them for common functions, let's look at some applications.

3.1 Linear Approximations

Suppose we have a function f and a point $(p, f(p))$ on f . Suppose L is the tangent line to f at that point. L is the *best linear approximation* to f at $(p, f(p))$ since $f(p) = L(p)$ and $f'(p) = L'(p)$ and no line can do better than agreeing with f both in the y-coordinate and the derivative. Therefore the tangent line to f at p is often called the best linear approximation or the *linearization* of f at p .

Problem 125. Find and sketch the linearization of the function at the indicated point.

1. $f(x) = 3x^2 - x + 2$ at $x = -2$.
2. $g(x) = \sin(x)$ at $x = \pi/4$.

Problem 126. Suppose f is a differentiable function and $(p, f(p))$ is a point on f . Find the equation of the tangent line to f at the point $(p, f(p))$.

Problem 127. Suppose $f(x) = x^3$. Find a parabola $p(x) = ax^2 + bx + c$ so that $f(1) = p(1)$ and $f'(1) = p'(1)$ and $f''(1) = p''(1)$. We would call p the best quadratic approximation to f at $(1, f(1))$.

Example 11. Approximating changes using linear approximations.

Suppose we have an ice cube floating in space that is expanding as cosmic dust forms on its exterior. Suppose the cube has an initial side length of $s = 3$ inches and we wish to approximate how much the volume ($V = s^3$) will change as the length of the side increases by .1 inch. Of course, we can compute the change in volume directly. But we can also approximate it using the fact that the derivative tells us the rate of change of the volume. Recall that

$$V'(3) \approx \frac{V(3.1) - V(3)}{3.1 - 3}.$$

If we think of the change in volume as $\Delta V = V(3.1) - V(3)$ and the change in side length as $\Delta s = 3.1 - 3$, then

$$V'(3) \approx \frac{\Delta V}{\Delta s}.$$

So,

$$\Delta V \approx V'(s)\Delta s = 3(3)^2 \cdot .1 = 2.7$$

and we expect the volume to change by this amount. In a case where you knew the rate of change of the volume, but did not know a formula for volume you could use this to estimate the change in volume over a small change in side length.

Problem 128. Compute the exact change in volume of the cube from the previous discussion and compare to our estimate.

Problem 129. Suppose the radius of a circle C is to be increased from an initial value of 10 by an amount $\Delta r = .01$. Estimate the corresponding increase in the circle's area ($A = \pi r^2$) and compare it to the exact change.

Problem 130. The surface area of a sphere ($S = 4\pi r^2$) is expanding as air is pumped in. Approximate the change in the surface area as the radius expands from $r = 2.5$ to $r = 2.6$ and compare this to the exact change.

3.2 Limits Involving Infinity and Asymptotes

We have already considered limits of the form $\lim_{x \rightarrow a} f(x) = L$ where both a and L were real numbers. Now we wish to consider the possibilities where a or L or both are $\pm\infty$. To determine the value for $\lim_{x \rightarrow \infty} f(x)$ is essentially to ask what is happening to the values of $f(x)$ as x increases without bound. This will help identify horizontal asymptotes. When we see limits of the form $\lim_{x \rightarrow a} f(x) = \infty$, where the values of $f(x)$ increase without bound as x approaches the number a , we see that $x = a$ must be a vertical asymptote. Our definition for asymptote is an informal one, like our definition for limits, because we don't give a mathematically valid explanation of what the word "approaches" means. Still, it will serve our purposes. A more precise definition might appear in a course titled "Real Analysis."

Definition 18. An **asymptote** of a function f is any curve (or line) that the graph of f approaches. A **horizontal asymptote** of f is a horizontal line that the graph of f approaches. A **vertical asymptote** of f is a vertical line that the graph of f approaches.

Problem 131. Graph $f(x) = \frac{3x}{x-2}$. What number does $f(x)$ approach as x becomes large and positive? As x becomes large and negative? What happens to the values of $f(x)$ as x approaches 2 from the left? From the right?

Definition 19. Intuitive Definition We say that $\lim_{x \rightarrow +\infty} f(x) = L$ provided that as x becomes arbitrarily large and positive, the values of $f(x)$ become arbitrarily closer to L .

Definition 20. Another Intuitive Definition We say that $\lim_{x \rightarrow \infty} f(x) = \infty$ provided that as x becomes arbitrarily large and positive the values of $f(x)$ become arbitrarily large and positive.

For the problems in this section, you may justify your answer using a graph or some algebra or limit tables or simply a sentence defending your answer, such as "the denominator gets larger and larger and the numerator is constant, so the limit must be zero."

Problem 132. Evaluate each of the following limits.

$$1. \lim_{x \rightarrow \infty} \frac{2x}{5x+3}$$

$$2. \lim_{x \rightarrow \infty} x^2 + 5$$

$$3. \lim_{t \rightarrow \infty} \frac{4t + \cos(t)}{t}$$

Problem 133. Evaluate each of the following limits or state why they do not exist.

$$1. \lim_{t \rightarrow -\infty} 2t^3 + 6$$

$$2. \lim_{z \rightarrow \infty} \frac{\sin(z) + 7z}{1 - z^2}$$

$$3. \lim_{\alpha \rightarrow -\infty} 3\sin(\alpha)$$

Here are precise restatements of our intuitive definitions.

Definition 21. Precise Restatement of Definition 19 Let L be a real number. We say that $\lim_{x \rightarrow \infty} f(x) = L$ provided for each positive real number ε there is a positive real number M such that for every $x > M$, we have $|f(x) - L| < \varepsilon$.

Definition 22. Precise Restatement of Definition 20 We say that $\lim_{x \rightarrow \infty} f(x) = \infty$ provided for each positive real number N there is a positive real number M such that for every $x > M$, we have $f(x) > N$.

Problem 134. Based on the definitions above, write an intuitive definition and a precise definition for $\lim_{x \rightarrow \infty} f(x) = -\infty$

Problem 135. Suppose k is some real number and P be the second-degree polynomial $P(x) = kx^2$. Does the $\lim_{x \rightarrow \infty} P(x)$ depend on the value of k ? What is $\lim_{x \rightarrow \infty} P(x)$? What is $\lim_{x \rightarrow -\infty} P(x)$? Suppose $C(x) = kx^3$. What is $\lim_{x \rightarrow \infty} C(x)$? What is $\lim_{x \rightarrow -\infty} C(x)$?

Definition 23. Yet Another Intuitive Definition We say that $\lim_{x \rightarrow a} f(x) = \infty$ if when x approaches a , the values of $f(x)$ become arbitrarily large.

The next definition makes this idea precise.

Definition 24. Precise Restatement of Definition 23 Let a be a real number. We say that $\lim_{x \rightarrow a} f(x) = \infty$ provided for each positive real number N there is positive number ε so that for all x satisfying $|x - a| < \varepsilon$, we have $f(x) > N$

Problem 136. Evaluate each of the following limits. Graph each function; include any vertical asymptotes.

$$1. \lim_{t \rightarrow -2} \frac{3}{(2+t)^2}$$

$$2. \lim_{x \rightarrow 1^+} \frac{x^2 - 5x + 6}{x^3 - 1}$$

Problem 137. Evaluate each of the following limits.

$$1. \lim_{x \rightarrow 0^-} \frac{1}{x} \text{ and } \lim_{x \rightarrow 0^+} \frac{1}{x}$$

$$2. \lim_{x \rightarrow -4^-} \frac{x-1}{x+4} \text{ and } \lim_{x \rightarrow -4^+} \frac{x-1}{x+4}$$

$$3. \lim_{x \rightarrow 0} \sin\left(\frac{1}{x}\right)$$

Problem 138. Find the asymptotes of each function and use them to sketch a rough graph of each.

$$1. G(x) = \frac{x-1}{x+4}$$

$$2. F(x) = \frac{3x}{5x-1}$$

$$3. g(x) = e^{-x} - 2$$

$$4. h(z) = -\frac{2z + \sin(z)}{z}$$

Problem 139. Write a statement that explains the relationship between limits and vertical asymptotes. Write a statement that explains the relationship between limits and horizontal asymptotes.

3.3 Graphing

Prior to an understanding of derivatives, graphing was largely a matter of:

1. plotting points,
2. memorizing the basic shapes of certain types of functions, and
3. trusting the graphs generated by technology.

With algebra and differentiation at our disposal, we can make a precise science out of graphing that will give more information than even our graphing software yields.

Example 12. I will work a few of these examples to provide intuition for the upcoming definitions.

$$1. p(x) = x^4 - 13x^2 + 36$$

$$2. q(x) = \frac{3}{x^2 + 1}$$

$$3. f(x) = 2x^{5/3} - 5x^{4/3}$$

$$4. r(x) = \frac{x^2 - 1}{2x + 1}$$

$$5. F(t) = t^{2/3} - 3$$

$$6. h(x) = \frac{3x^2}{x^2 + 1}$$

We list several questions that you should ask when graphing a function.

Question 1. What are the intercepts and asymptotes of the function?

Recall that an x -intercept is a point where the function crosses the x -axis (i.e. $y = 0$) and a y -intercept is a point where the function crosses the y -axis (i.e. $x = 0$).

Recall that asymptotes (Definition 18) are functions (usually lines) that f approaches. Limits are helpful in understanding and locating asymptotes. Review Problem 138 and Problem 139 when studying asymptotes.

Problem 140. List any intercepts and asymptotes for:

$$1. f(x) = \frac{x}{(x-2)(x+3)}$$

$$2. E(x) = 3 - e^{-x}$$

$$3. g(x) = \frac{3x}{\sqrt{x^2 + 3}}$$

Question 2. What are the local maxima and minima of the function?

The peaks and the valleys are called local maxima and local minima, respectively.

Definition 25. If f is a function, then we say that the point $(c, f(c))$ is a **local maximum** for f if there is some interval, (a, b) , containing c so that $f(x) < f(c)$ for all $x \in (a, b)$ except $x = c$. Local minimum is defined similarly.

We have seen examples of local minima and maxima. Problem 56 had a minimum at a point where the derivative did not exist. In problem 74 we found the local maxima and minima by finding the values where the tangent lines were horizontal. Hence points on the graph where the derivative is either zero or undefined are important points.

Definition 26. If f is a function, then any value x in the domain of f where $f'(x)$ is zero or undefined is called a **critical value** of f . We refer to $(x, f(x))$ as a **critical point** of f .

Critical points give us candidates for local maximum or minimum, but they do not guarantee a maximum or minimum. The function $f(x) = x^3 + 2$ has $(0, 2)$ as a critical point, but this point is neither a minimum nor a maximum.

Problem 141. Find the critical values for these functions.

$$1. g(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 12x$$

$$2. f(x) = \sin(2x) \text{ on } [0, 2\pi]$$

$$3. s(x) = (x - 3)^{1/3}$$

Question 3. Where is the function increasing? Decreasing?

Determining where a function is increasing or decreasing is perhaps the most useful graphing tool, so we'll call it *the Fundamental Theorem of Graphing*.

Definition 27. We say that a function f is **increasing** if, for every pair of numbers x and y in the domain of f with $x < y$, we have $f(x) < f(y)$. Decreasing, non-increasing, and non-decreasing are defined similarly.

If the slopes of all tangent lines to the function f are positive over an interval (a, b) , then f is increasing on the interval (a, b) and if the slopes of all tangent lines to the function f are negative over an interval (a, b) , then f is decreasing on the interval (a, b) .

Theorem 9. The Fundamental Theorem of Graphing. If f is a function and is differentiable on the interval (a, b) and $f'(x) > 0$ for all x in (a, b) , then f is increasing on (a, b) .

Problem 142. These are the functions you found the critical values for in the last problem. List the intervals on which these functions are increasing by finding where $f' > 0$.

1. $g(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 - 12x$

2. $f(x) = \sin(2x)$ on $[0, 2\pi]$

Question 4. Where is the function concave up? Concave down?

Sketch $f(x) = x^3 - x$. The graph of f looks somewhat like a parabola opening downward on the interval $(-\infty, 0)$ and somewhat like a parabola opening upward on the interval $(0, \infty)$. We say that f is *concave down* on $(-\infty, 0)$ and *concave up* on $(0, \infty)$. The defining characteristic for concavity is that f is concave down when the slopes of the tangent lines are decreasing and f is concave up when the slopes are increasing.

Definition 28. We say that f is **concave up** on (a, b) if f' is increasing on (a, b) and **concave down** on (a, b) if f' is decreasing on (a, b) .

Applying the *Fundamental Theorem of Graphing*, f will be concave up when f' is increasing which will be when f'' is positive.

Question 5. What are the inflection points for the function?

Inflection points are points on the curve in which the function switches either from concave up to concave down, or from concave down to concave up.

Definition 29. We say that $(x, f(x))$ is an **inflection point** for f if the concavity of f changes sign at x .

Problem 143. Let $f(x) = \frac{6}{x^2 + 3}$. Find the values where $f'' = 0$. Find the intervals where f is concave up. List the inflection points.

Question 6. What is the least upper bound of the function? The greatest lower bound?

From the graph of $f(x) = \frac{6}{x^2 + 1}$ in the last problem we see that $(0, 6)$ is the highest point on the graph, but there is no lowest point since the graph looks like an infinite bell – it tends towards zero as x takes on large positive or large negative values. We say that 6 is the *least upper bound* of f and that f *attains this value* since $f(0) = 6$. We say that 0 is the *greatest lower bound* and that f *does not attain this value* since there is no number x so that $f(x) = 0$.

Definition 30. If there is a number M satisfying $f(x) \leq M$ for all x in the domain of f , then we call M an **upper bound** of f . If there is no number less than M that satisfies this property, then we call M the **least upper bound**.

Problem 144. For each function (1) find the least upper bound or state why it does not exist, and (2) find the greatest lower bound or state why it does not exist.

1. $E(x) = 3 - e^{-x}$

2. $f(x) = x^4 - x^2$

Problem 145. Choose one of the functions below and attempt to list as much information as possible. Use this information to help you sketch a very precise graph of the function. Do not expect to be able to list all of the information for each function. You won't be able to answer every question for every function. Graphing is the art of deciding which questions will best help you graph which functions.

- x -intercepts and y -intercepts (where $x = 0$ or $y = 0$)
- horizontal asymptotes and vertical asymptotes (lines the function approaches)
- critical points (places where f' is either zero or undefined)
- intervals over which each function is increasing (intervals where f' is positive)
- intervals over which each function is decreasing (intervals where f' is negative)
- intervals over which each function is concave up (intervals where f'' is positive)
- intervals over which each function is concave down (intervals where f'' is negative)
- least upper bound and greatest lower bound for each function (bounds for the range)

1. $f(x) = 2x^3 - 3x^2$

2. $p(x) = 3x^4 + 4x^3 - 12x^2$

3. $g(x) = (1 - x)^3$

4. $r(x) = 3 - x^{1/3}$

5. $F(t) = \frac{2t}{t+1}$

6. $h(x) = \frac{5}{x^2 - x - 6}$

7. $g(x) = \frac{x^3 + 1}{x}$

8. $s(x) = x - \cos(x)$

9. $T(x) = \tan(2x)$

Problem 146. Construct two functions f and g , each having a least upper bound of 1, so that f attains its least upper bound, but g does not.

3.4 The Theorems of Calculus

In this section we will study several important theorems: the Intermediate Value Theorem, the Extreme Value Theorem, Rolle's Theorem and the Mean Value Theorem. When a statement is of the form "if P then Q ", P is called the hypothesis of the theorem and Q is called the conclusion.

Definition 31. $C_{[a,b]}$ denotes the set of all functions that are continuous at every point in $[a, b]$ and $C^1_{[a,b]}$ denotes the set of all functions that are differentiable at every point in $[a, b]$. Thus, when we write $f \in C_{[a,b]}$ we mean that f is an element of $C^1_{[a,b]}$, so f is a function and f is differentiable at every point in $[a, b]$.

Theorem 10. Intermediate Value Theorem. If $f \in C_{[a,b]}$ and $f(a) < f(b)$ and y is a number between $f(a)$ and $f(b)$, then there is a number $x \in [a, b]$ such that $f(x) = y$.

Problem 147. Does $f(x) = \frac{x^2 - 1}{x - 1}$ satisfy the hypothesis Intermediate Value Theorem on the interval $[-1, 3]$? Does f satisfy the conclusion?

Problem 148. Use the Intermediate Value Theorem with $f(x) = x^3 + 3x - 2$ to show that $x^3 + 3x - 2 = 0$ has a solution on the interval $[0, 1]$.

Problem 149. Does $t^3 \cos(t) + 6\sin^5(t) - 3 = 2$ have a solution on the interval $[0, 2\pi]$?

Theorem 11. Extreme Value Theorem. If $f \in C_{[a,b]}$, then there are numbers c and d in $[a, b]$ so that $f(c) \leq f(x)$ for every x in $[a, b]$ and $f(d) \geq f(x)$ for every x in $[a, b]$.

The Extreme Value Theorem and the Intermediate Value Theorem together give us the important result that if $f \in C_{[a,b]}$ then $f([a, b]) = [m, M]$ where m and M are the absolute minimum and maximum of f , respectively.

Problem 150. Find the maximum and minimum of each function on the indicated interval.

1. $f(x) = x^2 - 9$ on $[2, 3]$
2. $f(x) = e^x$ on $[2, 5]$
3. $f(x) = x^3 - x$ on $[-1, 5]$
4. $f(x) = \frac{4\sqrt{x}}{x^2 + 3}$ on $[0, 4]$

Theorem 12. Rolle's Theorem. If $f \in C^1_{[a,b]}$ and $f(a) = f(b)$, then there is a point c in the interval (a, b) such that $f'(c) = 0$.

Sort-of-Proof. We have seen that f has an absolute minimum and an absolute maximum on the closed interval $[a, b]$. Suppose an absolute minimum of f occurs at x_m and an absolute maximum of f occurs at x_M ; then $f(x_m) \leq f(x) \leq f(x_M)$ for each x in $[a, b]$. If x_m is neither a nor b , then $f'(x_m) = 0$ since it is an absolute minimum. Similarly, if x_M is neither a nor b , $f'(x_M) = 0$ since it is an absolute maximum. Otherwise, we assume that $x_m = a$ and $x_M = b$. Then we have $f(x_m) \leq f(x) \leq f(x_M)$ for every x in $[a, b]$ which implies that $f(x_m) = f(x) = f(x_M)$ for every x in $[a, b]$. Therefore, $f'(x) = 0$ for all x in $[a, b]$. The same argument applies if $x_m = b$ and $x_M = a$. *q.e.d.*

Problem 151. In each case, determine if the function and the given interval $[a, b]$ satisfy the hypothesis of Rolle's Theorem. If so, find all points c in (a, b) such that $f'(c) = 0$.

1. $f(x) = 3x^2 - 3$ on $[-3, 3]$
2. $F(t) = 6t^2 - t - 1$ on $[-4, 5]$
3. $g(\theta) = \cos(\theta)$ on $[-\pi, 3\pi]$

Theorem 13. Mean Value Theorem. If $f \in C^1_{[a,b]}$, and then there is a point c in the interval (a, b) such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

Problem 152. In each case, determine if the function and the given interval $[a, b]$ satisfy the hypothesis of the Mean Value Theorem. If so, find all points c in (a, b) such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

1. $f(x) = 3x^2 - 3$ on $[-1, 3]$
2. $F(t) = 6t^2 - t - 1$ on $[-4, 5]$
3. $h(\beta) = \sin(2\beta)$ on $[-\frac{\pi}{4}, \pi]$
4. $H(y) = 5y^{\frac{2}{3}}$ on $[-2, 2]$

We have already discussed this theorem in an applied context. If we average 80 miles per hour between Houston and St. Louis, then at some point we must have had a velocity of 80 miles per hour. In our applied problem, f represented the distance traveled, a the starting time, b the ending time, f' the velocity, and c the time that must exist when we were traveling at 80 miles per hour.

Problem 153. Suppose $f(t)$ is the distance in feet that Ted has traveled down a road at time t in seconds. Suppose $f(10) = 0$ and $f(20) = 880$. Show that Ted traveled at 60 miles per hour at some time during the 10 second period.

Problem 154. A car is stopped at a toll booth. 18 minutes later it is clocked 18 miles away traveling at 60 miles per hour. Sketch a graph that might represent the car's speed from time $t = 0$ to $t = 18$ minutes. Then use the Mean Value Theorem to prove that the car exceeded 60 miles per hour.

The next problem is a very slight modification of a problem that I shamelessly stole from my friend and colleague, Brian Loft!

Problem 155. The first two toll stations on the Hardy Toll Road are 8 miles apart. Dr. Mahavier's EasyPass says it took 6 minutes to get from one to the other on a Sunday drive last week. A few days later, a ticket came in the mail for exceeding the 75 mph speed limit. Can he fight this ticket? Does Dr. Mahavier leave his GPS on in his phone while driving? Have you read "1984?"

Example 13. L'Hôpital's Rule. Consider these three simple limits and discuss the implications.

1. $\lim_{x \rightarrow \infty} \frac{x^2}{x} = \infty$
2. $\lim_{x \rightarrow \infty} \frac{3x}{x} = 3$
3. $\lim_{x \rightarrow \infty} \frac{x}{x^2} = 0$

Therefore if the top and the bottom of a rational function both tend to infinity, then the limit can be 0, ∞ , or any real number! We call such limits *indeterminate forms* and these were three examples of the indeterminate form $\frac{\infty}{\infty}$ since both the numerator and the denominator tended to infinity. Here is the formal definition.

Definition 32. If f and g are two functions such that $\lim_{x \rightarrow c} f(x) = \infty$ and $\lim_{x \rightarrow c} g(x) = \infty$, then the function $\frac{f}{g}$ is said to have the **indeterminate form** $\frac{\infty}{\infty}$ at c .

There are formal definitions for other indeterminate forms, but I don't think we need them to proceed. The list of indeterminate forms is:

$$\frac{0}{0}, \frac{\infty}{\infty}, 0 \cdot \infty, 0^0, 0^\infty, \infty^0, 1^\infty \text{ and } \infty - \infty.$$

One tool that is helpful in evaluating indeterminate forms follows.

Theorem 14. L'Hôpital's Rule. Suppose f and g are differentiable functions on some interval (a, b) and c is a number in that interval. If $\lim_{x \rightarrow c} f(x) = 0$ and $\lim_{x \rightarrow c} g(x) = 0$ and $g'(x) \neq 0$ for $x \neq c$ and $\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$ exists, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$.

We won't state it formally, but you may also use L'Hôpital's Rule for the indeterminate form $\frac{0}{0}$.

Problem 156. Evaluate each of the following limits using L'Hôpital's rule. For each problem, is there another way you know how to do it?

$$1. \lim_{x \rightarrow 1} \frac{x^2 - 2x + 1}{x^2 - 1}$$

$$2. \lim_{x \rightarrow 0} \frac{\sin(x)}{x}$$

$$3. \lim_{x \rightarrow 0} \frac{6^x - 1}{x^2 + 2x}$$

$$4. \lim_{x \rightarrow -3} \frac{3x - \frac{2}{x}}{x^2 - x - 20}$$

Problem 157. Evaluate each of the following limits if they exist. If they do not exist, why not?

$$1. \lim_{x \rightarrow \infty} \frac{2x}{5x + 3}$$

$$2. \lim_{x \rightarrow \infty} x^2 + 5$$

$$3. \lim_{t \rightarrow \infty} \frac{4t + \cos(t)}{t} \text{ (Try it with, and without, L'Hôpital's rule. Read L'Hôpital's rule carefully.)}$$

$$4. \lim_{x \rightarrow \infty} \frac{1 - 4x + 3x^2 + x^5}{x^2 + x - 2}$$

$$5. \lim_{x \rightarrow 0^+} \frac{x}{\ln(x)}$$

$$6. \lim_{x \rightarrow 0^+} \sin^{-1}(3x) \cdot \csc(2x)$$

$$7. \lim_{x \rightarrow \infty} (\sqrt{x^2 + 1} - x)$$

Problem 158. Use common denominators and L'Hôpital's Rules to compute $\lim_{x \rightarrow 0} \left(\frac{1}{x} - \frac{1}{\sin(x)} \right)$.

We can use properties of the natural log to resolve limits of the forms, 0^0 , ∞^0 and 1^∞ .

Example 14. Compute $\lim_{x \rightarrow \infty} x^{\frac{1}{x}}$.

1. Suppose the answer is L : $L = \lim_{x \rightarrow \infty} x^{\frac{1}{x}}$.
2. Apply the natural log to both sides: $\ln(L) = \ln(\lim_{x \rightarrow \infty} x^{\frac{1}{x}})$.
3. Swap the limit and the $\ln$: $\ln(L) = \lim_{x \rightarrow \infty} \ln(x^{\frac{1}{x}})$. Why is this legal?
4. Do some algebra and compute the limit: $\ln(L) = \lim_{x \rightarrow \infty} \frac{1}{x} \ln x = \lim_{x \rightarrow \infty} \frac{\ln(x)}{x} = \lim_{x \rightarrow \infty} \frac{1}{x} = 0$.
5. Solve for L : $\ln(L) = 0$ so $L = e^0 = 1$.

Problem 159. Evaluate these using the clever trick just described.

1. $\lim_{x \rightarrow 0^+} x^{x^2}$
2. $\lim_{x \rightarrow \infty} (1 + \frac{1}{x})^x$

3.5 Maxima and Minima

Many problems from industry boil down to trying to find the maximum or minimum of a function. In the example that follows, if a mathematician can find a way to make a can cheaper than the competition, then the company will save money and the mathematician will get a raise. One way to save money would be to determine if the dimensions of the cans can be changed to hold the same amount of fluid, but use less material.

Example 15. *TedCo Bottling Company wants to find a way to create a can of Big Ted that has 355cc of soda but uses less material than a standard can. This is called a max/min problem because we want to minimize the surface area while holding the volume constant.*

The surface area of the can is given by

$$A = 2\pi r^2 + 2\pi rh$$

where r is the radius and h is the height in centimeters. The volume is given by

$$V = \pi r^2 h$$

Since the volume is 355 cc, we have

$$355 = \pi r^2 h$$

so we can eliminate h in the surface area equation by solving the volume equation for h and plugging in.

$$h = \frac{355}{\pi r^2}$$

Therefore

$$\begin{aligned} A(r) &= 2\pi r^2 + 2\pi r h \\ &= 2\pi r^2 + 2\pi r \frac{355}{\pi r^2} \end{aligned}$$

$$= 2(\pi r^2 + 355/r)$$

So, all we need is to find the radius that minimizes the function A . Taking the derivative of A and setting it equal to 0 we have:

$$r = \sqrt[3]{\frac{355}{2\pi}} \approx 3.837$$

We can solve for h by substituting this back into

$$h = \frac{355}{\pi r^2} \approx 7.67$$

Observing that h and r are in centimeters, how does this compare to a standard Coke can? If we note that

$$h = \frac{355}{\pi r^2} = \frac{2}{r^2} \cdot \frac{355}{2\pi} = \frac{2r^3}{r^2} = 2r,$$

then we see that this is an unusual sized can. Why don't the rest of the canning companies follow this material saving model?

Problem 160. A rock is thrown vertically. Its distance d in feet from the ground at time t seconds later is given by $d = -16t^2 + 640t$. How high will the rock travel before falling back down to the ground? What will its impact velocity be?

Problem 161. A rectangular field is to be bounded by a fence on three sides and by a straight section of lake on the fourth side. Find the dimensions of the field with maximum area that can be enclosed with M feet of fence.

Problem 162. The carp population in Table Rock Lake is growing according to the growth function $y = 2000/(1 + 49e^{-.3t})$, where y is the number of carp present after t months. Graph this function. What is the maximum number of carp that the pond will hold? At what time is the population of carp increasing most rapidly?

Problem 163. A rectangular plot of land is to be fenced, using two kinds of fencing. Two opposite sides will use heavy-duty fencing selling for \$4 a foot, while the remaining two sides will use standard fencing selling for \$3 a foot. What are the dimensions of the plot of greatest area that can be fenced with M dollars?

Problem 164. You are in your damaged dune buggy in a desert three miles due south of milepost 73 on Route 66, a highway running East and West. The nearest water is on this (presumably straight) highway and is located at milepost 79. Your dune buggy can travel 4 miles per hour on the highway but only 3 miles per hour in the sand. You figure that you have 2 hours and 10 minutes left before you die of thirst. Assuming that you can take any path from your present position to the water, how long will it take you to get to the water and what does your future look like?

Problem 165. A square sheet of cardboard with side length of M inches is used to make an open box by cutting squares of equal size from each of the four corners and folding up the sides. What size squares should be cut to obtain a box with largest possible volume ($V = \text{length} \times \text{width} \times \text{height}$)?

Problem 166. Geotech Industries owns an oil rig 12 miles off the shore of Galveston. This rig needs to be connected to the closest refinery which is 20 miles down the shore from the rig. If underwater pipe costs \$500,000 per mile and above-ground pipe costs \$300,000 per mile, the company would like to know which combination of the two will cost Geotech the least amount of money. Find a function that may be minimized to find this cost. (This problem also stolen shamelessly from Brian Loft.)

Problem 167. Find the dimensions of the rectangle with maximum area that can be inscribed in a circle of radius M .

3.6 Related Rates

Often two quantities are related and the rate at which each changes depends on the rate at which the other changes. Think of a turbo-charged car. You may push the accelerator to the floor at a constant rate. While the rate of acceleration of the car is related, it will not be constant because as the turbo kicks in the car will accelerate at a much faster rate even though you are still pushing the pedal down at a constant rate.

Example 16. A 20 foot boat hook is leaning against a wall and the base is sliding away from the wall at the rate of 5 feet per minute. This moves the top of the hook down the wall. How fast is the top moving when the base is 8 feet from the wall? How fast is the top moving at the instant when it strikes the floor?

Let

- $x(t)$ = the distance from the base of the hook to the wall at time t and
- $y(t)$ = the distance from the top of the hook to the floor at time t .

So

- $x'(t)$ = the velocity at which the base of the hook is moving and
- $y'(t)$ = the velocity at which the top of the hook is moving.

Note that x and y are related by the Pythagorean theorem,

$$x^2(t) + y^2(t) = 20^2.$$

Using implicit differentiation yields,

$$2x(t)x'(t) + 2y(t)y'(t) = 0.$$

If we consider t^* to be the time when the base of the pole is 8 feet from the wall, then $x(t^*) = 8$ and $x'(t^*) = 5$. Therefore, we have

$$2x(t^*)x'(t^*) + 2y(t^*)y'(t^*) = 0.$$

And solving for $y'(t^*)$ we have

$$y'(t^*) = \frac{-x(t^*)x'(t^*)}{y(t^*)} = \frac{-40}{\sqrt{400 - 64}} = -10\sqrt{21} \approx -2.2.$$

Why is $y'(t^*)$ negative?

Problem 168. A baseball diamond is a square with sides 90 feet long. A runner travels from first base to second base at 30 ft/sec. How fast is the runner's distance from home plate changing when the runner is 50 feet from first base?

Problem 169. Two planes leave an airport simultaneously. One travels north at 300 mph and the other travels west at 400 mph. How fast is the distance between them changing after 10 minutes?

Problem 170. If the temperature of a gas is held constant, then Boyle's Law guarantees that the pressure P and the volume V of the gas satisfy the equation $PV = c$, where c is a constant. Suppose the volume is increasing at the rate of 15 cubic inches per second. How fast is the pressure decreasing when the pressure is 125 pounds per square inch and the volume is 25 cubic inches?

Problem 171. Sand pouring from a chute at the rate of 10 cubic feet per minute forms a conical pile whose height is always one third of its radius. How fast is the height of the pile increasing at the instant when the pile is 5 feet high?

Problem 172. A balloon rises vertically from a point that is 150 feet from an observer at ground level. The observer notes that the angle of elevation is increasing at the rate of 15 degrees per second when the angle of elevation is 60 degrees. Find the speed of the balloon at this instant.

Problem 173. A snowball ($V = \frac{4}{3}\pi r^3$) melts at a rate that, at each instant, is proportional to the snowball's surface area ($A = 4\pi r^2$). Assuming that the snowball is always spherical, show that the radius decreases at a constant rate.

Problem 174. Herman is cleaning the exterior of a glass building with a squeegee. The base of the 10 foot long squeegee handle, which is resting on the ground, makes an angle of A with the horizontal. The top of the squeegee is x feet above the ground on the building. Herman pushes the bottom of the squeegee handle toward the wall. Find the rate at which x changes with respect to A when $A = 60$ degrees. Express the answer in feet per degree.

Chapter 4

Integrals

“We have to believe - before students can believe - that hard work pays off, that effort matters, that success depends not on your genes but on your sweat. We must convince students that they can do it when they have teachers who insist on quality work and give them extra remedial help when needed.” - Gene Bottoms

Integrals are used throughout engineering and industry to solve a wide variety of applied problems such as: the amount of work necessary to lift a 50 ton satellite, the total pressure on the Hoover dam, the probability that you will live to an age between 69 and 71, the volume formulae for various objects, the weight of objects that do not have uniform density, the work required to pump water out of a 55 gallon drum, and the area of an oil spill.

4.1 Anti-differentiation and the Indefinite Integral

We might denote the derivative of the function $f(x) = x^2$ as any of:

$$f', \frac{df}{dx}, (x^2)', \text{ or } \frac{d}{dx}(x^2).$$

If f is a function, then any function with derivative f is called an **anti-derivative** of f . For the *anti-derivative* of f we may write any of:

$$\int f, \int f(x) dx, \text{ or } \int x^2 dx.$$

Any function which has one anti-derivative has infinitely many anti-derivatives. To indicate all anti-derivatives of $f(x) = x^2$ we will write:

$$\int f(x) dx = \int x^2 dx = \frac{1}{3}x^3 + c, \quad c \in \mathbb{R}.$$

We would read this notation as “*the indefinite integral (or anti-derivative) of x^2 with respect to x is $\frac{1}{3}x^3 + c$ where c represents an arbitrary constant.*” It is misleading that mathematicians speak of “the indefinite integral” when in fact it is not a single function, but a class of functions. You may interpret $\int$ and “ dx ” as representing the beginning and end of the function we wish to find the anti-derivative of. We call this process *integrating* the function. The function to be integrated is called the *integrand*.

Problem 175. Evaluate the indicated indefinite integrals and then state the Sum Rule for Indefinite Integrals based on your results.

1. $\int 3x^4 + 5 \, dx$

2. $\int 3x^4 \, dx + \int 5 \, dx$

Problem 176. Evaluate the indicated indefinite integrals and then state the Constant Multiple Rule for Indefinite Integrals based on your results.

1. $\int 5 \sin(x) \, dx$

2. $5 \int \sin(x) \, dx$

Problem 177. Compute the following indefinite integrals.

1. $\int \frac{x+x^3}{x^5} \, dx$

2. $\int t(t^{1/2} + 8) \, dt$

3. $\int (s^3 + s)^3 \, ds$

4. $\int 3 \cos(x) - 2 \sin(4x) \, dx$

Like differentiation, anti-differentiation becomes more challenging when the *Chain Rule* appears in the problem.

Example 17. Compute $\int x^2(x^3 + 10)^9 \, dx$.

Our first guess will be $g(x) = (x^3 + 10)^{10}$. Then $g'(x) = 10(x^3 + 10)^9 \cdot 3x^2 = 30x^2(x^3 + 10)^9$. Our answer is correct except for the 30 in front. We take care of this by modifying our first guess to $g(x) = \frac{1}{30}(x^3 + 10)^{10}$. Now $g'(x) = \frac{1}{30} \cdot 10(x^3 + 10)^9 \cdot 3x^2 = x^2(x^3 + 10)^9$ and so our second guess is one anti-derivative of the integrand. Adding “+ c” to the end yields all possible anti-derivatives.

Problem 178. Compute the following indefinite integrals.

1. $\int (x+2)(3x^2 + 12x - 5)^{121} \, dx$

2. $\int (x+2)\sqrt[4]{3x^2 + 12x - 5} \, dx$

3. $\int p^2 \sqrt[3]{p^3 + 1} \, dp$

4. $\int \frac{3 \cos(\sqrt{t})}{\sqrt{t}} \, dt$

5. $\int \frac{x}{3x^2 + 4} \, dx$

Problem 179. *These look quite similar. Try both. What's the difference?*

$$1. \int \frac{\sin(t)}{\cos^2(t)} dt$$

$$2. \int \frac{\sin(t)}{\cos(t^2)} dt$$

4.2 Riemann Sums and Definite Integrals

You know how to compute the areas of certain shapes (rectangles, triangles, etc.) using formulas, but what about unusual shapes like the area trapped under a curve and above the x -axis or the area trapped between two curves? If we have a function that is always above the x -axis on the interval $[a, b]$, then Riemann sums give us a way to compute the area that is between the vertical lines $x = a$ and $x = b$, above the x -axis, and under the curve.

Example 18. *Determine the area under the curve $f(x) = x^2 + 1$, above the x -axis, and between the vertical lines $x = 0$ and $x = 2$.*

Graph f and subdivide $[0, 2]$ into 4 intervals each of width $\frac{1}{2}$ and with bases,

$$\left[0, \frac{1}{2}\right], \left[\frac{1}{2}, 1\right], \left[1, \frac{3}{2}\right], \text{ and } \left[\frac{3}{2}, 2\right].$$

Each has width $\frac{1}{2}$. These intervals are the bases of 4 rectangles. Make the height of each rectangle the value of f at the right end point of the base of that rectangle. Let's call the sum of the areas of these four rectangles U_4 . (If you are unfamiliar with summation notation, $\sum$, see Appendix 15.5 for a review.)

The sum of these rectangles is given by:

$$\begin{aligned} U_4 &= \frac{1}{2} \left(\left(\frac{1}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{2}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{3}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{4}{2}\right)^2 + 1 \right) \\ &= \frac{1}{2} \left[\left(\left(\frac{1}{2}\right)^2 + 1 \right) + \left(\left(\frac{2}{2}\right)^2 + 1 \right) + \left(\left(\frac{3}{2}\right)^2 + 1 \right) + \left(\left(\frac{4}{2}\right)^2 + 1 \right) \right] \\ &= \frac{1}{2} \sum_{i=1}^4 \left(\left(\frac{i}{2}\right)^2 + 1 \right) = \frac{23}{4} = 5.75 \end{aligned}$$

This estimate is too large because each rectangle we chose had area larger than the portion of the curve it approximated. Therefore, we make an estimate that is too small and average the two! An estimate that would be too small would have four rectangles each with the same bases, but with height the value of f at the *left* endpoint of its base. Let's call the sum of the areas of these four rectangles L_4 .

$$\begin{aligned} L_4 &= \frac{1}{2} \left(\left(\frac{0}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{1}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{2}{2}\right)^2 + 1 \right) + \frac{1}{2} \left(\left(\frac{3}{2}\right)^2 + 1 \right) \\ &= \frac{1}{2} \left[\left(\left(\frac{0}{2}\right)^2 + 1 \right) + \left(\left(\frac{1}{2}\right)^2 + 1 \right) + \left(\left(\frac{2}{2}\right)^2 + 1 \right) + \left(\left(\frac{3}{2}\right)^2 + 1 \right) \right] \\ &= \frac{1}{2} \sum_{i=0}^3 \left(\left(\frac{i}{2}\right)^2 + 1 \right) = \frac{15}{4} = 3.75 \end{aligned}$$

What you have just computed are referred to as *upper* and *lower Riemann sums*, respectively. Our estimate based on averaging the two would yield

$$\frac{U_4 + L_4}{2} = 4.75.$$

Problem 180. Let $f(x) = x^2 + 1$. Divide the interval $[0, 2]$ into eight equal divisions,

$$\left\{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, \dots, \frac{7}{4}, 2\right\}$$

and compute the upper and lower Riemann sums of f .

Subdividing an interval in this way is called *partitioning* the interval and the sub-intervals created via a partition may be of differing widths as indicated in the next definition.

Definition 33. A **partition of the interval $[a, b]$** is a collection of points, $\{x_0, x_1, \dots, x_n\}$ satisfying, $a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b$.

Definition 34. If f is a function defined on the interval $[a, b]$ and $P = \{x_0, x_1, \dots, x_n\}$ is a partition of $[a, b]$, then a **Riemann sum of f over $[a, b]$** is defined by

$$R(P, f) = \sum_{i=1}^n (x_i - x_{i-1}) \cdot f(\hat{x}_i) \quad (4.1)$$

where $\hat{x}_i \in [x_{i-1}, x_i]$.

In the definition of Riemann sum, $\hat{x}_i$ can be *any* value in the interval $[x_{i-1}, x_i]$. But look back at our example. When we computed U_4 we always chose $\hat{x}_i$ so that $f(\hat{x}_i)$ was the *maximum* for f on this interval to assure that our approximation was more than the desired area. When $\hat{x}$ is chosen in this manner, we call our sum an **upper Riemann sum**. When we computed L_4 we always chose $\hat{x}_i \in [x_{i-1}, x_i]$ so that $f(\hat{x}_i)$ was the *minimum* for f on that interval to assure that our approximation was less than the desired area. When $\hat{x}$ is chosen in this manner, we call our sum a **lower Riemann sum**.

Problem 181. Compute upper and lower Riemann sums for $g(x) = x^2 + 3x$ on $[0, 1]$ using the partition, $P = \{0, \frac{1}{3}, \frac{2}{3}, 1\}$.

Problem 182. Compute upper and lower Riemann sums for $g(x) = -3x^3 + 3x$ over $[0, 1]$ using the partition, $P = \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$. The heights of our rectangles do not always occur at the endpoints of the sub-intervals of our partition.

Returning to $f(x) = x^2 + 1$ on $[0, 2]$, continue to subdivide our interval into n subdivisions of equal length, letting $n = 10, n = 100, n = 1000, \dots$ and labeling our partitions as $P_{10}, P_{100}, P_{1,000}, \dots$. If we label our lower sums corresponding to these partitions as $L_{10}, L_{100}, L_{1,000}, \dots$ and label our upper sums as $U_{10}, U_{100}, U_{1,000}, \dots$, then we would have

$$L_4 < L_{10} < L_{100} < L_{1,000} < \dots < U_{1,000} < U_{100} < U_{10} < U_4.$$

Furthermore, $L_{1,000}$ and $U_{1,000}$ should be very close together and very close the area we desire to compute.

By increasing the number of rectangles and taking the limit as the number of rectangles tends to infinity, we will determine the *exact* area. Our notation for this area is:

$$\text{Area} = \int_0^2 x^2 + 1 \, dx = \lim_{n \rightarrow \infty} U_n = \lim_{n \rightarrow \infty} L_n.$$

For our function, these two limits are equal, but there are functions where these limits are not equal. When they are equal, we say f is *integrable* over the interval. If they are not equal, we say that f is not integrable. We refer to $\int_0^2 x^2 + 1 \, dx$ as “the definite integral from 0 to 2 of $x^2 + 1$.”

Problem 183. Give an example of a function and an interval where the function will not be integrable over that interval.

We now consider the upper Riemann sum for $f(x) = x^2 + 1$, above the x -axis, and between the vertical lines $x = 0$ and $x = 2$ assuming n equal divisions of the interval $[0, 2]$. Our partition of n divisions is now,

$$P = \left\{0, \frac{2}{n}, \frac{4}{n}, \dots, \frac{2(n-1)}{n}, 2\right\},$$

and our upper Riemann sum U_n that approximates the area is given by:

$$\begin{aligned} U_n &= \frac{2}{n} \sum_{i=1}^n \left(\left(\frac{2i}{n} \right)^2 + 1 \right) \\ &= \frac{2}{n} \left[\sum_{i=1}^n \frac{4i^2}{n^2} + \sum_{i=1}^n 1 \right] \\ &= \frac{2}{n} \sum_{i=1}^n \frac{4i^2}{n^2} + \frac{2}{n} \sum_{i=1}^n 1 \\ &= \frac{8}{n^3} \sum_{i=1}^n i^2 + \frac{2}{n} \cdot n \text{ by a formula from Appendix 15.5} \\ &= \frac{8}{n^3} \frac{n(n+1)(2n+1)}{6} + 2 \\ &= \frac{4n(n+1)(2n+1)}{3n^3} + 2 \\ &= \frac{8n^3 + 12n^2 + 4n}{3n^3} + 2 \\ &= \frac{8n^3}{3n^3} + \frac{12n^2}{3n^3} + \frac{4n}{3n^3} + 2 \\ &= \frac{8}{3} + \frac{12}{3n} + \frac{4}{3n^2} + 2 \end{aligned} \tag{4.2}$$

Now take the limit of the Riemann sums as the number of divisions tends to infinity to get the *exact* area under the curve.

$$\begin{aligned} \text{Area} &= \int_0^2 x^2 + 1 \, dx \\ &= \lim_{n \rightarrow \infty} U_n \\ &= \lim_{n \rightarrow \infty} \frac{8}{3} + \frac{12}{3n} + \frac{4}{3n^2} + 2 \\ &= \lim_{n \rightarrow \infty} \frac{8}{3} + \lim_{n \rightarrow \infty} \frac{12}{3n} + \lim_{n \rightarrow \infty} \frac{4}{3n^2} + \lim_{n \rightarrow \infty} 2 \\ &= \frac{8}{3} + 2 \\ &= 4\frac{2}{3} \end{aligned} \tag{4.3}$$

Look back at your answers from Problem 180 (U_8 and L_8) and the solutions from our Example (U_4 and L_4). Just as we expect,

$$L_4 < L_8 < 4\frac{2}{3} < U_8 < U_4.$$

We denote the number $4\frac{2}{3}$ by $\int_0^2 x^2 + 1 \, dx$ and call this *the definite integral of $x^2 + 1$ from $x = 0$ to $x = 2$* . What we have just done motivates the following definition of the definite integral.

Definition 35. If f is a function defined on $[a, b]$ so that $\lim_{n \rightarrow \infty} L_n$ and $\lim_{n \rightarrow \infty} U_n$ both exist and are equal, then we say f is **integrable on $[a, b]$** and we define **the definite integral** by $\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} L_n$. We refer to a as the **lower limit of integration** and b as the **upper limit of integration**.

For each of the next few problems, you will require some of the summation formulas from Appendix 15.5.

Problem 184. Repeat this procedure, but compute the area using **lower** Riemann sums. That is, compute and simplify L_n (the lower Riemann sum of $f(x) = x^2 + 1$ on the interval $[0, 2]$ assuming n equal divisions) and then compute $\lim_{n \rightarrow \infty} L_n$.

Problem 185. Compute each integral using Definition 35 (Riemann sums).

1. $\int_0^1 2x \, dx$

2. $\int_0^1 x^3 \, dx$

3. $\int_0^2 2x^2 - 3 \, dx$

Problem 186. Compute the area trapped between the curves $y = x$ and $y = x^2$ using Definition 35.

Which functions are integrable? That is a question better-suited for a course called *Advanced Calculus* or *Real Analysis*, but let's give the quick-and-dirty answer without proof. By Definition 35 a function is integrable if a certain limit exists, but this can be difficult to compute. It can be shown that every differentiable function is continuous and every continuous function is integrable. Thus,

$$f \text{ is differentiable} \Rightarrow f \text{ is continuous} \Rightarrow f \text{ is integrable.}$$

The next two theorems state this formally.

Theorem 15. A Differentiability Theorem. If f is differentiable on $[a, b]$, then f is continuous on $[a, b]$.

Theorem 16. An Integrability Theorem. If f is continuous on $[a, b]$, then f is integrable on $[a, b]$.

We defined integrals in terms of Riemann sums and Riemann sums in terms of partitions of the interval. Therefore the integral $\int_a^b f(x) \, dx$ (see Definition 35) is only defined when $a < b$. The following definitions define definite integrals where the lower limit of integration is larger than the upper limit of integration or the two endpoints of integration are equal.

Definition 36. If a and b are numbers with $a < b$ and f is integrable on $[a, b]$ then:

1. $\int_b^a f(x) \, dx = - \int_a^b f(x) \, dx$ and

2. $\int_a^a f(x) \, dx = 0$.

The next theorem we discuss can be proved via Riemann sums, but we will leave that for your first course in *Real Analysis* as well. That's where the real fun begins.

Theorem 17. Sum Rule for Definite Integrals. If $c \in [a, b]$, then $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$.

Problem 187. Compute each of the following integrals using Definition 35 (Riemann Sums). This illustrates Theorem 17, the Sum Rule for definite integrals.

1. $\int_0^1 x^2 dx$

2. $\int_1^2 x^2 dx$

3. $\int_0^2 x^2 dx$

4.3 The Fundamental Theorem of Calculus

The *Fundamental Theorem of Calculus* (FTC) is one of the most applied theorems in all of calculus as it enables us to compute an integral *without* using Riemann sums.

To prove the FTC it will be helpful for us to understand the *Mean Value Theorem for Integrals*. This theorem holds for any continuous function f even though we state it here only for positive functions, that is, functions whose graphs are above the x -axis. This theorem says that if f is a positive function defined on the closed interval $[a, b]$, then there is a rectangle with base the closed interval $[a, b]$ and height $f(c)$ for some c in $[a, b]$ so that the area of this rectangle $f(c)(b - a)$ is equal to the area under the curve. The height of this rectangle, $f(c)$, is called the average value of the function over the interval. This is one of those theorems that is “obvious” once you draw a careful picture and follows from two powerful theorems that we did not prove, namely the Extreme Value Theorem and the Intermediate Value Theorem.

Theorem 18. The Mean Value Theorem for Integrals. If $f \in C_{[a,b]}$, then there exists a number $c \in [a, b]$ such that $\int_a^b f(x) dx = f(c)(b - a)$.

Problem 188. Let $f \in C_{[a,b]}$ with maximum M and minimum m . Use a graph to show that $m(b - a) \leq \int_a^b f(x) dx \leq M(b - a)$. Now apply the Intermediate Value Theorem to the function $g(x) = f(x)(b - a)$ to prove the Mean Value Theorem for Integrals.

Problem 189. Let $f(x) = 2x^2 - 3$ on the interval $[0, 2]$. Compute the value of c from the Mean Value Theorem for Integrals. You computed $\int_0^2 f(x) dx$ in Problem 185.

Definition 37. If $f \in C_{[a,b]}$, then the **average value** of f over $[a, b]$ is the number $\frac{1}{b - a} \int_a^b f(x) dx$.

Problem 190. Find the average value of the function over the given interval.

1. $g(x) = x^2 - 3x$ over $[0, 4]$

2. $f(x) = |x - 2| + 3$ over $[0, 3]$

Theorem 19. The Fundamental Theorem of Calculus.

1. Let $f \in C_{[a,b]}$ and let x be any number in $[a, b]$. If F is the function defined by

$$F(x) = \int_a^x f(t) dt,$$

then

$$F'(x) = f(x).$$

2. Let $f \in C_{[a,b]}$. If F is any function such that $F'(x) = f(x)$ for all x in $[a, b]$, then

$$\int_a^b f(x) dx = F(b) - F(a).$$

Proof of Part 1. Let $f \in C_{[a,b]}$ and define the function F so that $F(x)$ is the definite integral of f from a to x . That is,

$$F(x) = \int_a^x f(t) dt \text{ for every } x \in [a, b].$$

If F' exists, then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} [F(x+h) - F(x)] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt. \end{aligned}$$

Applying the *Mean Value Theorem for Integrals* at this point, we have

$$\frac{1}{h} \int_x^{x+h} f(t) dt = \frac{1}{h} (x+h - x) \cdot f(c)$$

for some value c that is in the interval $[x, x+h]$. Since this the interval depends on the value of h , we replace c by c_h to indicate that c depends on h . As $h \rightarrow 0^+$, we see that $x+h \rightarrow x$ so this interval will shrink to the point x . Since $x < c_h < x+h$ and $h \rightarrow 0^+$, we have that $c_h \rightarrow x$ by Theorem 3 (The Squeeze Theorem). This observation allows us to finish the proof.

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt \\ &= \lim_{h \rightarrow 0} \frac{1}{h} (x+h - x) \cdot f(c_h) \text{ where } c_h \in [x, x+h] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \cdot h \cdot f(c_h) \\ &= \lim_{h \rightarrow 0} f(c_h) \\ &= f(x) \text{ because } c_h \rightarrow x \text{ and } f \text{ is continuous.} \end{aligned}$$

q.e.d.

Sketch of Proof of Part 2. Let $f \in C_{[a,b]}$. Let

$$F(x) = \int_a^x f(t) dt \text{ for every } x \in [a, b].$$

By the previous proof, F is an anti-derivative of f .

$$F(b) - F(a) = \int_a^b f(t) dt - \int_a^a f(t) dt = \int_a^b f(t) dt - 0 = \int_a^b f(t) dt.$$

It might appear that we are done, but we have not shown that the theorem is true for *any* anti-derivative of f , rather we have only shown that it was true for anti-derivatives of the form $F(x) = \int_a^x f(t) dt$. Suppose we have some other anti-derivative, G . Then $G' = f = F'$, so $G' = F'$. If two functions have the same derivative then they must have the same shape. Therefore, either they are the same function or one function is just the other function slid up or down along the y-axis. It can be shown using the Mean Value Theorem if $F' = G'$, then $G(x) = F(x) + k$. Therefore, for any anti-derivative G we have,

$$G(b) - G(a) = (F(b) + k) - (F(a) + k) = F(b) - F(a) = \int_a^b f(t) dt.$$

q.e.d.

Problem 191. Use the second part of the fundamental theorem.

1. Compute $\int_1^3 (3x - 1) dx$. Verify your result using simple formulas for area.
2. Compute $\int_0^{2\pi} \cos^5(x) \sin(x) dx$.
3. Compute $\int_2^3 x(10 - x^2)^3 dx$.
4. Compute $\int_0^3 (x^2 + 2)e^{x^3 + 6x} dx$.

Problem 192. Use the first part of the fundamental theorem.

1. Compute F' where $F(x) = \int_1^x t^2 - 3t dt$
2. Compute F' where $F(x) = \int_5^x \frac{4}{t^4 + 1} dt$
3. Compute F' where $F(x) = \int_3^{x^2} t(t^2 - 4)^2 dt$

Problem 193. Let $f(t) = |t^2 - t|$ and let F be a function defined by $F(x) = \int_{-4}^x f(t) dt$ where $x \in [-4, 4]$. Answer the following questions.

1. What is the derivative of F ?
2. On what intervals is F increasing?
3. On what intervals is F decreasing?
4. On what intervals is F concave upward?
5. On what intervals is F concave downward?

4.4 Area, Volume, Arc length and Center of Mass

Definite integrals and the Fundamental Theorem of Calculus can be used to solve a variety of problems in geometry and physical science. In this section, the focus is on solving problems related to area, volume, and arc length.

Area Between Curves

Example 19. Discussion on using area between curves to approximate the volume of an oil spill, followed by computing the area between $y = x^2 + 1$ and $y = 2x + 1$.

Problem 194. Consider the curves $y = x^3$ and $y = x$.

1. Graph these curves and shade the regions bounded by these curves.
2. Set up and evaluate the definite integrals that represent the area of each of the shaded regions.

Problem 195. Use techniques in the previous problem to compute the areas of the regions bounded by these curves.

1. $y = x^2 + 4$ and $x + y = 6$
2. $x = y^2$ and $y = x - 2$
3. $x = \cos(y)$ and $x = 0$ from $y = 0$ to $y = \pi$

Problem 196. Sketch the area bounded by $y = x^2$, $y = 0$, and $x = 2$. Compute the area of this region by computing $\int_0^2 x^2 dx$. Now fill in the blanks to integrate this along the y -axis: $\int_0^2 x^2 dx = \int_0^4 \text{----} dy$.

Volumes of Solids

We now develop a technique that will allow us to compute volumes of various three dimensional objects. A *solid* is a three dimensional object along with its interior; think bowling ball, not balloon. When we computed areas, we began with Riemann sums to approximate the area and as the number of rectangles tended to infinity, the integral expression turned out to be nothing more than integrating over the height of the function at each point. That is,

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)(x_i - x_{i-1}) = \int_a^b f(x) dx.$$

Thus what we integrated was simply the height of the line segment from $(x, 0)$ to $(x, f(x))$. Intuitively, we might think of this as adding up infinitely many line lengths to get the area. Of course, the only *mathematically valid* way to think of this is as the limit of sums of areas of rectangles. The same idea applies to solids. To be precise, we need to slice the volume into n pieces and sum the volumes of each of the n pieces. Then we would take the limit as $n \rightarrow \infty$. Intuitively, we might think of slicing the solid into infinitely many cross sections and adding up the area of all these cross sections. The volume of a solid is the integral of the function that tells us the area of each cross section. Let's call this the *method of slicing*.

Problem 197. Sketch the region bounded by the lines $y = \frac{1}{2}x$, $y = 0$, and $x = 4$ and the solid cone obtained by rotating this region about the x -axis.

1. Draw a two dimensional slice of this solid parallel to the base with center at the point $(x, 0)$.
2. Find the function $A(x) = \dots$ that gives the area of this two dimensional slice.
3. Compute $\int_0^4 A(x) dx$ to determine the volume of your solid.

Problem 198. Sketch a pyramid whose base is a square of length 4 and whose height is 10.

1. Write a function $A(h) = \dots$ that gives the area of the slice that is parallel to the base and is at height h .
2. Compute $\int_0^{10} A(h) dh$ to determine the volume of the pyramid.

Problem 199. Compute the volume ($V = \frac{1}{3}\pi r^2 h$) of the cone with a circular base of radius r and a height of h .

Problem 200. Compute the volume ($V = \frac{1}{3}b^2 h$) of the pyramid with a square base of side length b and a height of h .

Problem 201. Compute the volume of the solid whose base is the disk $x^2 + y^2 \leq 4$ and whose cross sections are squares perpendicular to the diameter of the base through $(-2, 0)$ and $(2, 0)$. Can you sketch this?

The method of slicing can be used to find the volumes of *surfaces of revolution* which are solids created by taking a two dimensional region (an area) and rotating this region about a line to form a three dimensional solid. For example, take the unit circle and spin it about the x -axis and you get a sphere along with its interior. Or, take a circle of radius one centered at $(2, 0)$ and spin it around the y -axis and you get a donut.

Example 20. Find the volume obtained by rotating the region bounded by $f(x) = x^3 - x$ and $y = 0$ about the x -axis.

Sketching this, we get two solids. Looking at the portion where $x > 0$, we see something shaped like a lime with ends at $(0, 0)$ and $(1, 0)$. Choose a value for x between 0 and 1. If we think about slicing the lime parallel to the y -axis and through the point $(x, 0)$, then the cross section (slice) will be a circle and the area of this circle is $A = \pi r^2 = \pi (f(x))^2$ since the radius of the circle is $f(x)$. Therefore by the method of slicing, the volume of the solid S is given by

$$V = \pi \int_a^b [f(x)]^2 dx.$$

This method of finding the volume of revolution is called the **disk method** since intuitively we are adding up the area of an infinite number of disks.

Problem 202. Compute the volume of the “lime” in the preceding example.

There is nothing special about the x -axis.

Problem 203. Consider the region bounded by $y = x^3$, $x = 0$, and $y = 27$. Sketch the solid created by rotating this region about the y -axis. Compute the area function $A(y) = \dots$ that tells the area of one horizontal slice at height y (this would be a circle with center the point $(0, y)$). Now compute the volume of this solid of revolution, $V = \pi \int_0^{27} A(y) dy$.

Problem 204.

1. Sketch the region bounded by the curves $y = x^2 + 1$, $x = -1$, $x = 2$, and $y = -1$.
2. Sketch the solid of revolution generated by revolving this region about the line $y = -1$.
3. Let $-1 < x < 2$ and sketch the slice of the solid at x that is perpendicular to the line $y = -1$.
4. Determine the area of this slice, $A(x)$.
5. Integrate this area function from $x = -1$ to $x = 2$ to determine the volume of the solid.

Problem 205. Find the volume of the solid generated by rotating the region bounded by $x = 0$, $y = 6 - x$, and $y = 3$ around the x -axis.

Problem 206. Suppose a and b are positive numbers with $a < b$. Find the volume of the solid generated by rotating about the x -axis the region bounded by: $x = 0$, $y = x$, $y = a$, and $y = b$.

Problem 207. Find the volume of the solid generated by revolving about the y -axis the region bounded by the line $y = 2x$ and the curve $y = x^2$.

Now that we know this method of slicing up the surfaces, one might ask “Is this the only way we can slice up the surface to find the volume?” The answer is that there are lots of ways to slice up the surfaces. Let’s look at another way.

Problem 208. Find the volume obtained by rotating the region bounded by $f(x) = x - x^4$ and the x -axis about the x -axis.

Example 21. Find the volume obtained by rotating around the y -axis the region bounded by $f(x) = x - x^4$ and the x -axis.

To use the disk method, we would need to rewrite the function f which has as its independent variable, x , as a function g which has as its independent variable, y . Go ahead and try – if you can, you can skip the rest of this section! If you can’t, read on!

We will still be using the method of slicing, but we will use a “circular knife” like a cookie cutter so that instead of circular slices, we will have cylindrical slices. We will add up the surface area of each cylindrical slice, just as before we added up the areas of each circular slice.

Choose a value x between 0 and 1 and draw the line segment from $(x, 0)$ to $(x, f(x))$. Now rotate this line about the y -axis. Do you see that this is a “cylindrical slice” of your solid? If we consider all such slices with x ranging from 0 to 1, then integrating the function that gives the surface area of each slice will (again) yield the volume of our solid. But what exactly is the surface area of our slice? The radius of the cylinder is x and the height is $f(x)$ so the area must be $A(x) = 2\pi x f(x)$. And the volume will be the integral of this quantity,

$$V = 2\pi \int_0^1 x f(x) dx = 2\pi \int_0^1 x(x - x^4) dx = \frac{\pi}{3}.$$

This method is called the *shell method* or the *cylindrical shell method*.

Problem 209. Find the volume of the solid generated by revolving about the x axis the region bounded by the line $y = \sqrt{x}$ the x axis, and the line $x = 4$ using

1. the shell method, and
2. the disk method.

Sometimes the best way to get a deep understanding of a concept is to take a simple problem and work it in every possible way. The next problem is just that sort of problem.

Problem 210. Consider the region in the first quadrant bounded by $y = x^2$, the y -axis, and $y = 4$. Find the volume of the solid obtained by rotating this region about:

1. the y -axis using the disk method,
2. the y -axis using the shell method,
3. the x -axis using the disk method, and
4. the x -axis using the shell method.

Problem 211. Use any method to find the volume of the solid generated by revolving about the y -axis the region bounded by the curves $y = 8 - x^2$ and the curve $y = x^2$.

Arc Length

Every one who roller blades or rides a skateboard knows that you can scrub speed off by slalom-ing back and forth as you proceed down a hill. In doing so, you travel a greater distance than the speed freak who takes a straight line. If we have a road of length 2π and one person takes the straight line path while a second follows the sine curve, how much farther does the second person travel?

Problem 212. Let $f(x) = \sin(x)$ from $x = 0$ to $x = 2\pi$.

1. Consider the partition of $[0, 2\pi]$, defined by $x_0 = 0$, $x_1 = \pi/2$, $x_2 = \pi$, $x_3 = 3\pi/2$, $x_4 = 2\pi$. Approximate the length of the curve by summing the lengths of the straight line segments from $(x_{i-1}, f(x_{i-1}))$ to $(x_i, f(x_i))$ for $i = 1, 2, 3, 4$.
2. Generalize this so that the partition has N divisions and take the limit as $N \rightarrow \infty$.

Definition 38. A function f is said to be **continuously differentiable** on $[a, b]$ if its derivative f' is continuous on $[a, b]$. If f is continuously differentiable on $[a, b]$, then its graph on the interval $[a, b]$ is called a **smooth curve**.

Problem 213. Suppose the function f and its derivative f' are continuous on $[a, b]$. Let s be the arc length of the curve f from the point $(a, f(a))$ to $(b, f(b))$.

1. Let $a = x_0 < x_1 < x_2 < \cdots < x_n = b$ be a partition of $[a, b]$.

$$\text{Show that } s \approx \sum_{i=1}^n (x_i - x_{i-1}) \sqrt{1 + \left(\frac{f(x_i) - f(x_{i-1})}{x_i - x_{i-1}} \right)^2}.$$

2. Show that $s = \int_a^b \sqrt{1 + [f'(x)]^2} dx$ by using the Mean Value Theorem for differentiation.

The result you just proved can be summarized in the following theorem.

Theorem 20. Arc Length Theorem. If s is the arc length of the smooth curve f on the interval $[a, b]$, then $s = \int_a^b \sqrt{1 + [f'(x)]^2} dx$.

Problem 214. For each of the following curves, sketch its graph and find the length of the arc with the indicated endpoints.

1. $y = x^{\frac{3}{2}}$ from the point $(0, 0)$ to the point $(4, 8)$.
2. $y = \frac{1}{4}x^3 + \frac{1}{3}x^{-1} + 2$ from the point where $x = 2$ to the point $x = 5$.
3. $x^2 = (y + 1)^3$ from the point where $x = 0$ to the point where $x = 3\sqrt{3}$. Hint: You may integrate with respect to x or y . Which is easier?

Center of Mass

Definition 39. One-dimensional Moments (Discrete Case) Place n weights with masses $m_1, m_2, \dots, m_n$ at the points $(x_1, 0), (x_2, 0), \dots, (x_n, 0)$ on the x -axis.

The **mass** is

$$M = \sum_{i=1}^n m_i ,$$

the **moment of mass** is

$$M_0 = \sum_{i=1}^n x_i m_i ,$$

and the **center of mass** is

$$\bar{x} = \frac{\sum_{i=1}^n x_i m_i}{\sum_{i=1}^n m_i} ,$$

which is the point on the x -axis where everything would balance.

Definition 40. One-dimensional Moments (Continuous Case). Let L be the length of a rod whose left hand endpoint is at the origin of the x axis and let ρ be a continuous function on $[0, L]$ such that $\rho(x)$ is the density at $x \in [0, L]$.

The **mass** is

$$M = \int_0^L \rho(x) \, dx ,$$

the **moment of mass** is

$$M_0 = \int_0^L x\rho(x) \, dx ,$$

and the **center of mass** is

$$\bar{x} = \frac{\int_0^L x\rho(x) \, dx}{\int_0^L \rho(x) \, dx} .$$

Problem 215. Find the moment of mass and the center of mass of each of the following systems.

1. $m_1 = 4$ at $x_1 = -3$; $m_2 = 1$ at $x_2 = -1$; $m_3 = 2$ at $x_3 = 3$; $m_4 = 1$ at $x_4 = -5$; $m_5 = 5$ at $x_5 = -4$; $m_6 = 6$ at $x_6 = 1$
2. A rod whose length is 5 meters with linear density $3x + 2$ kg/m where $x = 0$ is the left-hand endpoint of the rod.

Definition 41. Two-dimensional Moments (Discrete Case). Assume that the thickness and the weight of a sheet of material are negligible and there are n particles with mass $m_1, m_2, \dots, m_n$ located at $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ on the xy plane. Then the **total mass** of the system (M), the **total moment with respect to the x axis** (M_x), the **total moment with respect to the y axis** (M_y), and the **center of mass of the system** $((\bar{x}, \bar{y}))$, are defined by

$$M = \sum_{i=1}^n m_i \quad ,$$

$$(M_x, M_y) = \left(\sum_{i=1}^n y_i m_i, \sum_{i=1}^n x_i m_i \right) \quad , \text{ and}$$

$$(\bar{x}, \bar{y}) = \left(\frac{M_y}{M}, \frac{M_x}{M} \right) \quad .$$

Definition 42. Two-dimensional Moments (Continuous Case). A **lamina** is a two dimensional material of continuously distributed mass. A **homogeneous lamina** is a lamina with constant area density. Assume that a lamina L is bounded by the curve $y = f(x)$, the x axis, and the lines $x = a$ and $x = b$ such that f is continuous on $[a, b]$, $f(x) \geq 0$ for all x in $[a, b]$, and the **area density** function ρ is also continuous on $[a, b]$. Then the **total mass** of the system (M), the **total moment with respect to the x axis** (M_x), the **total moment with respect to the y axis** (M_y), and the **center of mass of the system** $((\bar{x}, \bar{y}))$, are defined by

$$M = \int_a^b \rho(x) f(x) dx \quad ,$$

$$(M_x, M_y) = \left(\frac{1}{2} \int_a^b \rho(x) [f(x)]^2 dx, \int_a^b x \rho(x) f(x) dx \right) \quad , \text{ and}$$

$$(\bar{x}, \bar{y}) = \left(\frac{M_y}{M}, \frac{M_x}{M} \right) \quad .$$

Problem 216. Find the total mass and center of mass of the collection of objects: $m_1 = 4$ at $(-1, 3)$; $m_2 = 1$ at $(0, 0)$; $m_3 = 2$ at $(2, -4)$; $m_4 = 1$ at $(4, -3)$.

Problem 217. Find the total mass and center of mass for a lamina bounded by the curve $y = x^2 - 4$ and the $y = 3x + 6$ with the area density function $\rho(x) = x + 1$.

Chapter 5

Techniques of Integration

“That student is taught the best who is told the least.” - R. L. Moore

Change of variable (substitution), partial fractions, integration by parts, and trigonometric substitutions are four techniques of integration that can be applied to evaluate indefinite integrals. More than one technique will often work on the same problem.

5.1 Change of Variable Method

The integrand can often be rewritten so that it is easier to see how one of our derivative theorems can be applied to find the antiderivative of the integrand. This method is called either a *change of variable* or *substitution* because you replace some part of the integrand with a new function.

Example 22. Consider $\int \frac{x}{\sqrt{x+5}} dx$. If we let $u(x) = \sqrt{x+5}$, then $u'(x) = \frac{1}{2\sqrt{x+5}}$ so

$$\int \frac{x}{\sqrt{x+5}} dx = 2 \int (u^2(x) - 5)u'(x) dx = 2 \int u^2(x)u'(x) - 5u'(x) dx = \frac{1}{3}u^3(x) - 5u(x) = \dots$$

Problem 218. Evaluate the indefinite integral, $\int x^2(2x^3 + 10)^{15} dx$, using the following steps.

1. Let $u(x) = 2x^3 + 10$ and compute u' .
2. Show that $\int x^2(2x^3 + 10)^{15} dx = \frac{1}{6} \int (u(x))^{15} u'(x) dx$.
3. Evaluate this indefinite integral and then replace $u(x)$ with $2x^3 + 10$ in your answer to eliminate the function u from your answer.
4. Verify your answer by taking the derivative.

Problem 219. Evaluate each of the following integrals.

1. $\int (2x + 7) \sqrt[3]{x^2 + 7x + 3} dx$

2. Three parts to this one!

(a) $\int \sqrt{2-x} dx$

$$(b) \int x\sqrt{2-x} \, dx$$

$$(c) \int x^2\sqrt{2-x} \, dx$$

$$3. \int \frac{(\ln x)^4}{x} \, dx$$

$$4. \int 5x^2 4^{x^3} \, dx$$

$$5. \int_0^{\frac{\pi}{8}} \sin^5(2x) \cos(2x) \, dx$$

6. *Three parts to this one!*

$$(a) \int \frac{x}{(x^2+2)^2} \, dx$$

$$(b) \int \frac{x}{\sqrt[4]{x^2+2}} \, dx$$

$$(c) \int \frac{x}{x^2+2} \, dx$$

5.2 Partial Fractions

This section focuses on integrating rational functions. The method of *partial fractions* may be used on rational functions to break one rational function into a sum of two or more rational functions, each of which has a lower degree denominator and is easier to anti-differentiate. When a rational function has a numerator with degree greater than the degree of the denominator, one can long divide to rewrite the integrand as the sum of a polynomial and another rational function with numerator of lower degree than the degree of the denominator. Sometimes the rational function displays features of our derivative theorems and thus is already in a form that is integrable.

Problem 220. *You will integrate this rational function by breaking it into two rational functions and integrating each of those functions separately.*

$$1. \text{ Find numbers } A \text{ and } B \text{ so that } \frac{1}{x^2-1} = \frac{A}{x-1} + \frac{B}{x+1}.$$

$$2. \text{ Evaluate } \int \frac{1}{x^2-1} \, dx = \int \frac{A}{x-1} \, dx + \int \frac{B}{x+1} \, dx = \dots$$

Problem 221. *Evaluate $\int \frac{4x}{x^2-2x-3} \, dx$ via partial fractions.*

Problem 222. *Evaluate $\int \frac{3x^2-2x+1}{x^2-2x-3} \, dx$. This is called an improper fraction because the degree of the numerator is greater than or equal to the degree of the denominator. In such cases, long divide first.*

Problem 223. *Evaluate $\int \frac{x^2+x-2}{x(x+1)^2} \, dx$ via partial fractions by first finding constants, A, B , and C so that $\frac{x^2+x-2}{x(x+1)^2} = \frac{A}{x} + \frac{B}{x+1} + \frac{C}{(x+1)^2}$.*

Problem 224. Evaluate $\int \frac{5x^2 + 3}{x^3 + x} dx$ via partial fractions by first finding constants A , B , and C so that $\frac{5x^2 + 3}{x^3 + x} = \frac{A}{x} + \frac{Bx + C}{x^2 + 1}$.

Problem 225. Evaluate these three indefinite integrals that look similar, but are not!

$$1. \int \frac{5}{x^2 - 1} dx$$

$$2. \int \frac{5x}{x^2 - 1} dx$$

$$3. \int \frac{5}{x^2 + 1} dx$$

5.3 Integration by Parts

Integration by parts is an application of the product rule – essentially the product rule in reverse! It is typically used when the integrand is the product of two functions, one which you can easily anti-differentiate and one which becomes simpler (or at least no more complicated) when you differentiate.

Problem 226. Find two functions f and g so that $\int f(x) \cdot g(x) dx \neq \int f(x) dx \cdot \int g(x) dx$.

Example 23. Evaluate the integral $\int x \ln(x) dx$.

We begin by guessing a function which has $x \ln(x)$ as a part of its derivative when we use the product rule. From the product rule for derivatives we have,

$$(x^2 \ln x)' = 2x \ln x + x^2 \left(\frac{1}{x}\right).$$

Integrating both sides yields,

$$\int (x^2 \ln x)' dx = \int 2x \ln x dx + \int x dx.$$

Now we have,

$$x^2 \ln x + k_1 = 2 \int x \ln x dx + \frac{1}{2}x^2 + k_2$$

Where k_1 and k_2 are constants. Solving for the integral we wanted yields,

$$\int x \ln x dx = \frac{1}{2}x^2 \ln x - \frac{1}{4}x^2 + \frac{1}{2}(k_1 - k_2).$$

Problem 227. Use the idea from the example to compute the following anti-derivatives.

$$1. \int x^{45} \ln x dx$$

$$2. \int x e^x dx$$

$$3. \int x \cos(x) dx$$

Problem 228. Let f and g be differentiable functions. Starting with the product rule for derivatives, show that

$$\int f(x)g'(x) dx = f(x) \cdot g(x) - \int g(x)f'(x) dx.$$

Problem 229. Evaluate each indefinite integral.

1. $\int x^2 e^x dx$

2. $\int x^2 \sin(x) dx$

3. $\int e^x \cos x dx$

4. $\int (\ln x)^2 dx$

5. $\int \sec^3 x dx$

Problem 230. Compute the derivatives of each.

1. $s(x) = \sin^{-1}(x)$ by differentiating the equation $\sin(s(x)) = x$ and solving for $s'(x)$

2. $c(x) = \sec^{-1}(x)$

3. $t(x) = \tan^{-1}(x)$

Now we address the integrals of the inverse trigonometric functions. Use integration by parts since you know the derivatives of each function.

Problem 231. Compute the following indefinite integrals via integration by parts.

1. $\int \sin^{-1}(x) dx$ using $\sin^{-1}(x) = \sin^{-1}(x) \cdot 1$.

2. $\int \tan^{-1}(x) dx$

3. $\int \sec^{-1}(x) dx$

Theorem 21. Integration by Parts. If each of f and g are functions that are differentiable on the interval $[a, b]$, then

$$\int f(x)g'(x) dx = f(x) \cdot g(x) - \int g(x)f'(x) dx$$

and

$$\int_a^b f(x)g'(x) dx = f(x) \cdot g(x)|_a^b - \int_a^b g(x)f'(x) dx$$

Writing $f(x)$ as u and $g(x)$ as v and using Leibnitz notation, this formula is often written as:

$$\int u dv = uv - \int v du.$$

5.4 Integrals of Trigonometric Functions

When integrating trigonometric functions you will often use a combination of substitution, integration by parts, and trigonometric identities. Appendix 15.4 has the trigonometric identities that will be necessary for this section: the Pythagorean, double angle, half angle, and product identities.

Problem 232. Evaluate $\int \cos^3(x) dx$ by using the fact that $\cos^3 = \cos^2 \cdot \cos$ and the Pythagorean identity for $\cos^2$.

Problem 233. Evaluate $\int \cos^5(x) dx$ by using the fact that $\cos^5 = \cos^4 \cdot \cos$

Problem 234. Evaluate $\int \cos^{2n+1}(x) dx$ and $\int \sin^{2n+1}(x) dx$ where n is a positive integer.

Problem 235. Evaluate $\int \sin^4(x) dx$ using the half-angle identity for $\sin$.

Problem 236. Evaluate each of the following integrals.

1. $\int \sin^3(x) \cos^2(x) dx$

2. $\int \sin^3(x) \cos^5(x) dx$

3. $\int \sin^4(x) \cos^2(x) dx$

4. $\int \sin(3x) \cos(6x) dx$

Problem 237. Evaluate $\int \sec(x) dx$ by first multiplying by $\frac{\sec(x) + \tan(x)}{\sec(x) + \tan(x)}$.

Problem 238. Evaluate $\int \tan^3(x) \sec^4(x) dx$ in the two different ways indicated.

1. Use $\tan^3 \cdot \sec^4 = \tan^2 \cdot \sec^3 \cdot \tan \cdot \sec$ and the Pythagorean identity for $\tan^2$.

2. Use $\tan^3 \cdot \sec^4 = \tan^3 \cdot \sec^2 \cdot \sec^2$ and the Pythagorean identity for $\sec^2$.

Problem 239. Evaluate each of the following integrals.

1. $\int \cot^5(x) dx$

2. $\int_{\frac{3\pi}{4}}^{\frac{5\pi}{4}} \sec^4(x) dx$

3. $\int \cot^3(x) \csc^5(x) dx$

4. $\int \tan^2(x) \sec^4(x) dx$

5. $\int \tan^4(x) \sec^3(x) dx$

Problem 240. Not for presentation. Integrate the first, second, third, and fourth power of each of the six trigonometric functions. It is a well-documented fact that one of these (24) problems has appeared on at least one test in every Calculus II course since the days of Newton and Leibnitz.

5.5 Integration by Trigonometric Substitution

We have already used substitution (change of variables) to transform difficult integration problems into easier ones. Trigonometric substitutions are an extension of this same theme.

Problem 241. Derive the formula ($A = \pi r^2$) for the area of a circle of radius r using integration as follows.

1. Compute the definite integral $\int_{-3}^3 \sqrt{9-x^2} dx$ by making the substitution, $x(t) = 3 \sin(t)$.
2. Show that the area inside of a circle $x^2 + y^2 = r^2$ is πr^2 by setting up and evaluating an appropriate definite integral using the substitution $x(t) = r \sin(t)$.

Problem 242. Evaluate each of the following integrals.

1. $\int \frac{x^2}{\sqrt{x^2+4}} dx$ by using the substitution $x = 2 \tan(\theta)$
2. $\int_0^4 5x \sqrt{16-x^2} dx$
3. $\int_2^3 \frac{1}{4x^3 \sqrt{x^2-1}} dx$ by using the substitution $x = \sec(\theta)$
4. $\int \frac{1}{\sqrt{x^2-2x+5}} dx$ by completing square and letting $u = x-1$
5. $\int \frac{2x+5}{x^2+4x+1} dx$

Definition 43. If a and b are non-zero numbers, then the set of points in the plane satisfying

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

is an ellipse centered at the origin.

Problem 243. Consider the ellipse, $\frac{x^2}{25} + \frac{y^2}{16} = 1$. Set up and evaluate a definite integral that represents the area (or half the area) inside this ellipse.

Problem 244. Solve each of the following problems. Assume that a is a real number.

1. Evaluate $\int \frac{1}{\sqrt{a^2-x^2}} dx$
2. Evaluate $\int \frac{1}{x^2+a^2} dx$
3. Evaluate $\int \frac{1}{\sqrt{x^2-a^2}} dx$

Problem 245. Verify the formula $A = \pi ab$ for the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ where a and b are positive numbers.

5.6 The Hyperbolic Functions

Similar to the trigonometric functions in Euclidean trigonometry, the following functions are important in **hyperbolic trigonometry**. These occur in engineering problems because they can be used to represent the curve that results when hanging a rope between two poles (think telephone lines).

Definition 44. The **hyperbolic sine and hyperbolic cosine** functions are denoted by ***sinh*** and ***cosh*** and defined by:

$$\sinh(x) = \frac{e^x - e^{-x}}{2} \text{ and } \cosh(x) = \frac{e^x + e^{-x}}{2}.$$

The **hyperbolic tangent, hyperbolic cotangent, hyperbolic secant, and hyperbolic cosecant** functions are denoted by ***tanh***, ***coth***, ***sech***, ***csch*** and defined by:

$$\tanh(x) = \frac{\sinh(x)}{\cosh(x)}, \quad \coth(x) = \frac{\cosh(x)}{\sinh(x)}, \quad \operatorname{sech}(x) = \frac{1}{\cosh(x)}, \quad \text{and} \quad \operatorname{csch}(x) = \frac{1}{\sinh(x)}.$$

Problem 246. Assume that x is a number and show that each of the following is true.

1. $\cosh^2(x) - \sinh^2(x) = 1$
2. $1 - \tanh^2(x) = \operatorname{sech}^2(x)$
3. $1 - \coth^2(x) = -\operatorname{csch}^2(x)$

Problem 247. Assume that x is a number and show that each of the following is true.

1. $\frac{d}{dx} \sinh(x) = \cosh(x)$
2. $\frac{d}{dx} \cosh(x) = \sinh(x)$
3. $\frac{d}{dx} \tanh(x) = \operatorname{sech}^2(x)$
4. $\frac{d}{dx} \coth(x) = -\operatorname{csch}^2(x)$
5. $\frac{d}{dx} \operatorname{sech}(x) = -\operatorname{sech}(x) \cdot \tanh(x)$
6. $\frac{d}{dx} \operatorname{csch}(x) = -\operatorname{csch}(x) \cdot \coth(x)$

Problem 248. Based on parts 1 through 5 of the last problem, write the corresponding antiderivative formulas. For example, 1. $\int \cosh(x) \, dx = \sinh(x) + C$ is the integral formula corresponding to part 1 in the last problem.

While many inverse functions (such as $\tan^{-1}$) cannot be written out explicitly as a function of x only, the inverse hyperbolic trigonometric functions can be.

Problem 249. Rewrite $\cosh^{-1}$ as a function of x and without using any hyperbolic functions by solving $y = \frac{e^x + e^{-x}}{2}$ for x . (This can be done for all six of the hyperbolic trig functions.)

Problem 250. Show that $\frac{d}{dx} \sinh^{-1}(x) = \frac{1}{\sqrt{x^2+1}}$ by letting $y = \sinh^{-1}(x)$, applying implicit differentiation to $\sinh(y) = x$, and using the identity $\cosh^2(x) - \sinh^2(x) = 1$.

Problem 251. Verify each of the following. Why are there restrictions on the domains?

1. $\frac{d}{dx} \cosh^{-1}(x) = \frac{1}{\sqrt{x^2-1}}$ for $x > 1$
2. $\frac{d}{dx} \tanh^{-1}(x) = \frac{1}{1-x^2}$ for $|x| < 1$
3. $\frac{d}{dx} \operatorname{cotanh}^{-1}(x) = \frac{1}{1-x^2}$ for $|x| > 1$
4. $\frac{d}{dx} \operatorname{sech}^{-1}(x) = -\frac{1}{x\sqrt{1-x^2}}$ for $0 < x \leq 1$
5. $\frac{d}{dx} \operatorname{csch}^{-1}(x) = -\frac{1}{|x|\sqrt{1+x^2}}$ for $x \neq 0$

5.7 Improper Integrals

In our definition of the definite integral $\int_a^b f(x) dx$, it was assumed that f was defined on all of $[a, b]$ where a and b were numbers. We wish to extend our definition to consider the possibilities where a and b are $\pm\infty$ and where f might not be defined at every point in $[a, b]$. When any of these possibilities occur, the integral is called an **improper integral**.

Problem 252. Let $f(x) = \frac{1}{x^2}$.

1. Sketch a graph of f .
2. Compute $\int_1^{10} f(x) dx$, $\int_1^{50} f(x) dx$, and $\int_1^{100} f(x) dx$.
3. Compute $\lim_{N \rightarrow \infty} \int_1^N f(x) dx$.

Definition 45. If a is a number and f is continuous for all $x \geq a$, then we define the **improper integral** $\int_a^\infty f(x) dx$ to be $\lim_{N \rightarrow \infty} \int_a^N f(x) dx$. If this limit exists, we say the integral **converges** and if it does not exist, we say the integral **diverges**.

Problem 253. Write down definitions for $\int_{-\infty}^a f(x) dx$ and $\int_{-\infty}^\infty f(x) dx$.

Problem 254. Let $f(x) = \frac{4}{(x-1)^2}$.

1. Sketch the graph of f .
2. Evaluate $\int_2^{100} f(x) dx$

3. Evaluate $\int_2^{100,000} f(x) dx$

4. Evaluate $\int_2^{\infty} f(x) dx$

5. Evaluate $\int_a^{\infty} f(x) dx$ where $a > 1$.

Problem 255. Sketch $f(x) = \frac{1}{x-1}$ and evaluate $\int_2^{\infty} f(x) dx$.

Problem 256. Sketch $f(x) = \frac{1}{4+x^2}$ and evaluate $\int_{-\infty}^{\infty} f(x) dx$.

Problem 257. Evaluate $\int_{-\infty}^{-2} \frac{5}{x\sqrt{x^2-1}} dx$.

Definition 46. If f is continuous on the interval $[a, b)$, $\lim_{x \rightarrow b^-} |f(x)| = \infty$, then we define $\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx$.

Problem 258. Definitions.

1. Suppose f is continuous on the interval $(a, b]$ and $\lim_{x \rightarrow a^+} |f(x)| = \infty$. Write a definition for

$$\int_a^b f(x) dx.$$

2. Suppose f is continuous on the interval $[a, b]$ except at c where $a < c < b$. Suppose $\lim_{x \rightarrow c^-} =$

$$\lim_{x \rightarrow c^+} f(x) = \pm\infty. \text{ Write a definition for } \int_a^b f(x) dx.$$

Problem 259. Evaluate each of the following improper integrals.

1. $\int_0^2 \frac{1}{(x-2)^2} dx$

2. $\int_3^5 \frac{x}{4-x} dx$

3. $\int_0^{\frac{\pi}{2}} \cot x dx$

4. $\int_0^e \frac{\ln x}{x} dx$

5. $\int_1^{\infty} \frac{dx}{-2x\sqrt{x^2-1}}$

Chapter 6

Sequences & Series

“All the instruments have been tried save one, the only one precisely that can succeed: well-regulated freedom.” - Jean-Jacques Rousseau

Intuitively speaking, a *sequence* may be thought of as a list of numbers and a *series* may be thought of as the sum of a sequence of numbers. The sequence below is approaching 0. Does the series below add up to a number or as we keep adding terms do the sums tend to infinity?

Sequence: $1, 1/2, 1/3, 1/4, \dots$

Series: $1 + 1/2 + 1/4 + 1/8 + \dots$

6.1 Sequences

Definition 47. A **sequence** is a function with domain a subset of the natural numbers and range a subset of the real numbers.

Example 24. We will denote our sequences by listing elements of the range of the sequence.

- $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$ denotes the range of the sequence $f(n) = \frac{1}{2^n}$, $n = 1, 2, 3, \dots$
- $a_1, a_2, a_3, \dots$ denotes the range of the sequence $f(n) = a_n$, $n = 1, 2, 3, \dots$
- $\{x_n\}_{n=1}^{\infty}$ denotes the range of the sequence $f(n) = x_n$, $n = 1, 2, 3, \dots$
- $\{x_n = \frac{n}{n+1}\}_{n=1}^{\infty}$ denotes the range of the sequence $f(n) = \frac{n}{n+1}$, $n = 1, 2, 3, \dots$

We mathematicians abuse the definition of function by referring to $y = x^3$ as a function, when in fact the function is the collection of ordered pairs generated by this equation. In a similar abuse, while $\{x_n\}_{n=1}^{\infty}$ or $x_1, x_2, x_3, \dots$ represents the elements of the range of the sequence, we refer to it as though it were the sequence itself.

Definition 48. The sequence $\{x_n\}_{n=1}^{\infty}$ is said to be **bounded above** if there is number M such that $x_n \leq M$ for all $n \geq 1$. If such a number exists, then M is called an **upper bound** for the sequence. **Bounded below** and **lower bound** are defined in the analogous way. The sequence is **bounded** if it is bounded above and bounded below.

Problem 260. Show that the sequence, $\{x_n = \frac{1}{n}\}_{n=1}^{\infty}$ is bounded above and below.

Definition 49. The sequence $\{x_n\}_{n=1}^{\infty}$ is **increasing** (often called strictly increasing) if $x_n < x_{n+1}$ for all natural numbers n . The sequence $\{x_n\}_{n=1}^{\infty}$ is **non-decreasing** if $x_n \leq x_{n+1}$ for all natural numbers n . **Decreasing** and **non-increasing** are defined similarly. A sequence is said to be **monotonic** if it is either increasing, decreasing, non-increasing, or non-decreasing.

Definition 50. The **factorial** function is defined for all non-negative integers as follows:

1. $0! = 1$
2. $n! = n \cdot (n-1)! \quad \text{for } n = 1, 2, 3, \dots$

The first few factorials are:

$$0! = 1, \quad 1! = 1, \quad 2! = 2 \cdot 1 = 2, \quad 3! = 3 \cdot 2 \cdot 1 = 6, \quad 4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24, \dots$$

Problem 261. For each of the following, prove that it is monotonic or show why it is not monotonic.

1. $\{\frac{n+3}{4n+2}\}_{n=1}^{\infty}$
2. $\{\sin(\frac{n\pi}{3})\}_{n=1}^{\infty}$
3. $\{\frac{n!}{2^n}\}_{n=0}^{\infty}$

Definition 51. Given the sequence, $\{x_n\}_{n=1}^{\infty}$, the number G is called the **greatest lower bound** of the sequence if G is a lower bound for the sequence and no other lower bound is larger than G . The number L is called the **least upper bound** if L is an upper bound and no other upper bound of the sequence is less than L .

Problem 262. Find the least upper bound and the greatest lower bound for these sequences.

1. $\{\frac{n}{n+1}\}_{n=1}^{\infty}$
2. $\{\frac{2^n}{n!}\}_{n=2}^{\infty}$
3. $\{\frac{2n+1}{n}\}_{n=2}^{\infty}$

Problem 263. For each sequence, write out the first five terms. Is there a number that the values of x_n approach as $n \rightarrow \infty$? If so, what is it. If not, why not?

1. $\{x_n = \frac{1}{n} - \frac{1}{n-1}\} \quad \text{for } n = 2, 3, 4, \dots$
2. $\{x_n = (-1)^n(2 - \frac{1}{n})\} \quad \text{for } n = 1, 2, 3, \dots$
3. $\{\frac{n^2}{n!}\}_{n=5}^{\infty}$

Definition 52. Given a sequence, $\{x_n\}_{n=k}^{\infty}$, we say that the sequence **converges** to the number L if for every positive number ϵ , there is a natural number N so that for all $n \geq N$, we have $|x_n - L| < \epsilon$.

If $\{x_n\}_{n=k}^{\infty}$ converges to L , we may write “ $\lim_{n \rightarrow \infty} x_n = L$ ” or “ $x_n \rightarrow L$ as $n \rightarrow \infty$.” We call a sequence **convergent** if there is a number to which it converges, and **divergent** otherwise. A divergent sequence might tend to $\pm\infty$ or might “oscillate.” In either case, there is no single number that the range of the sequence approaches.

Problem 264. Consider the sequence, $\frac{5n+4}{3n+1}$. Let $\varepsilon = 0.00002$ and find the smallest natural number N so that $|\frac{5n+4}{3n+1} - \frac{5}{3}| < \varepsilon$ for every $n \geq N$. Repeat for $\varepsilon = 0.00001$.

Problem 265. Consider the sequence, $\frac{n^2}{n^2+1}$. Let $\varepsilon = .0004$ and find the smallest natural number N such that $|\frac{n^2}{n^2+1} - 1| < \varepsilon$ for all $n \geq N$. Repeat for $\varepsilon = .0002$.

Problem 266. Let ε represent a positive number less than one. Find the smallest natural number N such that $|\frac{5n+4}{3n+1} - \frac{5}{3}| < \varepsilon$ for every $n \geq N$. The number N will depend on ε since a smaller value for ε may result in a larger value for N .

Problem 267. Let $\{x_n\}_{n=1}^{\infty}$ be the sequence so that $x_n = 0$ if n is odd and $x_n = 1$ if n is even.

1. Show that the sequence is bounded.
2. Show that the sequence does not converge to 0.
3. What would we need to prove in order to show that the sequence is divergent?

Problem 268. Let ε represent a positive number less than one. Find the smallest positive integer N such that $|\frac{n^2}{n^2+1} - 1| < \varepsilon$ for all $n \geq N$.

Problem 269. Show that if $\{a_n\}_{n=1}^{\infty}$ is a convergent sequence then it is a bounded sequence.

Definition 53. The sequence $\{x_n\}_{n=k}^{\infty}$ is called a **Cauchy sequence** if for every $\varepsilon > 0$, there is a natural number N so that $|x_n - x_m| < \varepsilon$ for all $m, n \geq N$.

Problem 270. Show that if a and b are numbers then $|a + b| \leq |a| + |b|$.

Problem 271. Show that if $\{a_n\}_{n=1}^{\infty}$ is a convergent sequence then it is a Cauchy sequence.

It is also true that if $\{a_n\}_{n=1}^{\infty}$ is a Cauchy sequence then it is a convergent sequence. That's a bit more interesting problem than we have time to explore – take *Analysis*.

Definition 54. The sequence $\{x_n\}_{n=k}^{\infty}$ is called a **C sequence** if for every $\varepsilon > 0$, there is natural number N such that $|x_N - x_n| < \varepsilon$ for all $n \geq N$.

Problem 272. Two parts.

1. Suppose that $\{a_n\}_{n=1}^{\infty}$ is a C sequence. Is it a Cauchy sequence?
2. Suppose that $\{a_n\}_{n=1}^{\infty}$ is a Cauchy sequence. Is it a C sequence?

An **axiom** is a statement that we use without proof. All of mathematics is built on axioms since one must make certain assumptions just to get started. There are entire branches of mathematics devoted to determining what of the mathematics that we use now would still be true if we changed the underlying axiom system that mathematicians (more or less) universally agree on.

The **Completeness Axiom** says that if we have a set of numbers and there is some number that is greater than every number in our set, then there must be a *smallest* number that is greater than or equal to every number in the set.

Axiom 1. Completeness Axiom *If a non-empty set S of real numbers has an upper bound, then S has a least upper bound.*

Theorem 22. Monotonic Convergence Theorem. *Every bounded monotonic sequence is convergent.*

Problem 273. *Show that $\{\frac{2^n}{n!}\}_{n=1}^{\infty}$ is convergent.*

The next theorem formalizes the relationship between sequences and limits of functions.

Theorem 23. *Let $\{x_n\}_{n=1}^{\infty}$ be a sequence and f be a function defined for all $x \geq 1$ such that $f(n) = x_n$ for all $n = 1, 2, 3, \dots$. The only difference between the function f and the function x (remember, a sequence is a function) is that the domain of f includes real numbers and the domain of x is only natural numbers.*

1. *If $\lim_{x \rightarrow \infty} f(x) = L$, then $\lim_{n \rightarrow \infty} x_n = L$.*
2. *If $\lim_{x \rightarrow \infty} f(x) = \pm\infty$, then $\lim_{n \rightarrow \infty} x_n = \pm\infty$.*
3. *If $\lim_{n \rightarrow \infty} x_n = L$ and $\lim_{n \rightarrow \infty} y_n = M$, then*
 - (a) $\lim_{n \rightarrow \infty} (x_n \pm y_n) = L \pm M$;
 - (b) $\lim_{n \rightarrow \infty} (x_n \cdot y_n) = LM$;
 - (c) $\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{L}{M}$ for $M \neq 0$.

Problem 274. *Consider the sequence $x_n = \frac{1}{n}$ for all $n = 1, 2, 3, \dots$. Sketch and define a function, f , so that f agrees with x_n at all the natural numbers, but $\lim_{x \rightarrow \infty} f(x) \neq 0$.*

Problem 275. *Evaluate each of the following limits, whenever it exists.*

1. $\lim_{n \rightarrow \infty} \frac{n^2 + 2n + 3}{3n^2 + 4n - 5}$
2. $\lim_{n \rightarrow \infty} \frac{6n + 100}{n^3 + 4}$
3. $\lim_{n \rightarrow \infty} \frac{\log_3(n)}{n}$
4. $\lim_{n \rightarrow \infty} (1 + \frac{3}{n})^n$

6.2 Series with Positive Constant Terms

Example 25. Bouncing balls.

Suppose we drop two balls at the same time from a height of one foot and the first one bounces to the height of one foot with each bounce. (Question for my Mechanical Engineers. Why is it that if you invent this ball, you can retire?) The second ball bounces to half the height that it fell from each time. What distance does each ball travel? Answer:

Ball 1: $1 + 1 + 1 + 1 + 1 + \dots$

and

$$\text{Ball 2: } 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{4} + \frac{1}{4} + \dots = 1 + 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots$$

Such sums are called *series* or *infinite series* and the individual numbers that are added are called the *terms* of the series. The first sum does not “add up,” so we’ll say it *diverges*. The second sum does “add up,” so we’ll say it *converges*. To see that the second sum converges, let

$$S_1 = 1, S_2 = 1 + 1, S_3 = 1 + 1 + \frac{1}{2}, S_4 = 1 + 1 + \frac{1}{2} + \frac{1}{4}, \dots$$

and consider the limit of this new sequence that we have created. The new sequence we have generated, $S_1, S_2, S_3, \dots$ is called the *sequence of partial sums* of the series and looks like this

$$S_1 = 1, S_2 = 2, S_3 = 2\frac{1}{2}, S_4 = 2\frac{3}{4}, \dots$$

Problem 276. Write down the simplest expression you can find for the N^{th} partial sum, S_N , for the series $1 + 1 + 1 + \dots$ and compute $\lim_{N \rightarrow \infty} S_N$.

Problem 277. Write down the simplest expression you can find for the N^{th} partial sum, S_N , for the series $1 + 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$ and compute $\lim_{N \rightarrow \infty} S_N$.

This notion of adding up partial sums of the series is the motivation for the definition of a series. We say that an infinite series *converges* if the limit of the partial sums exists.

Definition 55. Given a sequence $\{a_n\}_{n=k}^{\infty}$, where k is a positive integer or zero, we define the N^{th} **partial sum** of this sequence to be the number $S_N = \sum_{n=k}^N a_n$. We denote the limit of the sequence of partial sums by the notation, $\sum_{n=k}^{\infty} a_n = \lim_{N \rightarrow \infty} S_N$, and call this the *infinite series* associated with the sequence.

Definition 56. Given a sequence, $\{a_n\}_{n=k}^{\infty}$, the series $\sum_{n=k}^{\infty} a_n$ is said to **converge** if $\lim_{N \rightarrow \infty} S_N$ exists and is said to **diverge** if $\lim_{N \rightarrow \infty} S_N$ does not exist.

Problem 278. Let $a_0 = 1 - \frac{1}{2}$, $a_1 = \frac{1}{2} - \frac{1}{3}$, $a_2 = \frac{1}{3} - \frac{1}{4}$, $\dots$.

1. Find a formula for the n^{th} term of the sequence, a_n .
2. Compute the limit of the terms of the sequence, a_n .
3. Find a formula for the N^{th} partial sum of the sequence, S_N .
4. Compute the limit of the partial sums, $\lim_{N \rightarrow \infty} S_N = \sum_{n=1}^{\infty} a_n$.

We are about to develop many theorems (called “tests”) to tell us if a series converges or diverges. Like the various techniques of integration, each test is easy to apply, but figuring out which test to use on a particular series may be challenging.

In our bouncing ball example, you found a formula for the n^{th} partial sum, S_N and showed that $\lim_{N \rightarrow \infty} S_N$ was a real number. Since the sequence of partial sums converged, we say that our series converges. For this example, we were able to write down an exact expression for the N^{th} partial sum and we were able to compute the limit. In many cases, we will be able to determine whether a particular series converges, but not know from the test what it converges to. Knowing that the series converges is valuable, because as long as we *know* that it converges then we can use our calculators or computer algebra systems to add enough terms to get as close to the answer as we want.

Problem 279. Let $a_0 = 3$, $a_1 = 3(\frac{1}{2})$, $a_2 = 3(\frac{1}{2})^2$, $\dots$.

1. Find a formula for the n^{th} term, a_n .
2. Compute the limit of the terms, $\lim_{n \rightarrow \infty} a_n$.
3. Find a formula for the N^{th} partial sum, S_N .
4. Compute the limit of the partial sums, $\lim_{N \rightarrow \infty} S_N$ or $\sum_{n=0}^{\infty} a_n$.

Problem 280. Let $a_0 = 5$, $a_1 = \frac{5}{3}$, $a_2 = \frac{5}{9}$, $\dots$.

1. Find a formula for the n^{th} term, a_n .
2. Compute the limit of the terms, $\lim_{n \rightarrow \infty} a_n$.
3. Find a formula for the N^{th} partial sum, S_N .
4. Compute the limit of the partial sums, $\lim_{N \rightarrow \infty} S_N$ or $\sum_{n=0}^{\infty} a_n$.

Problem 281. Let $a_0 = a$, $a_1 = ar$, $a_2 = ar^2$, $\dots$.

1. Find a formula for the n^{th} term, a_n .
2. Compute the limit of the terms, $\lim_{n \rightarrow \infty} a_n$.
3. Find a formula for the N^{th} partial sum, S_N .
4. Compute the limit of the partial sums, $\lim_{N \rightarrow \infty} S_N$ or $\sum_{n=0}^{\infty} a_n$ where $r < 1$. (What happens when $r = 1$ or $r > 1$?).

In many series problems, it is useful to write out a few terms to see if there is something common that you can factor out of the sum. This is trick 49.2.b section 8 from the intergalactic “Math Book” that they give out when you get your Ph.D.

Problem 282. Determine whether these geometric series converge and if so to what number.

1. $\sum_{n=3}^{\infty} \frac{3}{4^n}$
2. $\sum_{n=3}^{\infty} \frac{2^n}{3^{n+4}}$
3. $\sum_{n=2}^{\infty} \frac{4^n}{3^n}$
4. $\sum_{n=3}^{\infty} \frac{\pi}{e^{n-1}}$

If you only do one problem in Calculus, do this one.

Problem 283. If a banker tells you that you will earn 6% interest compounded monthly, then you will actually earn $\frac{6}{12}\%$ each month, which is better than 6% annually since you get interest on your interest. Suppose you save \$300 per month and earn $\frac{6}{12}\%$ interest on the amount of money in the bank at the end of each month. How much money do you have in 30 years? How much was savings? How much was interest?

Theorem 24. The n^{th} Term Test. If $\sum_{n=k}^{\infty} a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$.

This test is essentially useless as written, but it's contrapositive is quite useful. The **contrapositive** of the statement “if P then Q” is “if not Q then not P.” To make this simple, the converse of “if you are good, then I will take you for ice cream” is “if I did not take you for ice cream, then you were not good.” Great. Now I'm hungry. We will look at a proof of the n^{th} term test later.

Problem 284. Assume that $a_n > 0$ for all $n = 1, 2, 3, \dots$. Prove that if $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series, $\sum_{n=k}^{\infty} a_n$ diverges. This is a special case of the n^{th} term test where the terms are positive.

Problem 285. In your mechanical engineering lab, you create a ball with a coefficient of elasticity so that if you dropped the ball from a height of one foot then the ball would bounce exactly to height $\frac{1}{2^n}$ feet on the n^{th} bounce. How far does the ball travel?

The n^{th} term test told us that if the terms don't approach zero, then the series diverges. This does not imply that if the terms do approach zero, then the series converges. The following problem illustrates two important points. First it shows that a series can have terms that approach zero, but still not converge. Second it shows a way to help determine whether a series converges by making a comparison between the series and a closely related integral.

Problem 286. The Harmonic Series. Let $a_1 = 1, a_2 = \frac{1}{2}, a_3 = \frac{1}{3}, \dots$

1. Find $\lim_{n \rightarrow \infty} a_n$.
2. Let $f(x) = \frac{1}{x}$ be defined on $[1, \infty)$ and sketch a graph of both f and our sequence, $a(n) = \frac{1}{n}$ for $n = 1, 2, 3, \dots$ on the same coordinate axes.
3. Use your picture to show that $S_N = \sum_{n=1}^N a_n \geq \int_1^N \frac{1}{x} dx$.

4. Show that $\int_1^N \frac{1}{x} dx$ diverges, so $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

Problem 287. Let $a_n = \frac{1}{n^2}$ for $n = 1, 2, 3, \dots$

1. Find $\lim_{n \rightarrow \infty} a_n$.

2. Use a graph to show that for each N ,

$$S_N = \sum_{n=1}^N \frac{1}{n^2} \leq 1 + \int_1^N \frac{1}{x^2} dx \leq 1 + \int_1^{\infty} \frac{1}{x^2} dx$$

so that the sequence of partial sums $\{S_n\}_{n=1}^{\infty}$ is bounded above.

3. Conclude that $\sum_{n=1}^{\infty} a_n$ converges because $\{S_n\}_{n=1}^{\infty}$ satisfies the hypothesis of Theorem 22.

Theorem 25. Integral Test. Let f be a continuous function such that $f(x) \geq 0$ and f is decreasing for all $x \geq 1$ and $f(n) = a_n$ for all $n = 1, 2, 3, \dots$. Then

$$\sum_{n=1}^{\infty} a_n \text{ converges if and only if } \int_1^{\infty} f(x) dx \text{ converges.}$$

Problem 288. Apply the integral test to each series to determine whether it converges or diverges.

1. $\sum_{n=4}^{\infty} \frac{1}{4n^3}$

2. $\sum_{n=2}^{\infty} \frac{3}{n^2}$

3. $\sum_{n=1}^{\infty} \frac{5}{n^{1.001}}$

4. $\sum_{n=1}^{\infty} \frac{4}{\sqrt{2n}}$

Problem 289. Prove the integral test as follows. Let f be a continuous function such that $f(x) \geq 0$ and is decreasing for all $x \geq 1$, and let $f(n) = a_n$. Partition the interval $[1, N]$ into $N - 1$ equal intervals.

1. Express the total area of the inscribed rectangles in sigma notation; express the total area of the circumscribed rectangles in sigma notation.

2. From (1), establish these two inequalities geometrically.

$$\sum_{n=1}^N a_n \leq a_1 + \int_1^N f(x) dx \text{ and } \int_1^N f(x) dx \leq \sum_{n=1}^N a_n$$

3. Suppose $\int_1^{\infty} f(x) dx = L$. Prove that $\sum_{n=1}^{\infty} a_n$ converges by showing that S_n is bounded above and increasing.

Theorem 26. *p*-Series Test. The series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if $p > 1$ and diverges if $p \leq 1$.

Problem 290. Use the integral test to prove the *p*-series test for the case when $p > 1$.

Problem 291. Determine whether each series converges. If it does, use Problem 289 to give an upper bound for the infinite sum.

1. $\sum_{n=4}^{\infty} \frac{5}{n^4}$
2. $\sum_{n=1}^{\infty} \frac{1}{2n^3} + \frac{2}{n^2}$
3. $\sum_{n=1}^{\infty} \frac{1}{\sqrt{2n^3}}$

We now have two types of series for which we can easily determine convergence or divergence: *geometric series* and *p-series*. We will use these to determine the convergence or divergence of many other series. If a series with positive terms is less than some convergent series, then it must converge. On the other hand, if a series is larger than a series with positive terms that diverges then it must diverge.

Problem 292. Consider the series $\sum_{n=1}^{\infty} \frac{1}{n^3 + 2}$.

1. Show that for all $n \geq 1$, we have $\frac{1}{n^3 + 2} > 0$.
2. Show that for all $n \geq 1$, we have $\frac{1}{n^3 + 2} < \frac{1}{n^3}$.
3. Why is this series monotonic and bounded and therefore convergent?

Theorem 27. Direct Comparison Test. Suppose that each of $\sum_{n=k}^{\infty} a_n$ and $\sum_{n=k}^{\infty} b_n$ are series such that $0 \leq a_n \leq b_n$ for all integers $n \geq k$.

1. If $\sum_{n=k}^{\infty} b_n$ converges, then $\sum_{n=k}^{\infty} a_n$ converges.
2. If $\sum_{n=k}^{\infty} a_n$ diverges, then $\sum_{n=k}^{\infty} b_n$ diverges.

Problem 293. Apply the direct comparison test to each of these series.

1. $\sum_{n=2}^{\infty} \frac{1}{3^n + 5}$
2. $\sum_{n=2}^{\infty} \frac{1}{3^n - 5}$
3. $\sum_{n=3}^{\infty} \frac{3}{n^2 \ln(n)}$

$$4. \sum_{n=3}^{\infty} \frac{e}{5(n^5 + 32)}$$

Example 26. Limit Comparison Test

Consider the series, $\sum_{n=1}^{\infty} \frac{n}{2^n(3n-2)}$. The terms converge to zero, so the n^{th} term test tells us nothing. It is not a geometric series either. It's not obvious what other series we might use the direct comparison test with, although that is a possibility. Still, each term of this series is the product of $\frac{1}{2^n}$ and $\frac{n}{3n-2}$ and we know that $\lim_{n \rightarrow \infty} \frac{n}{3n-2} = \frac{1}{3}$, so for large values of n , the n^{th} term is approximately equal to $\frac{1}{2^n} \cdot \frac{1}{3}$. Hence, our series behaves (for large n) a lot like a geometric series and that's a good intuitive reason to believe it converges. When we have a series like this that "looks like" a series that we know converges, in this case $\sum_{n=1}^{\infty} \frac{1}{3} \frac{1}{2^n}$, then if the limit of the ratio of the terms of the two series equals a positive number, then the series either both converge or both diverge. The next theorem makes this precise.

Theorem 28. Limit Comparison Test. Let $\{a_n\}_{n=k}^{\infty}$ and $\{b_n\}_{n=k}^{\infty}$ be sequences of positive terms such that $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ where c is a positive real number (i.e. $c = \infty$ is not allowed). Then the series $\sum_{n=k}^{\infty} a_n$ and $\sum_{n=k}^{\infty} b_n$ either both converge or both diverge.

Problem 294. Apply the limit comparison test to each series to determine if it converges or diverges.

$$1. \sum_{n=2}^{\infty} \frac{1}{3^n - 5}$$

$$2. \sum_{n=3}^{\infty} \frac{3}{n + \pi}$$

$$3. \sum_{n=1}^{\infty} \sin\left(\frac{1}{n}\right)$$

$$4. \sum_{n=1}^{\infty} \left(\frac{n+1}{2n+3}\right) \left(\frac{3^n}{5^n+1}\right)$$

Problem 295. In the real world of computer science and operations research, the "average" of a series is an important notion. Suppose we have a sequence, $\{a_n\}_{n=1}^{\infty}$ where $0 \leq a_n \leq 1$ for all positive integers, n . Consider the sequence given by

$$A_N = \frac{1}{N} \sum_{n=1}^N a_n.$$

1. Suppose $a_1 = a_2 = a_3 = \dots = 1$ Does the sequence $\{A_N\}_{N=1}^{\infty}$ converge? To what?
2. Suppose $a_1 = 1, a_2 = 0, a_3 = 1, \dots$ Does the sequence $\{A_N\}_{N=1}^{\infty}$ converge? To what?
3. Do you think that $\{A_N\}_{N=1}^{\infty}$ converges no matter what $a_1, a_2, \dots$ is? If so give an argument, if not, give an example for $a_1, a_2, \dots$ and show that for this sequence the A_n 's do not converge.

Problem 296. Determine the convergence or divergence of the following series by using any of the previously stated tests.

$$1. \sum_{n=1}^{\infty} \frac{2}{3^n + n}$$

$$2. \sum_{n=1}^{\infty} \frac{n^2 - n + 1}{2n^2 - n - 3}$$

$$3. \sum_{n=4}^{\infty} \frac{-2}{n^2 - 4n + 3}$$

$$4. \sum_{n=4}^{\infty} \frac{1}{n!}$$

$$5. \sum_{n=1}^{\infty} \cos(n\pi)$$

Problem 297. Determine the convergence or divergence of each of the following series. If any of them converge, compute the number to which it converges.

$$1. \sum_{n=1}^{\infty} \frac{|\sin(\frac{n\pi}{2})|}{2^n}$$

$$2. \sum_{n=1}^{\infty} (2n+1)^{\frac{1}{n}}$$

$$3. \sum_{n=2}^{\infty} \frac{n^2 + 2n + 1}{3n - 4}$$

$$4. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{e^n}$$

6.3 Alternating Series

We now consider series where every other term of the series is positive and every other term of the series is negative. To keep things simple, we will write each term of an alternating series as the product of two parts: the positive part, $a_n, n = 1, 2, 3, \dots$ and the alternating part, $(-1)^n$ or $(-1)^{n+1}$.

Definition 57. If $a_n > 0$ for all positive integers n , then the series $\sum_{n=1}^{\infty} (-1)^n a_n$ and the series

$\sum_{n=1}^{\infty} (-1)^{n+1} a_n$ are called **alternating series**.

Example 27. The alternating Harmonic Series.

Recall the Harmonic Series, $\sum_{n=1}^{\infty} \frac{1}{n}$, where the terms approached zero, but the series still diverged. We can modify this series to create the alternating series

$$S = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$$

By grouping the terms,

$$0 \leq (1 - \frac{1}{2}) + (\frac{1}{3} - \frac{1}{4}) + \cdots = S = 1 - (\frac{1}{2} - \frac{1}{3}) - (\frac{1}{4} - \frac{1}{5}) - \cdots \leq 1$$

so this series is between 0 and 1. The even partial sums can be written as:

$$S_2 = 1 - \frac{1}{2}, S_4 = (1 - \frac{1}{2}) + (\frac{1}{3} - \frac{1}{4}), S_6 = \dots$$

so

$$S_{2N} = (1 - \frac{1}{2}) + (\frac{1}{3} - \frac{1}{4}) + \cdots + (\frac{1}{2N-1} - \frac{1}{2N}).$$

This implies that the sequence $\{S_{2n}\}_{n=1}^{\infty}$ is increasing and bounded above by 1, so the sequence of even terms converges to some number, let's call it L . A similar argument shows that the odd terms form a decreasing sequence that is bounded below, so the odd terms converge to some number, call it G . Draw these terms on a number line and you will see that

$$S_2 < S_4 < \cdots < L \leq G < \cdots < S_5 < S_3 < S_1.$$

This means that

$$|L - G| \leq |S_N - S_{N+1}| = 1/N.$$

Since $\lim_{N \rightarrow \infty} 1/N = 0$ we have that $L = G$ so this alternating series converges.

Conclusion. This alternating series has the property that if we take the absolute value of the terms then the resulting series is the Harmonic Series which diverges. But the alternating series itself converges. We call this *conditionally convergent* and we classify all alternating series as either **divergent** (series and absolute value of series diverge), **conditionally convergent** (series converges, but absolute value of series diverges), or **absolutely convergent** (series converges and absolute value of series converges).

Definition 58. The series $\sum_{n=k}^{\infty} a_n$ is said to be **absolutely convergent** if the series $\sum_{n=k}^{\infty} |a_n|$ converges and the series $\sum_{n=k}^{\infty} a_n$ is said to be **conditionally convergent** if it converges but the series $\sum_{n=k}^{\infty} |a_n|$ diverges.

Here are two tests for checking the convergence of an alternating sequence. The first says that if the sum of the absolute value of the terms converge, then the alternating series also converges. The second test gives us another test for when an alternating series converges *and* gives us a powerful tool for saying how accurate the N^{th} partial sum is. This can tell us how many terms we need to add up in order to get an accurate approximation to the infinite sum.

Theorem 29. Absolute Convergence Test. Let $\{a_n\}_{n=k}^{\infty}$ be a sequence of positive terms. If $\sum_{n=k}^{\infty} a_n$ converges, then the alternating series $\sum_{n=k}^{\infty} (-1)^n a_n$ and $\sum_{n=k}^{\infty} (-1)^{n-1} a_n$ both converge.

Theorem 30. Alternating Series Test. Let $\{a_n\}_{n=k}^{\infty}$ be a non-increasing sequence of positive terms such that $\lim_{n \rightarrow \infty} a_n = 0$. Then

1. the alternating series $\sum_{n=k}^{\infty} (-1)^n a_n$ and $\sum_{n=k}^{\infty} (-1)^{n-1} a_n$ both converge and

2. if S is the sum of the series and S_N is a partial sum, then $|S_N - S| \leq a_{N+1}$.

Problem 298. Determine the convergence or divergence of each of the following alternating series. If the series converges, then find a positive integer N so that S_N will be within .01 of the series.

1. $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2 + 1}$

2. $\sum_{n=1}^{\infty} \sin\left(\frac{(2n-1)\pi}{2}\right)$

3. $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt[3]{n}}$

4. $\sum_{n=1}^{\infty} (-1)^n \frac{2^n}{n^2}$

5. $\sum_{n=1}^{\infty} \left(\frac{-1}{n}\right)^{n+1}$

Recall that it is possible for a series to have positive terms that approach zero, but the series still diverges – for example, the Harmonic Series. It turns out that you can have an alternating series where the limit of the terms is zero, but the alternating series still diverges. The next problem asks for you to find one!

Problem 299. Construct a sequence of non-negative terms, $\{a_n\}_{n=1}^{\infty}$, so that the $\lim_{n \rightarrow \infty} a_n = 0$ and $\sum_{n=1}^{\infty} (-1)^n a_n$ diverges.

Problem 300. Write out the first few terms for $\sum_{n=1}^{\infty} (-1)^n$. Does it converge or diverge? Why?

Problem 301. Consider the sequence of non-negative terms where $a_{2n-1} = n$ and $a_{2n} = n + \frac{1}{(n+1)^2}$. Determine whether $\sum_{n=1}^{\infty} (-1)^n a_n$ converges or diverges by writing out both the first 6 terms $a_1, a_2, a_3, a_4, a_5, a_6$ and then the first six partial sums $S_1, S_2, S_3, S_4, S_5, S_6$.

The integral, direct comparison and the limit comparison tests can only be used for series with positive terms. The next few theorems provide us with tests that we can use regardless of whether the terms are positive or not. First, let's recall the n^{th} **Term Test** which said that if $\sum_{n=1}^{\infty} a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$. Let's prove that this works for any series regardless of whether the terms are positive or not. If the series converges, let's call it $S = \sum_{n=1}^{\infty} a_n$. Since it converges, $S = \lim_{n \rightarrow \infty} S_n$. But if this is the case, then isn't it true that $S = \lim_{n \rightarrow \infty} S_{n-1}$? From this we have

$$0 = S - S = \lim_{n \rightarrow \infty} S_n - \lim_{n \rightarrow \infty} S_{n-1} = \lim_{n \rightarrow \infty} (S_n - S_{n-1}) = \lim_{n \rightarrow \infty} a_n.$$

Since we didn't use anywhere that the terms were positive, the n^{th} term test may be applied to any series whether the terms are positive or not. *q.e.d.*

The next two tests, the **Ratio Test** and the **Root Test** also work on series regardless of whether the terms are positive or not.

Theorem 31. Ratio Test. Let $\{a_n\}_{n=k}^{\infty}$ be a sequence such that $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = r$ (where we allow the case $r = \infty$). Then

- the series $\sum_{n=k}^{\infty} a_n$ converges absolutely if $r < 1$,
- the series $\sum_{n=k}^{\infty} a_n$ diverges if $r > 1$ or $r = \infty$, and
- the test fails if $r = 1$.

Sketch of Proof of Ratio Test. This is a quite advanced proof, worthy of a course in Analysis. If you read it and are curious, I'll happily discuss it or fill in the missing details. Let's consider the case where $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = r$ and $r < 1$. Since $\left| \frac{a_{n+1}}{a_n} \right|$ is getting closer and closer to r then we can choose a positive integer N large enough to assure that $\left| \frac{a_{N+1}}{a_N} \right| < 1$. Therefore we can find a positive real number R so that $|a_{N+1}| < R|a_N|$. A little bit of algebra shows that for all positive integers k we have $|a_{N+k}| < R^k|a_N|$. This implies that $\sum_{k=1}^{\infty} |a_{N+k}|$ converges and we can conclude that $\sum_{n=1}^{\infty} a_n$ converges absolutely. *q.e.d.*

Problem 302. Apply the ratio test to each series in an attempt to determine the convergence or divergence of each series. If the ratio test fails, apply another test to determine convergence or divergence.

1. $\sum_{n=2}^{\infty} \frac{n^2}{n!}$
2. $\sum_{n=3}^{\infty} \frac{2^n}{n^3}$
3. $\sum_{n=3}^{\infty} \frac{\ln(n)}{n}$

Problem 303. Using any test you like, determine whether the series is absolutely convergent, conditionally convergent, or divergent.

1. $\sum_{n=1}^{\infty} \frac{(-1)^n}{\ln(n+1)}$
2. $\sum_{n=1}^{\infty} (-1)^n \frac{n!}{3^{n+1}}$
3. $\sum_{n=1}^{\infty} (-1)^n \frac{n^2+3}{4n^3}$

Here is one last test that the pundits believe you should be aware of.

Theorem 32. Root Test. Let $\{a_n\}_{n=k}^{\infty}$ be a sequence such that $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = r$ (where we allow the case $r = \infty$). Then

- the series $\sum_{n=k}^{\infty} a_n$ converges absolutely if $r < 1$,
- the series $\sum_{n=k}^{\infty} a_n$ diverges if $r > 1$ or $r = \infty$, and
- the test fails if $r = 1$.

6.4 Power Series

From the first semester of calculus, we know that the best linear (1st degree polynomial) approximation to the function, $f(x) = e^x$ at the point $P = (2, e^2)$ is the tangent line to the function at this point. Let's call that approximation L and note that $L(2) = f(2)$ and $L'(2) = f'(2)$. That is, L and f intersect at P and L and f share the same derivative at P . What is the best quadratic (2nd degree polynomial) approximation? If the best first degree approximation to the curve agrees at the point and in the first derivative, then the best second degree approximation should agree with the function at the point, in the first derivative, and in the second derivative.

Example 28. Compute the best linear and quadratic approximations to $f(x) = x^3 - 16x$.

Problem 304. Let $f(x) = e^x$. Find the best linear approximation, $L(x) = mx + b$, to f at $(2, e^2)$. Find the best quadratic approximation, $Q(x) = ax^2 + bx + c$, to f at $(2, e^2)$. Graph all three functions on the same pair of coordinate axes.

Problem 305. Let $f(x) = \cos(x)$. Find the best linear (L , 1st degree), quadratic (Q , 2nd degree), and quartic (C , 4th degree) approximations to f at $(\pi, \cos(\pi))$. Sketch the graph of f and all three approximations on the same pair of coordinate axes. Compute and compare $f(\frac{3\pi}{2})$, $L(\frac{3\pi}{2})$, $Q(\frac{3\pi}{2})$, and $C(\frac{3\pi}{2})$.

From our work on geometric series, we know that

$$\sum_{n=0}^{\infty} t^n = \frac{1}{1-t}$$

for $|t| < 1$. Therefore the function $g(t) = \frac{1}{1-t}$ where $|t| < 1$ can be written as a series,

$$g(t) = \frac{1}{1-t} = 1 + t + t^2 + t^3 + \cdots = \sum_{n=0}^{\infty} t^n.$$

Replacing t by $-x^2$, we have

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - \cdots + (-1)^{n-1} x^{2n-2} + \cdots.$$

Based on these two observations, we see that we can write at least two rational functions as infinite series (or infinite polynomials). And we can write every polynomial as an infinite series, since

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_Nx^N = \sum_{i=0}^{\infty} a_i x^i$$

where $a_i = 0$ for $i > N$. Our goal is a systematic way to write any differentiable function (like $\sin$ or $\cos$ or $\ln$) as an infinite series.

Definition 59. $0^0 = 1$

Notation. To keep from confusing the powers of f with the derivatives of f , we use $f^{(n)}$ to represent f if $n = 0$ and the n^{th} derivative of f if $n \geq 1$. Therefore when $n \geq 1$, $f^{(n)}(c)$ means the n^{th} derivative of f evaluated at the number c .

Problem 306. Let $f(x) = \sum_{i=0}^{\infty} a_i x^i$. Write out S_6 , the 6th partial sum of this series. Compute S'_6 and S''_6 . Compute $S'_6(0)$ and $S''_6(0)$.

Problem 307. Let $f(x) = \sum_{i=0}^{\infty} a_i x^i$. Compute f' and f'' . Compute $f(0)$, $f'(0)$, $f''(0)$, ...? If n is any positive integer, what is the n^{th} derivative at 0, $f^{(n)}(0)$?

Definition 60. If f is a function with N derivatives, then the N^{th} **degree Taylor polynomial of f expanded at c** is defined by

$$T_N(x) = \sum_{n=0}^N \frac{f^{(n)}(c)}{n!} (x - c)^n.$$

Example 29. Compute the Taylor series polynomials of degree 1, 2, 3 and 4 for $f(x) = 1/x$ expanded at 1. Compute the Taylor series polynomial for the same function expanded at 1.

Problem 308. Suppose that $T(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$. Write out the first few terms of this series and compute $f'(c)$, $f''(c)$, $f'''(c)$, ...

Definition 61. The series in the last problem is called the **Taylor Series for f expanded at c** . When $c = 0$ this is called the **McLaurin Series for f** .

Problem 309. Let $f(x) = e^x$. Compute the first, second, and third degree Taylor polynomials for f expanded at 2. Compare to the result of problem 304.

Example 30. Compute the Taylor series for $f(x) = 1/x$ at $c = 1$ and use the Ratio Test to determine the interval of convergence. Also check convergence at both endpoints of the interval of convergence.

Problem 310. For each function, write the infinite degree Taylor polynomial as a series.

1. $f(x) = e^x$ expanded at 0
2. $f(x) = \sin(x)$ expanded at 0
3. $f(x) = \frac{\cos(x)}{x}$ expanded at 0 (Divide the series for $\cos(x)$ by x .)

Problem 311. Let $f(x) = \ln(x)$. Compute the first, second, and third degree Taylor polynomial for f expanded at 1. Compute $|f(2.5) - T_N(2.5)|$ for $N = 1, 2, 3$.

We now can write certain functions as infinite series, but we know that not every infinite series converges. Therefore, our expressions only make sense for the values of x for which the series converges. For this reason, every time we write down an infinite series to represent a function, we need to determine the values of x for which the series converges. This is called interval of convergence (or domain) of the series.

Definition 62. A **power series expanded at c** is any series of the form $\sum_{n=0}^{\infty} a_n(x - c)^n$. Taylor and McLaurin series are special cases of power series. The largest interval for which a power series converges is called its **interval of convergence**. Half the length of this interval is called the **radius of convergence**.

Problem 312. Use the ratio test to find the interval and radius of convergence for each power series.

$$1. \sum_{n=0}^{\infty} \frac{x^n}{n+1}$$

$$2. \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$3. \sum_{n=0}^{\infty} n!(x+2)^n$$

The next theorem says that the interval of convergence of a power series expanded at the point c is either (i) only the one point, c (bad, since this means the series is useless), (ii) an interval of radius R centered at c , (good, and the endpoints might be contained in the domain), or (iii) all real numbers (very good).

Theorem 33. Power Series Theorem. For the power series $\sum_{n=0}^{\infty} a_n(x - c)^n$, one and only one of the following is true.

1. The series converges only at c . (bad)

2. There exists a positive number R called the **radius of convergence** such that $\sum_{n=0}^{\infty} a_n(x - c)^n$ converges absolutely for $|x - c| < R$. (good)

3. The series converges absolutely for all x . (great)

When we write the power series for a function (like e^x) and it converges for all values of x then we simply have a different way to write the function that turns out to be both computationally useful, because of convergence, and theoretically useful. Speaking loosely, e^x is just an infinite polynomial – how cool is that?!

Problem 313. Find the interval of convergence for each of the following power series.

$$1. \sum_{n=0}^{\infty} \frac{(x-3)^n}{\ln(n+2)}$$

$$2. \sum_{n=1}^{\infty} \frac{(-x)^n}{n}$$

$$3. \sum_{n=0}^{\infty} \frac{\log(n+1)}{(n+1)^4} (x+1)^n$$

$$4. \sum_{n=0}^{\infty} \frac{n^3}{3^n} (x+2)^n$$

The following theorem says that if you have a power series, then its derivative is exactly what you want it to be! Just take the derivative of the series term by term and the new infinite series you get is the derivative of the first and is defined on the same domain.

Theorem 34. Power Series Derivative Theorem. If $f(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$ for all x in the interval of convergence $(c-R, c+R)$, then

1. f is differentiable in $(c-R, c+R)$,
2. $f'(x) = \sum_{n=1}^{\infty} na_n(x-c)^{n-1}$ for all x in $(c-R, c+R)$, and
3. the interval of convergence of f' is also $(c-R, c+R)$.

Problem 314. Suppose that $f(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$ with radius of convergence $r > 0$. Show that a_n must equal $\frac{f^{(n)}(c)}{n!}$ for $n = 0, 1, 2$, and 3 .

How does the calculator on your phone approximate numbers like $\sqrt{5}$ or $\sin(\pi/7)$? It uses Taylor series. For example, if we want to approximate $\sqrt{5}$ accurate to 3 decimal places, we could compute the Taylor series for $f(x) = \sqrt{x}$ expanded about $c = 4$ because 4 is the integer closest to 5 for which we actually know the value of $\sqrt{4}$. Then we can use this polynomial to approximate $\sqrt{5}$. But there's a problem. While the Taylor series approximates the function, it's pretty hard to store an infinite polynomial in a phone. So, how can we determine what degree Taylor polynomial we need to compute in order to assure a given accuracy? We look at the remainder of the series. For any positive integer, N , f can be rewritten as the sum of the Taylor polynomial of degree N plus the remainder of the series by writing,

$$f(x) = \sum_{n=0}^N \frac{f^{(n)}(c)}{n!} (x-c)^n + \sum_{n=N+1}^{\infty} \frac{f^{(n)}(c)}{n!} (x-c)^n$$

or

$$f(x) = T_N(x) + R_N(x),$$

where

$$T_N(x) = \sum_{n=0}^N \frac{f^{(n)}(c)}{n!} (x-c)^n \text{ is the } N^{\text{th}} \text{ degree Taylor polynomial}$$

and

$$R_N(x) = \sum_{n=N+1}^{\infty} \frac{f^{(n)}(c)}{n!} (x-c)^n \text{ is called the } N^{\text{th}} \text{ degree Taylor remainder.}$$

Since $f(x) = T_N(x) + R_N(x)$, for all x in the interval of convergence, the error for any x in the interval of convergence is just $|f(x) - T_N(x)| = |R_N(x)|$. If we could just approximate the size of the remainder term, we would know how accurate the N^{th} degree Taylor polynomial was. The next theorem says that the remainder, even though it is an infinite sum, can be written completely in terms of the $(N+1)^{\text{st}}$ derivative, so if we can find a bound for the $(N+1)^{\text{st}}$ derivative then we can get a bound on the error of the Taylor series.

Theorem 35. Taylor Series Error Theorem. Let f be a function such that $f^{(i)}$ is continuous for each $i = 0, 1, 2, \dots, N$ on $[a, b]$ and $f^{(N+1)}(x)$ exists for all x in (a, b) . Then $R_N(x) = \frac{f^{(N+1)}(k)}{(N+1)!}(x - c)^{N+1}$ for some k between x and c .

Problem 315. Let $f(x) = \sqrt{x}$ and compute $T_3(x)$, the third degree Taylor series for f expanded at 4. Estimate the error $|f(5) - T_3(5)|$ by using Theorem 35 to find the maximum of $|R_3(5)|$. Now use your calculator to compute $|f(5) - T_3(5)|$ and see how this compares to your error estimate.

Problem 316. Let $f(x) = \sin(x)$ and compute $T_3(x)$, the third degree McLaurin series f . Estimate the error $|f(\frac{\pi}{7}) - T_3(\frac{\pi}{7})|$ by using Theorem 35 to find the maximum of $|R_3(\frac{\pi}{7})|$. Now use your calculator to compute $|f(\frac{\pi}{7}) - T_3(\frac{\pi}{7})|$ and see how this compares to your error estimate.

Chapter 7

Conic Sections, Parametric Equations, and Polar Coordinates

“Learning is not compulsory... neither is survival.” - W. Edwards Deming

7.1 Conic Sections

In your mathematical life, you have no doubt seen circles, ellipses, parabolas, and possibly even hyperbolas. Here we make precise the geometric and algebraic definitions of these objects.

Definition 63. Consider two identical, infinitely tall, right circular cones placed vertex to vertex so that they share the same axis of symmetry (two infinite ice cream cones placed point-to-point). A **conic section** is the intersection of this three dimensional surface with any plane that does not pass through the vertex where the two cones meet.

These intersections are called circles (when the plane is perpendicular to the axis of symmetry), parabolas (when the plane is parallel to one side of one cone), hyperbolas (when the plane is parallel to the axis of symmetry), and ellipses (when the plane does not meet any of the three previous criteria). This is a geometric interpretation of the conic sections, but each conic sections may also be represented by a quadratic equation in two variables.

Problem 317. Algebraically. Graph the set of all points (x,y) in the plane satisfying $Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$ where:

1. $A = B = C = 0, D = 1, E = 2$ and $F = 3$.
2. $A = 1/16, C = 1/9, B = D = E = 0$ and $F = -1$.
3. $A = B = C = D = E = 0$ and $F = 1$.
4. $A = B = C = D = E = F = 0$.

Every conic section may be obtained by making appropriate choices for A, B, C, D, E , and F . Now let's return to the definition you might have seen in high school for a circle.

Definition 64. Given a plane, a point (h,k) in the plane, and a positive number r , the **circle** with **center** (h,k) and **radius** r is the set of all points in the plane at a distance r from (h,k) .

Problem 318. Determine values for $A, B, \dots, E$ and F so that $Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$ represents a circle. A single point. A parabola.

Problem 319. Use the method of completing the square to rewrite the equation $4x^2 + 4y^2 + 6x - 8y - 1 = 0$ in the form $(x - h)^2 + (y - k)^2 = r^2$ for some numbers h, k , and r . Generalize your work to rewrite the equation $x^2 + y^2 + Dx + Ey + F = 0$ in the same form.

Definition 65. The **distance between a point P and a line L** (denoted by $d(P,L)$) is defined to be the length of the line segment perpendicular to L that has one end on L and the other end on P .

Definition 66. Given a point P and a line L in the same plane, the **parabola** defined by P and L is the set of all points in the plane Q so that $d(P,Q) = d(Q,L)$. The point P is called the **focus** of the parabola and the line L is called the **directrix** of the parabola.

Problem 320. Parabolas.

1. Let p be a positive number and graph the parabola that has focus $(0, p)$ and directrix $y = -p$.
2. Let (x, y) be an arbitrary point of the parabola with focus $(0, p)$ and directrix $y = -p$. Show that (x, y) satisfies $x^2 = 4py$.
3. Let (x, y) be an arbitrary point of the parabola with focus $(p, 0)$ and directrix $x = -p$. Show that (x, y) satisfies $y^2 = 4px$.
4. Graph the parabola, focus, and directrix for the parabola $y = -(x - 2)^2 + 3$.

Problem 321. Find the vertex, focus, and directrix of each of the parabolas defined by the following equations.

1. $y = 3x^2 - 12x + 27$
2. $y = x^2 + cx + d$ (Assume that c and d are numbers.)

Definition 67. Given two points, F_1 and F_2 in the plane and a positive number d , the **ellipse** determined by F_1, F_2 and d is the set of all points Q in the plane so that $d(F_1, Q) + d(F_2, Q) = d$. F_1 and F_2 are called the **foci** of the ellipse. The **major axis** of the ellipse is the line passing through the foci. The **vertices** of the ellipse are the points where the ellipse intersects the major axis. The **minor axis** is the line perpendicular to the major axis and passing through the midpoint of the foci.

Problem 322. Let $F_1 = (c, 0), F_2 = (-c, 0)$, and $d = 2a$. Show that if (x, y) is a point of the ellipse determined by F_1, F_2 and d , then (x, y) satisfies the equation $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ where $c^2 = a^2 - b^2$. What are the restrictions on a, b , and c ?

Problem 323. Ellipses.

1. Graph the ellipses defined by $\frac{x^2}{25} + \frac{y^2}{9} = 1$ and $\frac{(x+1)^2}{25} + \frac{(y-2)^2}{9} = 1$. List the foci for each, the vertices for each, and the major axis for each. Compare the two graphs and find the similarities and differences between the two graphs.
2. For the ellipses defined by $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$, list the foci for each, the vertices for each, and the major axis for each.

Definition 68. Given two points, F_1 and F_2 in the plane and a positive number d , the **hyperbola** determined by F_1, F_2 and d is the set of all points Q in the plane so that $|d(F_1, Q) - d(F_2, Q)| = d$. F_1 and F_2 are called the **foci** of the hyperbola.

Problem 324. Let $F_1 = (c, 0), F_2 = (-c, 0)$, and $d = 2a$. Show that the equation of the hyperbola determined by F_1, F_2 and d is given by $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ where $c^2 = a^2 + b^2$. What are the restrictions on a, b , and c ?

Problem 325. Graph two hyperbolas (just pick choices for a and b). Solve the equation

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

for y . Show that as $x \rightarrow \infty$ the hyperbola approaches the lines $L(x) = \pm \frac{b}{a}x$; i.e., show that the asymptotic behavior of a hyperbola is linear.

Problem 326. Hyperbolas.

1. Graph $\frac{x^2}{25} - \frac{y^2}{9} = 1$ and $\frac{(x+1)^2}{25} - \frac{(y-2)^2}{9} = 1$. Compare the two graphs and find the similarities and differences between the two hyperbolas. Find the foci and the equations of the asymptotes of these hyperbolas.
2. Find the vertices and the asymptotes of $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ and $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$.

Problem 327. Use the method of completing the square to identify whether the set of points represents a circle, a parabola, an ellipse, or a hyperbola. Sketch the conic section and list the relevant data for that conic section from this list: center, radius, vertex (vertices), focus (foci), directrix, and asymptotes.

1. $x^2 - 4y^2 - 6x - 32y - 59 = 0$
2. $9x^2 + 4y^2 + 18x - 16y = 11$
3. $x^2 + y^2 + 10 = 6y - 4 + 4x$
4. $x^2 + y^2 + 6x - 14y = 6$
5. $3x^2 - 2x + y = 5$

7.2 Parametric Equations

Parametric equations represent a mathematical way to model the position of an object in space at a given time, for example, a satellite traveling through space or a fish swimming in the ocean. In this section, we restrict our study to objects traveling in the plane. In the next chapter we will extend this notion to three-space.

Example 31. Define two functions x and y on the interval $[0, 2\pi]$ by $x(t) = r \cos(t)$ and $y(t) = r \sin(t)$. Now define C on the same interval by $C(t) = (x(t), y(t))$. If we assume that t represents time, then we may consider $C(t)$ to represent the position of some object at that time. $C(t) = (x(t), y(t))$ is called a **parametric equation** with **parameter** t and defines a function from $[0, 2\pi]$ into $\mathbb{R}^2$. C is a way to represent the circle centered at $(0, 0)$ with radius r since for any t we have $(x(t))^2 + (y(t))^2 = r^2 \cos^2(t) + r^2 \sin^2(t) = r^2$. In this mathematical model, each point on the circle models the position of the object at a specific time.

Definition 69. If each of f and g is a continuous function, then the curve in the plane defined by the range of C , where $C(t) = (f(t), g(t))$, is called a **parametric curve** or a **planar curve**.

Problem 328. Sketch each pair of planar curves by plotting points. Compare.

1. $C(t) = (t, \sin(t))$ and $D(t) = (\sin(t), t)$ with $0 \leq t \leq 2\pi$

2. $C(t) = (2\sin(t), 3\cos(t))$ and $D(t) = (2\cos(t), 3\sin(t))$ with $0 \leq t \leq 2\pi$

Problem 329. Sketch the planar curve $C(t) = (3t + 2, 9t^2 + 1)$. Sketch the tangent line to this curve when $t = 1$. Write the equation of this tangent line as a function $y = mx + b$ and as a planar curve, $D(t) = (---, ---)$.

Definition 70. If $C(t) = (f(t), g(t))$ is a parametric equation where both of f and g are differentiable at the point t , then the **derivative** of C is defined by $C'(t) = (f'(t), g'(t))$.

Problem 330. Let $C(t) = (2t + 3, 4(2t + 5)^2)$.

1. Sketch a graph of C between the times $t = 0$ and $t = 2$.

2. Compute $C'(t)$ and $C'(1)$.

Problem 331. Let $C(t) = (2t + 3, 4(2t + 5)^2)$.

1. Let $x = 2t + 3$ and $y = 4(2t + 5)^2$ and eliminate the parameter, t , to write this in terms of x and y only.

2. Compute the derivative of y with respect to x and compute y' when $x = 5$.

3. Compute the derivative of y with respect to t and compute y' when $t = 1$.

4. Compute the derivative of x with respect to t and compute x' when $t = 1$.

5. Observe that the derivative of y with respect to x at $x = 5$ equals the quotient of the derivative of y with respect to t at $t = 1$ and the derivative of x with respect to t when $t = 1$.

The preceding problem illustrates an important theorem that relates the derivatives of two different, but related, functions:

1. the derivative of the parametric curve $c = (f, g)$ and

2. the derivative of the function y that we obtain when we eliminate the parameter.

Theorem 36. Parametric Derivative Theorem. If $c = (f, g)$ and y is the function obtained by eliminating the parameter, then $y'(f(t)) = \frac{g'(t)}{f'(t)}$. Since $f(t)$ represents our x coordinate, many books would write this as $y'(x) = \frac{g'(t)}{f'(t)}$.

To justify this theorem, we offer three levels of “proof.” An intuitive argument, a slightly more precise argument, and an actual proof.

Intuitively. Suppose we have a parametric equation, $C(t) = (f(t), g(t))$ where f and g are differentiable functions. If we let $x = f(t)$ and $y = g(t)$ then $y'(x)$ is the rate of change of y with respect to x . The rate of change of y with respect to time is $g'(t)$ and the rate of change of x with respect to time is $f'(t)$. The rate of change of y over the rate of change of x is $y'(x) = \frac{g'(t)}{f'(t)}$.
q.e.d.

More precisely. Suppose we have a parametric equation,

$$C(t) = (f(t), g(t))$$

where f and g are differentiable functions. If we can eliminate the parameter t (as we did in the last problem), then we can write a relationship that yields a differentiable function, y so that for every t ,

$$y(f(t)) = g(t).$$

Then using implicit differentiation, we can see that

$$y'(f(t)) \cdot f'(t) = g'(t).$$

Dividing through by $f'(t)$ we have the desired result,

$$y'(f(t)) = \frac{g'(t)}{f'(t)}.$$

If we put $x = f(t)$, then we have

$$y'(x) = \frac{g'(t)}{f'(t)}.$$

q.e.d.

Before proceeding to give a true proof of this theorem, we need one more tool, which the next problem illustrates.

Notation. The expression $^{-1}$ means *inverse* when applied to a function (see the left-hand side of the equation in part 2 of Problem 332) and *reciprocal* when applied to a number (see the right-hand side of the equation in part 2 of Problem 332). When a function has an inverse, we say it is invertible.

Problem 332. Let $f(x) = x^3 - 1$ and $(x, y) = (3, 26)$.

1. Compute f' and f^{-1} .
2. Show that $(f^{-1})'(26) = (f'(3))^{-1}$.

Theorem 37. Inverse Derivative Theorem. Let f be a function that is differentiable, and has an inverse that is differentiable. If $y = f(x)$ then $(f^{-1})'(y) = (f'(x))^{-1}$.

Problem 333. Prove Theorem 37 by computing the derivative of $f^{-1}(f(x)) = x$.

Now that we have Theorem 37 we can give a full proof of Theorem 36.

A Proof. Suppose we have a parametric equation,

$$C(t) = (f(t), g(t))$$

where f and g are differentiable functions whose derivatives are continuous. Suppose further that there is a value t^* where $f'(t^*) \neq 0$. Now, since $f'(t^*) \neq 0$ it is either positive or negative. Let's assume it's positive. (We also need to do the case where it is negative, but it turns out that the argument is almost identical, so we omit it.) Since f' is continuous and $f'(t^*) > 0$, there must be some open interval (a, b) containing t^* where $f'(t) > 0$ for all t in (a, b) . Now, since f' is positive on this interval f must be increasing on this interval and therefore f must be invertible on this interval. Suppose (t, x) is a point of f with t in (a, b) . Thus, $t = f^{-1}(x)$ and x is in $(f(a), f(b))$. Now, let's define a new function, y by

$$y(x) = g(f^{-1}(x)).$$

Computing the derivative via the chain rule as we did in the last ‘proof’ yields,

$$y'(x) = g'(f^{-1}(x)) \cdot (f^{-1})'(x).$$

Applying Theorem 37, we know that

$$(f^{-1})'(y) = (f'(x))^{-1}$$

so we have

$$y'(x) = g'(f^{-1}(x)) \cdot (f^{-1})'(x) = g'(t) \cdot \frac{1}{f'(t)} = \frac{g'(t)}{f'(t)}.$$

q.e.d.

Problem 334. Assume that C , f , and g are as above and compute $y''(x)$ to show that

$$y''(x) = \frac{g''(t)f'(t) - g'(t)f''(t)}{(f'(t))^3}.$$

What we really want to compute is the derivative of the function

y' and we know that $y'(f(t)) = \frac{g'(t)}{f'(t)}$, so compute the derivative of both sides of this equation with respect to t .

The next theorem summarizes our results about derivatives of parametric equations.

Theorem 38. Parametric Second Derivative Theorem. Suppose we have a parametric equation, $C(t) = (f(t), g(t))$ where f and g are differentiable functions. Then, at any value of t where $f'(t) \neq 0$ if we let $x = f(t)$ and $y = g(t)$ we have:

1. $y'(x) = \frac{g'(t)}{f'(t)}$ and
2. $y''(x) = \frac{g''(t)f'(t) - g'(t)f''(t)}{(f'(t))^3}.$

This theorem is often written in textbooks using Leibnitz notation as:

1. $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$
2. $\frac{d^2y}{dx^2} = \frac{\frac{d^2y}{dt^2}}{\frac{dx}{dt}}$ where $y' = \frac{dy}{dx}$ and $\frac{dx}{dt} \neq 0$.

Problem 335. Let $C(t) = (t^2, t^3)$. Eliminate the parameter to write this parametric equation as a function y in terms of x . Compute each of y' and y'' at $t = 2$ both directly and by applying the theorem.

Theorem 39. Arc Length Theorem. Suppose we have a parametric equation, $C(t) = (f(t), g(t))$ where f and g are differentiable functions with continuous derivatives for all $a \leq t \leq b$. If the curve C is traversed exactly once as t increases from a to b , then the curve's **arc length** is given by

$$L = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt.$$

This is one of the theorems in calculus with a very nice straightforward proof. If I don't remember to show you the proof, ask!

Problem 336. Let $C(t) = (2t, t^2)$ from $t = 0$ to $t = 2$. Use the arc length formula to determine the length of this curve. Now, eliminate the parameter to write this parametric equation as a function f in terms of x . Use the formula we had for the arc length of a function, $L = \int_a^b \sqrt{1 + (f'(x))^2} dx$, to verify your answer.

Problem 337. Let $C(t) = (5\cos(t) - \cos(5t), 5\sin(t) - \sin(5t))$. Compute the arc length of the range of C over the interval $[0, 2\pi]$.

7.3 Polar Coordinates

Like the Cartesian (rectangular) coordinate system that you are familiar with, the polar coordinate system is a way to specify the location of points in the plane. Some curves have simpler equations in polar coordinates, while others have simpler equations in the Cartesian coordinate system.

Example 32. Convert a few points from polar coordinates to rectangular coordinates and vice-versa – perhaps $(3, 4)$, $(0, 1)$, $(5\pi/6, 2)$, and $(3\pi/2, -3)$. Convert a few equations from polar to rectangular coordinates and vice-versa – perhaps $5x - 3y = 9$, $3x^2 = y$, $r = 2\sin(\theta)$, and $r = 4$.

Definition 71. Given a point P in the plane we may associate P with an ordered pair (r, θ) where r is the distance from $(0, 0)$ to P and θ is the angle between the x -axis and the ray $\overrightarrow{OP}$ as measured in the counter-clockwise direction. The pair (r, θ) is called a **polar coordinate representation** for the point P .

Warning. There are many polar coordinate representations for a point. For example, if (r, θ) is a polar coordinate representation for some point in the plane, then by adding 2π to the angle, $(r, \theta + 2\pi)$ is a second polar coordinate representation for the same point. We also allow $r \leq 0$ so that by negating r and adding π to the angle, $(-r, \theta + \pi)$ is a third polar coordinate representation for the same point.

Problem 338. Plot the following points in the polar coordinate system and find their coordinates in the Cartesian coordinate system.

1. $(1, \pi)$
2. $\left(3, \frac{5\pi}{4}\right)$
3. $\left(-3, \frac{\pi}{4}\right)$
4. $\left(-2, \frac{-\pi}{6}\right)$

Problem 339. Suppose (r, θ) is a point in the plane in polar coordinates and (x, y) are the coordinates in the Cartesian coordinate system. Write formulas for x and y in terms of r and θ .

Problem 340. Suppose (x, y) is a point in the plane in Cartesian coordinates. Write formulas (in terms of x and y) for

1. the distance r from the origin to (x, y) and
2. the angles between the positive x -axis and the line containing the origin and (x, y) .

Problem 341. Graphing.

1. Graph the curve $y = \sin(x)$ for x in $[0, 2\pi]$ in the Cartesian coordinate system.
2. Graph the curve $r = \sin(\theta)$ for θ in $[0, 2\pi]$ in the polar coordinate system.

Problem 342. Each of the following equations is written in the Cartesian coordinate system. Use the relationships you found in Problem 339 to write each in polar coordinates.

1. $2x - 3y = 6$
2. $x^2 + y^2 = 9$
3. $25x^2 + 4y^2 = 100$
4. $x = 4y^2$
5. $xy = 36$

Problem 343. Each of the following equations is written in the polar coordinate system. Use the relationships you found in Problem 340 to write each in Cartesian coordinates.

1. $r = 9\cos(\theta)$
2. $r^2 = 2\sin(\theta)$
3. $r = \frac{1}{3\cos(\theta) - 2\sin(\theta)}$
4. $r = 8$
5. $\theta = \frac{\pi}{4}$

Problem 344. Graph, in a polar coordinate system, the curve defined by each of the following equations.

1. $r = 1 + 2\cos(\theta)$
2. $r = 2 + 2\sin(\theta)$
3. $r = 3\sin(2\theta)$
4. $r = 2\cos(3\theta)$
5. $r^2 = 4\sin(2\theta)$ (If you know how to use a graphing calculator to plot polar curves, try it.)

Problem 345. Find the points of intersection of the graphs of the given polar equations. (When I set them equal without graphing them, I always miss at least one point of intersection.)

1. $r = 4 - 4\cos(\theta)$ and $r = 4\cos(\theta)$
2. $r = 2\cos(2\theta)$ and $r = \sqrt{3}$
3. $r = \sin(\theta)$ and $r = \sqrt{3}\cos(\theta)$
4. $r = 4 + 2\cos(\theta)$ and $r = 5$
5. $r = 1 - 2\cos(\theta)$ and $r = 1$

Example 33. Derive the formula for the arc length of a function starting with the distance formula. Derive the formula for the arc length of a planar curve defined by a parametric equation.

Example 34. Compute the arc length (using the equation they will derive next) for the spiral $r = \theta$ from $\theta = 0$ to $\theta = 2\pi$. Compute the area of one of the four “petals” of $r = 3\sin(2\theta)$.

Problem 346. If we display the independent variable θ , then Cartesian and polar coordinates are related by $x(\theta) = r(\theta)\cos(\theta)$ and $y(\theta) = r(\theta)\sin(\theta)$. Use the formula for the arc length of a parametric curve

$$L = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt$$

to derive the formula for the arc length of a curve in polar coordinates

$$L = \int_{\alpha}^{\beta} \sqrt{(r(\theta))^2 + (r'(\theta))^2} d\theta.$$

Problem 347. Use the arc length formula from the previous problem and the half-angle identity to find the length of the cardioid defined by $r = 3(1 + \cos(\theta))$.

Problem 348. Recall that $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$. Use this to find the tangent line to the curve in a polar plane $r = 2 + 2\sin(\theta)$ at $\theta = \pi/4$.

Now we can find arc length and tangent lines to curves written in polar coordinates. How about finding the area bounded by a polar curve?

Let $r = f(\theta)$ be a continuous and non-negative function in the interval $[\alpha, \beta]$ and A be the area of the region bounded by the graph of f and the radial lines $\theta = \alpha$ and $\theta = \beta$. To find A , partition the interval $[\alpha, \beta]$ into n subintervals $[\theta_0, \theta_1], [\theta_1, \theta_2], \dots, [\theta_{n-1}, \theta_n]$ of equal lengths where $\alpha = \theta_0 < \theta_1 < \dots < \theta_n = \beta$. Since the area of a circular sector of radius r and angle θ is $\frac{1}{2}r^2\theta$, the area of each of our sectors is $\frac{1}{2}(\theta_i - \theta_{i-1})^2$ so the total area is approximately the sum of these sectors. Thus $A = \lim_{n \rightarrow \infty} \frac{1}{2} \sum_{i=1}^n [f(\theta_i)]^2 (\theta_i - \theta_{i-1}) = \frac{1}{2} \int_{\alpha}^{\beta} [f(\theta)]^2 d\theta$.

Problem 349. Find the area of each region in a polar plane.

1. the region bounded by the graph of $r = -2\cos(\theta)$
2. the region bounded by one loop of the graph of the equation $r = 4\cos(2\theta)$
3. the region bounded by the inner loop of the curve $r = 2 + 3\cos(\theta)$
4. the region bounded by the intersection of the two graphs $r = 4\sin(2\theta)$ and $r = 4\cos(2\theta)$
5. the region that lies inside the cardioid $r = 2 - 2\cos(\theta)$ and outside the circle $r = 1$

Chapter 8

Vectors and Lines

“The only way of discovering the limits of the possible is to venture a little way past them into the impossible.” - Arthur C. Clarke

Welcome to calculus in three dimensions. The beauty of this material is how closely it parallels your first semester of calculus. After a brief introduction to the coordinate plane, you learned how to graph lines and parabolas. After a brief introduction to three-space, we will be graphing planes and paraboloids. Just as we defined continuity in terms of limits in Calculus I, we will define continuity of functions of several variables in terms of limits in this course. Lines were important as they allowed you to define tangent lines to functions and the derivative. Tangent planes to functions and surfaces will aid our definition of derivative. Once you understood the derivative, you used it to find maxima and minima of real valued functions and we will use the derivative of functions in three-space to find maxima and minima in our applications as well. Two of the most central ideas of your first calculus course were the chain rule and the fundamental theorem of calculus. We will extend your notion of the chain rule and the fundamental theorem in this course. Only near the end of the course will the work we do not have a parallel to your first course. At the end, we'll cover Green's and Gauss' Theorems, which are necessary tools in physics and engineering, but which had no parallel in Calculus I.

Definition 72. $\mathbb{N}$ is the set of all **Natural Numbers**.

Definition 73. $\mathbb{R}$ is the set of all **Real Numbers**.

We will also use the notation, “ $\in$,” to mean “is an element of.” Thus, “ $x \in \mathbb{R}$ ” means “ x is an element of $\mathbb{R}$,” or “ x is a real number.” Similarly, $x, y \in \mathbb{R}$ means “ x and y are real numbers.” In Calculus I and II, you lived in two-space, or $\mathbb{R}^2 = \{(x, y) : x, y \in \mathbb{R}\}$. Now you have graduated to 3-space!

Definition 74. *Three dimensional space (Euclidean 3-space or $\mathbb{R}^3$), is the set of all ordered sequences of 3 real numbers. That is*

$$\mathbb{R}^3 = \{(x_1, x_2, x_3) : x_1, x_2, x_3 \in \mathbb{R}\}.$$

Of course, there is no reason to stop with the number 3. More generally, n -space or $\mathbb{R}^n$ is the set of all ordered sequences of n real numbers, but we will spend most of our time concerned with only $\mathbb{R}, \mathbb{R}^2, \mathbb{R}^3$, and occasionally, $\mathbb{R}^4$. Euclidean 4-space is handy since one might want to consider an object or shape in 3-space that is moving with respect to time, thus adding a 4th dimension.

We will write elements in $\mathbb{R}^3$ just as letters; hence, by $x \in \mathbb{R}^3$ we mean the element, $x = (x_1, x_2, x_3)$ where $x_1, x_2, x_3 \in \mathbb{R}$. The **origin** is the element, $o = (0, 0, 0)$.

Definition 75. If $x, y \in \mathbb{R}^3$, then $\overrightarrow{xy}$ is the directed line segment from x to y . We abbreviate $\overrightarrow{ox}$ by $\overrightarrow{x}$. Directed line segments are referred to as **vectors**.

Physicists and mathematicians often speak of a vector's *magnitude* and *direction*. Given a vector, $\overrightarrow{x}$, by **magnitude (or norm)** we mean the distance between the point, x , and the origin, o . By **direction** we mean the direction determined by the directed line segment $\overrightarrow{x}$ that has base at o and tip at the point, x . When we say to “sketch the vector $\overrightarrow{x}$ ” we mean to draw the directed line segment from the origin to the point, x .

We are making a distinction between *points* and *vectors*. Points are the actual elements of 3-space and vectors are directed line segments. The word **scalar** will be used to refer to real numbers (and later in your mathematical career as complex numbers or elements of any field). The word, **point** may be used to mean a real number, an element of $\mathbb{R}^2$, an element of $\mathbb{R}^3$, etc.

Having carefully made clear the distinction between *point* and *vector* you will have to work hard to keep me honest; I tend to use the two more or less interchangeably.

Definition 76. If $x, y \in \mathbb{R}^3$, with $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$, and $\alpha \in \mathbb{R}$ then:

- $x + y = (x_1 + y_1, x_2 + y_2, x_3 + y_3)$ “Addition in $\mathbb{R}^3$ ”
- $\alpha x = (\alpha x_1, \alpha x_2, \alpha x_3)$ “Scalar Multiplication in $\mathbb{R}^3$ ”

Now, to be precise, we should also define *vector addition* and *scalar multiplication for vectors*. Since the definition is identical (except for placing arrows above the x and y) we omit this. As you can see, always making a distinction between a *point* and a *vector* can be cumbersome.

Problem 350. Let $\overrightarrow{x} = \overrightarrow{(1, 2)}$ and $\overrightarrow{y} = \overrightarrow{(5, 2)}$ and sketch $\overrightarrow{x}$, $\overrightarrow{y}$, $-\overrightarrow{x}$, $2\overrightarrow{y}$.

Consider the two vectors, $\overrightarrow{xy}$ and $\overrightarrow{y} - \overrightarrow{x}$. Both vectors have the same direction and the same magnitude. They are different because the vector $\overrightarrow{y} - \overrightarrow{x}$ has its base at the origin and its tip at the point $y - x$ while $\overrightarrow{xy}$ has its base at x and its tip at y .

Problem 351. Sketch $\overrightarrow{x} + \overrightarrow{y}$, $\overrightarrow{x} - \overrightarrow{y}$ and $\overrightarrow{xy}$ where $\overrightarrow{x} = \overrightarrow{(2, 3)}$ and $\overrightarrow{y} = \overrightarrow{(4, 2)}$.

Definition 77. Let $n \in \mathbb{N}$. A function from $\mathbb{R}$ to $\mathbb{R}^n$ is called a **parametric curve**.

We will be concerned primarily with vector valued functions where the range is $\mathbb{R}^2$ or $\mathbb{R}^3$. If $n = 2$, then such functions are also called *planar curves* and if $n = 3$ they are called *space curves*. We'll put a vector symbol, $\rightarrow$, over such functions to remind us that the range is not a real number but a point in $\mathbb{R}^n$.

Problem 352. Let $\overrightarrow{l}(t) = (2t, 3t)$. Sketch $\overrightarrow{l}$ for all $t \in \mathbb{R}$. If t represents time and $\overrightarrow{l}(t)$ represents the position of a llama at time t , then how fast is the llama traveling?

Problem 353. Let $\overrightarrow{l}(t) = (1, 2) + (4, 5)t$. Sketch $\overrightarrow{l}$ for all $t \in \mathbb{R}$ and give the speed of a lemur whose position in the plane at time t seconds is given by $\overrightarrow{l}(t)$.

Problem 354. What is the distance between the point $(1, 2, 3)$ and the origin? What is the distance between $(1, 2, 3)$ and $(4, 5, 6)$?

Problem 355. Sketch $\overrightarrow{r}(t) = t(1, 2, 3) + (1 - t)(3, 4, -5)$ for all $t \in \mathbb{R}$. Compute $\overrightarrow{r}(0)$ and $\overrightarrow{r}(1)$.

Problem 356. What is the distance between $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$?

Problem 357. Let $x = (2, 5, -1)$ and $y = (1, 2, 4)$.

1. Write an equation, $\vec{l}$, for the line in $\mathbb{R}^3$ passing through x and y with $\vec{l}(0) = x$ and $\vec{l}(1) = y$.
2. Write an equation, $\vec{m}$, for the line in $\mathbb{R}^3$ passing through x and y so that $\vec{m}(0) = x$ and the speed of an object with position determined by the line is twice the speed of an object with position determined by the line in part 1 of this problem.

Problem 358. Find infinitely many parametric equations for the line passing through (a, b, c) in the direction $\overrightarrow{(x, y, z)}$.

Problem 359. Plot some points in order to graph $\vec{f}(t) = (\sin(t), \cos(t), t)$ for $t \in [0, 6\pi]$. How would you write the set representing the range of $\vec{f}$?

In Calculus I and II you studied functions from $\mathbb{R}$ to $\mathbb{R}$, for example $f(x) = x^3$, and functions from $\mathbb{R}$ to $\mathbb{R}^2$, for example $\vec{r}(t) = (\cos(t), \sin(t))$. Now we have added functions from $\mathbb{R}$ to $\mathbb{R}^3$ such as the examples in the previous few problems. We wish to add functions from $\mathbb{R}^2$ to $\mathbb{R}$ to our list next. Such functions are called *real-valued functions of several variables*.

Definition 78. A *real valued function of several variables* is a function f from $\mathbb{R}^n \rightarrow \mathbb{R}$. We will primarily consider $n = 2$ and $n = 3$ in this course.

Example 35. Functions we will encounter and their names.

1. real valued functions defined on $\mathbb{R}$
 $f : \mathbb{R} \rightarrow \mathbb{R}, f(t) = t^2 + e^t$
2. parametric curves (also called vector valued functions) defined on $\mathbb{R}$
 - (a) planar curves
 $\vec{f} : \mathbb{R} \rightarrow \mathbb{R}^2, \vec{f}(t) = (e^{2t+1}, 3t + 1)$
 - (b) space curves
 $\vec{f} : \mathbb{R} \rightarrow \mathbb{R}^3, \vec{f}(t) = (2t + 1, (3t + 1)^3, 4t + 1)$
3. real-valued functions of several variables, or *multivariate* functions
 - (a) real valued functions of two variables
 $f : \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = x^2 + \sqrt{y}$
 - (b) vector valued functions of three variables
 $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f(x, y, z) = (2xy, 3x + 4y^2)$
 - (c) vector valued functions of several variables
 $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ where $m, n \geq 2$

Problem 360. Graph the set of all points (x, y, z) in $\mathbb{R}^3$ that satisfy $x + y + z = 1$.

Problem 361. Graph the set of all points (x, y, z) in $\mathbb{R}^3$ that satisfy $z = 4$.

Definition 79. If $\vec{x}$ is a vector in $\mathbb{R}^3$, then $|\vec{x}| = \sqrt{x_1^2 + x_2^2 + x_3^2}$. This is called the **magnitude**, **length**, or **norm** of $\vec{x}$.

We defined the norm on vectors, but the same definition is valid for *points* in $\mathbb{R}^3$.

Theorem 40. Law of Cosines. Given any triangle with sides of lengths a, b , and c , and having an angle of measure α opposite the side of length a , the following equation holds: $a^2 = b^2 + c^2 - 2bc \cos(\alpha)$.

Problem 362. Sketch in $\mathbb{R}^2$ the vectors $\overrightarrow{(1,2)}$ and $\overrightarrow{(3,5)}$, and find the angle between these vectors by using the law of cosines.

Problem 363. Sketch in $\mathbb{R}^3$ the vectors $\overrightarrow{(1,2,3)}$ and $\overrightarrow{(-2,1,0)}$, and find the angle between these vectors by using the law of cosines.

Definition 80. If $\vec{x} = \overrightarrow{(x_1, x_2, x_3)}$ and $\vec{y} = \overrightarrow{(y_1, y_2, y_3)}$, then the **dot product** of $\vec{x}$ and $\vec{y}$ is defined by $\vec{x} \cdot \vec{y} = x_1y_1 + x_2y_2 + x_3y_3$

Again, we defined the dot product on vectors, but the same definition is valid for points in $\mathbb{R}^3$.

Definition 81. We say the vectors $\vec{x}$ and $\vec{y}$ are **orthogonal** if $\vec{x} \cdot \vec{y} = 0$.

Problem 364. Show that if $\vec{x}$ and $\vec{y}$ are vectors in $\mathbb{R}^3$ then

$$|\vec{x} - \vec{y}|^2 = |\vec{x}|^2 - 2\vec{x} \cdot \vec{y} + |\vec{y}|^2.$$

Problem 365. Find two vectors orthogonal to $\overrightarrow{(1,2)}$. How many are there?

Problem 366. Find three vectors orthogonal to $\overrightarrow{(1,2,3)}$. How many are there?

Problem 367. Use Theorem 40 and Problem 364 to show that if $\vec{x}, \vec{y} \in \mathbb{R}^3$ and θ is the angle between $\vec{x}$ and $\vec{y}$, then $\vec{x} \cdot \vec{y} = |\vec{x}| |\vec{y}| \cos \theta$.

Problem 368. Find two vectors orthogonal to both $\overrightarrow{(1,4,3)}$ and $\overrightarrow{(2,-3,4)}$. Sketch all four vectors.

Problem 369. Show that if two non-zero vectors $\vec{x}$ and $\vec{y}$ are orthogonal then the angle between them is 90° . Hence any two orthogonal vectors are perpendicular vectors.

Problem 370. Given the vectors $\vec{u}, \vec{v} \in \mathbb{R}^3$, find the area of the parallelogram with sides $\vec{u}$ and $\vec{v}$ and diagonals $\vec{u} + \vec{v}$ and $\vec{u} - \vec{v}$. The vertices of this parallelogram are the points: the origin, u , v , and $u+v$.

Problem 371. Assume $\vec{u}, \vec{v} \in \mathbb{R}^3$. Find a vector $\vec{x} = (x, y, z)$ so that $\vec{x} \perp \vec{u}$ and $\vec{x} \perp \vec{v}$ and $x + y + z = 1$.

Problem 372. Prove or give a counter example to each of the following where $\vec{u}, \vec{v} \in \mathbb{R}^3$ and $c \in \mathbb{R}$:

1. $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$
2. $\vec{u}(\vec{w} \cdot \vec{v}) = (\vec{u} \cdot \vec{w})(\vec{u} \cdot \vec{v})$
3. $c(\vec{u} \cdot \vec{v}) = (c\vec{u}) \cdot \vec{v} = \vec{u} \cdot (c\vec{v})$
4. $\vec{u} + (\vec{v} \cdot \vec{w}) = (\vec{u} + \vec{v}) \cdot (\vec{u} + \vec{w})$
5. $\vec{u} \cdot \vec{u} = |\vec{u}|^2$

Problem 373. Let $f(x, y) = x^2 + y^2$. Sketch the intersection of the graph of f with the planes: $z = 0$; $z = 4$; $z = 9$; $y = 0$; $y = -1$; $y = 1$; $x = 0$; $x = -1$; $x = 1$. Now sketch all of these together in one 3-D graph.

In the previous problem, the intersection of the graph with $z = 0, z = 4$, and $z = 9$ are called *level curves* because each represents the path you would take if you walked around the graph always remaining at a certain height or level. Recall the Chain Rule from Calculus I.

Theorem 41. Chain Rule. If $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions, then $(f \circ g)(t) = f(g(t))$ and $(f \circ g)'(t) = f'(g(t))g'(t)$

Problem 374. Let $g(x,y) = x^2 + y^3$ and $\vec{l}(t) = (0,1)t$. Compute $g \circ \vec{l}$. Graph g , $\vec{l}$, and $g \circ \vec{l}$.

Problem 375. Compute $(g \circ \vec{l})'(t)$ and $(g \circ \vec{l})'(2)$. What is the significance of this number with respect to your graphs from the previous problem?

Here is a reminder from Calculus II of the definition of arc length.

Definition 82. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function which is differentiable on $[a, b]$, then the **arc length of f on $[a, b]$** is $\int_a^b \sqrt{1 + (f'(x))^2} dx$. If $\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ is a vector valued function which is differentiable on $[a, b]$ so that $\vec{c}(t) = (x(t), y(t))$, then the **arc length of $\vec{c}$ on $[a, b]$** is $\int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt$.

Problem 376. A man walks along a path on the surface $f(x,y) = 4 - 2x^2 - 3y^2$ from one point on the x -axis to a second point on the x -axis, always remaining directly above the x -axis. Graph the path and write an integral expression for the distance he walked and compute the distance he walked.

Problem 377. A lady walks along the surface from the previous problem staying exactly 3 units above the xy -plane. Write an integral expression for the distance she walks if she starts and stops at $(0, \frac{1}{\sqrt{3}}, 3)$ and never retraces her steps?

Problem 378. Redo Problem 374 and Problem 375 with $\vec{l}(t) = (-1, 1)t$.

Problem 379. Find the slope of the line tangent to $f(x,y) = x^3 + 3y^2$ at $(1, 2, 13)$ that lies above the line $\vec{l}(t) = (1, 2) + (1, 1)t$.

Problem 380. Given $\vec{a} = \overrightarrow{(4, 3)}$, $\vec{b} = \overrightarrow{(1, -1)}$, and $\vec{c} = \overrightarrow{(6, -4)}$, determine the angle between $\vec{ba}$ and $\vec{bc}$.

In the next problem the notation, $|\cdot|$, is used for both the absolute value (on the left side of the equation) and the norm (on the right side of the equation). Is this bad notation? Consider the definition for the norm, that

$$|(x_1, x_2)| = \sqrt{x_1^2 + x_2^2}.$$

Suppose we take the norm of a vector in $\mathbb{R}^1$, such as (x_1) . Then,

$$|(x_1)| = \sqrt{x_1^2} = \text{the absolute value of the number } x_1.$$

Thus, the absolute value of x is the norm of x so you have been studying norms since high school (elementary school?) without knowing it!

Problem 381. Show that if $\vec{u}$ and $\vec{v}$ are vectors in $\mathbb{R}^2$ then $|\vec{u} \cdot \vec{v}| \leq |\vec{u}| |\vec{v}|$. Two approaches follow:

1. Let $\vec{u} = (u_1, u_2)$ and $\vec{v} = (v_1, v_2)$. Substitute into both sides and simplify.
2. Observe that for any constant k , $(\vec{u} - k\vec{v}) \cdot (\vec{u} - k\vec{v}) \geq 0$. Simplify this expression and use the quadratic formula.

The next result is known as the *Triangle Inequality* and it states essentially that the shortest distance between two points is the straight line. Look at a graph of $\vec{u}$, $\vec{v}$, $\vec{u+\vec{v}}$, and $\vec{u(u+\vec{v})}$. If you travel from the origin, along the vector $\vec{u}$ and then along the vector $\vec{u(u+\vec{v})}$, then you have traveled further than if you traveled along the vector $\vec{u+\vec{v}}$.

Problem 382. Triangle Inequality. Show that if each of $\vec{u}$ and $\vec{v}$ are vectors in $\mathbb{R}^2$ then $|\vec{u} + \vec{v}| \leq |\vec{u}| + |\vec{v}|$.

Chapter 9

Cross Product and Planes

“Happiness lies in the joy of achievement and the thrill of creative effort.” - Franklin D. Roosevelt

In Problem 371, you were given two vectors and asked to find a vector that was perpendicular to both. Because there are infinitely many vectors perpendicular to any two given vectors, we added another condition ($x_1 + x_2 + x_3 = 1$) so that the answer would be unique. This problem was a warm-up for the definition of the *cross product*.

Definition 83. The **cross product** of two vectors is the vector that is perpendicular to both of them and has length that is the area of the parallelogram defined by the two vectors.

If $\vec{u} = (u_1, u_2, u_3)$ and $\vec{v} = (v_1, v_2, v_3)$, then the cross product of $\vec{u}$ and $\vec{v}$ may be computed by one of two methods:

Method One.
$$\vec{u} \times \vec{v} = (u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1)$$

For the second method, you need a tool from linear algebra, determinants. If I have not done so already, ask me during class to show you how to compute the determinant of 2 by 2 and 3 by 3 matrices.

Method Two. If we define $\vec{i} = \overrightarrow{(1, 0, 0)}$, $\vec{j} = \overrightarrow{(0, 1, 0)}$, and $\vec{k} = \overrightarrow{(0, 0, 1)}$, then we may compute the cross product as:

$$\vec{u} \times \vec{v} = \det \begin{pmatrix} \vec{i} & \vec{j} & \vec{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{pmatrix}.$$

Problem 383. Prove or give a counter example for each statement, assuming $\vec{u} = (u_1, u_2, u_3)$, $\vec{v} = (v_1, v_2, v_3)$, $\vec{w} = (w_1, w_2, w_3)$ and $k \in \mathbb{R}$.

1. $k(\vec{u} \times \vec{v}) = (k\vec{u}) \times \vec{v} = \vec{u} \times (k\vec{v})$
2. $\vec{u} \times \vec{v} = -(\vec{v} \times \vec{u})$
3. $\vec{u} \times \vec{u} = \vec{u} \times \vec{0} = \vec{0} \times \vec{u} = \vec{0}$
4. $(\vec{u} \times \vec{v}) \cdot \vec{w} = \vec{u} \cdot (\vec{v} \times \vec{w})$
5. $k + (\vec{u} \times \vec{v}) = (k + \vec{u}) \times (k + \vec{v})$

$$6. \vec{u} \times (\vec{v} + \vec{w}) = (\vec{u} \times \vec{v}) + (\vec{u} \times \vec{w})$$

$$7. \vec{u} \times (\vec{v} \cdot \vec{w}) = (\vec{u} \times \vec{v}) \cdot (\vec{u} \times \vec{w})$$

$$8. \vec{u} \times (\vec{v} \times \vec{w}) = (\vec{u} \cdot \vec{w})\vec{v} - (\vec{u} \cdot \vec{v})\vec{w}$$

One might think of planes in $\mathbb{R}^3$ as analogous to lines in $\mathbb{R}^2$. This is because a line is a one-dimensional object in $\mathbb{R}^2$ – that is it has dimension one less than the dimension of the space. In $\mathbb{R}^3$ a plane is two-dimensional – it has dimension one less than the dimension of the space.

Definition 84. Given $a, b, c \in \mathbb{R}$ where a and b are not both zero, the **line** determined by a, b , and c is the collection of all points $(x, y) \in \mathbb{R}^2$ satisfying $ax + by = c$.

Given this definition we can define a *plane* in the same manner.

Definition 85. Given $a, b, c, d \in \mathbb{R}$ where a, b and c are not all zero, the **plane** determined by a, b, c and d is the collection of all points $(x, y, z) \in \mathbb{R}^3$ satisfying $ax + by + cz = d$.

If algebraically we think of a plane as all $(x, y, z) \in \mathbb{R}^3$ satisfying $ax + by + cz = d$ where not all of a, b and c are zero, then geometrically we can think of a plane as uniquely determined by a vector and a point where the plane is perpendicular to the vector and contains the point. We need both because there are infinitely many planes perpendicular to a given vector, but knowing one point in the plane uniquely determines the plane.

Problem 384. Show that $(3, -2, 5)$ is perpendicular to the plane $3x - 2y + 5z = 7$ by choosing two points, $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$, in the plane and showing that $(3, -2, 5) \cdot \overrightarrow{x - y} = 0$.

Problem 385. Show that $(a, b, c) \in \mathbb{R}^3$ is orthogonal to the plane $ax + by + cz = d$ by showing that if $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$ are two points on the plane, then $(a, b, c) \cdot \overrightarrow{x - y} = 0$.

Problem 386. Determine whether these two planes are parallel.

$$1. 2x - 3y + \frac{5}{2}z = 9$$

$$2. x - \frac{3}{2}y + \frac{5}{4}z = 12$$

Problem 387. Write in standard form ($ax + by + cz = d$) the equation of a plane $\perp$ to the first plane from previous problem and containing the point $(9, 2, 3)$.

Problem 388. Find all the planes parallel to the plane $x + y - z = 4$ and at a distance of one unit away from the plane. When does the “distance between two planes” make sense?

Problem 389. Find both angles between these two planes:

$$1. 2x - 3y + 4z = 10$$

$$2. 4x + 3y - 6z = -4$$

Problem 390. Find the equation of the plane containing $(2, 3, 4)$, $(1, 2, 3)$ and $(6, -2, 5)$.

For the next two problems, there is a formula on the web or in some book, but it’s probably wrong because of a typographical error. Find a vector $\perp$ to both planes, determine the equation of a parametric line, $\vec{l}$, passing through both planes. Find the points where $\vec{l}$ intersects each plane. Find the distance between these points.

Problem 391. Find the distance between the two planes, $x + y + z = 1$ and $x + y + z = 2$.

Problem 392. Find the distance between the two planes, $3x - 4y + 5z = 9$ and $3x - 4y + 5z = 4$.

Problem 393. Find an equation for the distance between two planes, $ax + by + cz = e$ and $ax + by + cz = f$.

Problem 394. Find the equation of the plane containing the line $\vec{l}(t) = (1 + 2t, -1 + 3t, 4 + t)$ and the point $(1, -1, 5)$.

The next problem asks for the intersection of two planes. What are all the possibilities for the intersection of any two planes? One possibility is a line. To find the equation of the line, there are a couple of ways we could think about this. First, we could find two points in the intersection and then find the equation of that line. Or we could observe that the intersection of two planes must be contained in each plane and thus is parallel to both planes. How can we find a vector that is parallel to both planes?

Problem 395. Find the intersection of the two planes $3x - 2y + 6z = 1$ and $3x - 4y + 5z = 1$.

Chapter 10

Limits and Derivatives

“Real knowledge is to know the extent of one’s ignorance.” - Confucius

For the next two definitions, suppose that x, y , and $z : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions.

Definition 86. The **limit of the vector valued function of one variable** $\vec{f}(t) = (x(t), y(t), z(t))$, as t approaches a is defined by

$$\lim_{t \rightarrow a} \vec{f}(t) = \left(\lim_{t \rightarrow a} x(t), \lim_{t \rightarrow a} y(t), \lim_{t \rightarrow a} z(t) \right)$$

as long as each of these limits exists. If any one of the limits does not exist, then the limit of $\vec{f}$ does not exist at a .

Definition 87. The **derivative of the vector valued function of one variable** $\vec{f}(t) = (x(t), y(t), z(t))$, is defined by

$$\vec{f}'(t) = (x'(t), y'(t), z'(t))$$

as long as each of the derivatives exists. If any one of the derivatives does not exist, then the derivative of $\vec{f}$ does not exist.

Problem 396. Let $\vec{f}(t) = \left(t^2 - 4, \frac{\sin(t)}{t}, \frac{4t^3}{e^t} \right)$. Compute $\lim_{t \rightarrow 0} \vec{f}(t)$.

Problem 397. Compute $\vec{f}'$ and $\vec{f}'(0)$ for $\vec{f}$ from the previous problem.

We now wish to develop the rules for limits and derivatives that parallel the rules from calculus in one dimension. Of course, we know you have not *forgotten* any of these rules, so the ones we develop should look familiar! The good news is that the really hard work in proving these was done in the first semester of calculus and thus the work here is more notational than mathematical!

Problem 398. Let $\vec{f}(t) = (t^2, t^3 - 1, \sqrt{t-1})$ and $\vec{g}(t) = (2 - t^2, t^3, \sqrt{t+1})$.

1. Compute $\lim_{t \rightarrow 2} \vec{f}(t)$.
2. Compute $\lim_{t \rightarrow 2} \vec{g}(t)$.
3. Compute $\lim_{t \rightarrow 2} \vec{f}(t) + \lim_{t \rightarrow 2} \vec{g}(t)$.

4. Compute $\lim_{t \rightarrow 2} [\vec{f}(t) + \vec{g}(t)]$.

5. What can you conjecture about $\lim_{t \rightarrow a} [\vec{f}(t) + \vec{g}(t)]$ for arbitrary choices of a , $\vec{f}$, and $\vec{g}$?

Problem 399. State 5 rules for limits of the vector valued functions, $\vec{f}, \vec{g} : \mathbb{R} \rightarrow \mathbb{R}^3$ that parallel the limit rules from Calculus I and prove one of these conjectures. You may grab a book or look on the web to remind you of the rules from Calculus I.

Problem 400. Compute $\vec{f}'(2)$ and $\vec{g}'(2)$ where $\vec{f}$ and $\vec{g}$ are from Problem 398. Compute $(\vec{f} + \vec{g})'(2)$. What can you conjecture about $(\vec{f} + \vec{g})'(t)$ for arbitrary choices of $\vec{f}$ and $\vec{g}$?

Problem 401. State 5 rules for derivatives of vector valued functions that parallel the derivative rules from Calculus I. Prove one of these conjectures. You may grab a book or look on the net to remind you of the rules from Calculus I.

You may assume that which ever one you do not prove will end up on the next test. Yes, it is a well known fact that all calculus teachers can read students' minds. How else would we always be able to schedule our tests on the same days as your physics tests?

Limits of Functions of Several Variables

Recall from Calculus I the various ways in which you computed limits. If possible, you substituted a value into the function. If not, perhaps you simplified the function via some algebra or computed a limit table or graphed the function. Or you might have applied L'Hôpital's Rule. Your instructor probably used the Squeeze Theorem to obtain the result that $\lim_{t \rightarrow 0} \frac{\sin(t)}{t} = 1$. Recall that if the left hand limit equaled the right hand limit then the limit existed.

In three dimensions, the difficulty is that there are more paths to consider than merely left and right. For the limit to exist at a point (a, b) , we need that the limit as (x, y) approaches (a, b) exists regardless of the path we take as we approach (a, b) . We could approach (a, b) along the x-axis for example, setting $y = 0$ and taking the limit as $x \rightarrow 0$. Or we could take the limit along the line, $y = x$. The limit exists if the limit as $(x, y) \rightarrow (a, b)$ along every possible path exists. We will see an example where the limit toward (a, b) exists along every straight line, but does not exist along certain non-linear paths!

Definition 88. If $(a, b) \in \mathbb{R}^2$ and $L \in \mathbb{R}$ and $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a function, then we say that

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L$$

if $f(x, y)$ approaches L as (x, y) approaches (a, b) along **every** possible path.

Problem 402. Sketch $f(x, y) = x^2 + y^2$, indicate the point $(2, 3, f(2, 3))$ and compute $\lim_{(x,y) \rightarrow (2,3)} x^2 + y^2$.

Problem 403. Use any free web-based software to graph the function from the previous example near $(0, 0)$. Print and use a high-lighter to mark the paths $x = 0$, $y = 0$, and $y = x$. (Google Chrome on a Windows box will graph functions like $z = x^2 + y^2$ just by typing it into the search bar!)

Problem 404. Convert the previous problem to polar coordinates via the substitution $x = r \cos(\theta)$ and $y = r \sin(\theta)$ and then compute the limit as $r \rightarrow 0$.

Problem 405. Let $f(x, y) = \frac{x + y^2 + 2}{x - y + 2}$.

1. Graph f using any software and state the domain.
2. Compute $\lim_{(x,y) \rightarrow (-1,2)} f(x,y)$.

Recall your Calculus I definition of continuity for functions of one variable.

Definition 89. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous at $a \in \mathbb{R}$ if $\lim_{x \rightarrow a} f(x) = f(a)$.

This definition says that for f to be continuous at a three things must happen. First, the function must be defined at a . This means that a must be in the domain of the function so that $f(a)$ is a number. Second, the limit of the function as we approach a must exist. And third, $f(a)$ must equal the limit of f at a . The same statement defines continuity for all functions.

Definition 90. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous at $a \in \mathbb{R}^n$ if $\lim_{x \rightarrow a} f(x) = f(a)$.

Problem 406. Consider $\lim_{(x,y) \rightarrow (0,0)} \frac{x^4 - y^4}{x^2 + y^2}$.

1. Compute this limit along the lines: $x = 0$, $y = 0$, $y = x$, and $y = -x$.
2. Convert to polar coordinates and check the limit.
3. Graph using any software.
4. Why isn't this function continuous at $(0,0)$?
5. How can you modify f in such a way as to make it continuous at $(0,0)$?

Problem 407. Determine whether $\lim_{(x,y) \rightarrow (0,0)} \frac{xy + y^3}{x^2 + y^2}$ exists and if so, state the limit.

Problem 408. Determine whether the function f is continuous at $(x,y) = (0,0)$ by considering the paths $y = kx^2$ for several choices of k .
$$f(x,y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} & (x,y) \neq (0,0) \\ 0 & (x,y) = (0,0) \end{cases}$$

Directional Derivatives

Suppose we have the function $f(x,y) = x^2 + y^2$ and we are sitting on that function at some point $(a,b, f(a,b))$ other than $(0,0,0)$. Then there are many directions we can walk while remaining on the surface. Depending on the direction of our path, the rate of increase of our height, or slope of our path, may vary. Some paths will move us uphill and others downhill. Suppose while we sit at the point, $(a,b, f(a,b))$, we decide to walk in a direction that will not change the y coordinate, but only changes the x coordinate. Thus, we are walking on the surface and staying within the plane, $y = b$. Walking in this way, we could go in one of two directions. Either we could go in the direction that increases x or decreases x . Let's go in the direction that increases x . Now, consider the tangent line to the curve at this point that lies in the plane, $y = b$. As we take our first step along the curve our rate of increase in height, z , will be the same as the slope of that tangent line. This slope is the *directional derivative of the function at the point (a,b) in the x direction*. If we had decided to fix $x = a$ and walk in the direction that increases y , then the slope of the line tangent to the function and in the plane $x = a$ is the *directional derivative of f at (a,b) in the y direction*.

The next definition formalizes this discussion and the problem immediately following it is an example that will make these notions of directional derivative precise!

Definition 91. If $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a function and (a, b) is in the domain of f , then the **derivative of f in the $(1, 0)$ direction at (a, b)** is the slope of the line tangent to f at the point $(a, b, f(a, b))$ and in the plane, $y = b$.

When it exists, the derivative of f with respect to x at (a, b) can be computed via the limit:

$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$$

or, since y is being held constant and x is changing, one may just compute the derivative of f as if x is the variable and y is a constant.

Notation. Suppose that $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a function as in the previous definition. There are many phrases and notations used to denote the **derivative of f in the $(1, 0)$ direction at (a, b)** . For example,

- f_1 – the **derivative of f with respect to the first variable**
- f_x – the **derivative of f with respect to x**
- $\frac{\partial f}{\partial x}$ – the **partial derivative of f with respect to x**

Similarly, f_2 , f_y , and $\frac{\partial f}{\partial y}$ would denote the same concepts where the derivative was taken in the $(0, 1)$ direction.

Problem 409. Let $f(x, y) = x^2 + y^2$ and $(a, b) = (2, -3)$.

1. Sketch f and sketch the line tangent to f at the point $(2, -3, f(2, -3))$ that is in the plane, $y = -3$.
2. Let $g(x) = f(x, -3)$ and compute $g'(2)$.
3. What part of the graph of f is the graph of g ?

Problem 410. Let $f(x, y) = \frac{y}{x}$. Compute $f_x(1, 2)$ using the limit described following Definition 91.

Problem 411. Compute f_x and f_y for each function.

1. $f(x, y) = x^3 - 4x^2$
2. $f(x, y) = e^{xy^2}$
3. $f(x, y) = \frac{x^2}{\sin(xy)}$
4. $f(x, y) = e^{x^2y} \sin(x - y)$

Definition 92. Just as in Calculus I, “second derivatives” are merely derivatives of the first derivatives. Thus $f_{xx} = (f_x)_x$. I.e. f_{xx} is the derivative of f_x with respect to x . Similarly, $f_{yy} = (f_y)_y$, $f_{xy} = (f_x)_y$ and $f_{yx} = (f_y)_x$.

Other standard notations are:

$$f_x = \frac{\partial f}{\partial x}, \quad f_y = \frac{\partial f}{\partial y}, \quad f_{xx} = \frac{\partial^2 f}{\partial x^2}, \quad f_{yy} = \frac{\partial^2 f}{\partial y^2}, \quad \text{and} \quad f_{xy} = \frac{\partial^2 f}{\partial x \partial y}$$

Theorem 42. Clairaut's Theorem. For any function, f , whose second derivatives exist, we have $f_{xy} = f_{yx}$.

Problem 412. Find f_{xx} , f_{xy} , f_{yx} , and f_{yy} for each function.

1. $f(x,y) = e^{x^2+y^2}$
2. $f(x,y) = \sin(xy + y^3)$
3. $f(x,y) = \sqrt{3x^2 - 2y^3}$
4. $f(x,y) = \cot\left(\frac{x}{y}\right)$

The study of partial differential equations is the process of finding functions that satisfy some equation that has derivatives with respect to multiple variables. For example *Laplace's Equation* is the equation $u_{xx} + u_{yy} + u_{zz} = 0$ and solutions give us information about the steady state of heat flow in a three dimensional object.

Problem 413. Which of these functions satisfy $u_{xx}(x,y) + u_{yy}(x,y) = 0$ for all $(x,y) \in \mathbb{R}^2$?

1. $u(x,y) = x^2 + y^2$
2. $u(x,y) = x^2 - y^2$
3. $u(x,y) = x^3 + 3xy^2$
4. $u(x,y) = \ln(\sqrt{x^2 + y^2})$
5. $u(x,y) = \sin(x)\cosh(y) + \cos(x)\sinh(y)$
6. $u(x,y) = e^{-x}\cos(y) - e^{-y}\cos(x)$
7. Find a solution to this equation other than the ones listed above.

In the last two problems, we studied functions of two variables and we defined derivatives in each direction, the x direction and the y direction. If f were a function of three variables, then there would be partial derivatives with respect to each of x, y , and z . Let's extend the notion of the partial derivatives with respect to x and y to functions with domain $\mathbb{R}^n$ where $n > 2$. If $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then the domain of f is $\mathbb{R}^n$ so there is a partial derivative of f with respect to the first variable, the second variable, and so on, up to the partial derivative of f with respect to the n^{th} variable. We use the notation, $f_1, f_2, f_3, \dots, f_n$ or $\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \frac{\partial f}{\partial x_3}, \dots, \frac{\partial f}{\partial x_n}$ to denote the derivative of f with respect to each variable.

Problem 414. Let $f(x_1, x_2, x_3, x_4, x_5) = x_3\sqrt{(x_1)^3 + (x_2)^2} + x_4e^{x_5x_3}$. Compute the five partial derivatives, $f_1, f_2, \dots, f_5$.

This allows us to define the derivative for functions of n variables.

Definition 93. If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a function and each partial of f exists, then the **gradient** of f is the function

$$\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

and is defined by

$$\nabla f = (f_1, f_2, f_3, \dots, f_n) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \frac{\partial f}{\partial x_3}, \dots, \frac{\partial f}{\partial x_n}\right).$$

Problem 415. For each function below, state the domain of the function, compute the gradient and state the domain of the gradient.

1. $f(x, y) = x^3y^2 - x^2y^3$
2. $g(x, y, z) = zx^3 - 3xyz + \ln(x^2yz^3)$

For any function of two variables, f , let's assume once again that we are at a point on the function, $(a, b, f(a, b))$. We know that the slope of the line tangent to f at $(a, b, f(a, b))$ and above the line $y = b$ is $f_x(a, b)$. And we know that the slope of the line tangent to f at $(a, b, f(a, b))$ and above the line $x = a$ is $f_y(a, b)$. Now consider a line in the xy -plane passing through (a, b) in some direction $(\vec{c}, \vec{d})$ that is not parallel to either the x axis or the y axis. What would the slope of the line tangent to f at $(a, b, f(a, b))$ and above this line be? From the point (a, b) there are infinitely many directions that we might travel, not just the directions parallel to the x and y axes. We can define such a direction from (a, b) by a vector, $(\vec{c}, \vec{d})$. The slopes of the lines tangent to f at $(a, b, f(a, b))$ in the direction $(\vec{c}, \vec{d})$ are called the *directional derivatives* of f at (a, b) in the direction $(\vec{c}, \vec{d})$. The partial derivatives of f with respect to x and y are your first examples of directional derivatives where the directions were $\vec{i} = (1, 0)$ and $\vec{j} = (0, 1)$. Here is the formal definition of the directional derivative.

Definition 94. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function. The **directional derivative** of f at $\vec{u}$ in the direction $\vec{v}$ is given by:

$$D_{\vec{v}} f(\vec{u}) = \lim_{h \rightarrow 0} \frac{f(\vec{u} + h\vec{v}) - f(\vec{u})}{h},$$

where $\vec{v}$ must be a unit vector.

Of course, not every limit exists, so directional derivatives may exist in some directions but not others.

Problem 416. Using the definition just stated, compute the directional derivative of $f(x, y) = 4x^2 + y$ at the point $\vec{u} = (1, 2)$ in the direction $\vec{v}$ for each $\vec{v}$ defined below.

1. $\vec{v} = (0, 1)$
2. $\vec{v} = (1, 0)$
3. $\vec{v} = (1, 1)$
4. $\vec{v} = (\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$

Important. In the previous problem, you got different answers for the directional derivatives in parts 3 and 4 even though those represent the same direction. For this reason, it is a convention to always use a *unit* vector for the direction when computing directional derivatives.

Problem 417. Use Definition 94, Definition 91, and the discussion immediately following Definition 91 to show that if $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\vec{i} = (1, 0)$, then $D_{\vec{i}} f(\vec{v}) = f_x(\vec{v})$.

Non-Definition. An analytical definition of **f is differentiable** at u is beyond the scope of this course, but a geometrical definition is not. In two dimensions (Calculus I), f was differentiable at a if there was a **tangent line** to f at $(a, f(a))$. In three dimensions (Calculus III), f is differentiable at $\vec{u}$ if there is a **tangent plane** to f at $(u, f(u))$.

Definition 95. An ε -neighborhood of $u \in \mathbb{R}^n$ is the set of all points with a distance from u of less than ε . I.e. $N_\varepsilon(u) = \{v \in \mathbb{R}^n : |u - v| < \varepsilon\}$.

How will we be able to tell when a function is “nice,” that is, when a function has a derivative?

Theorem 43. A Differentiability Theorem. If ∇f exists at u and at all points in some ε -neighborhood of u , then f is differentiable at u .

Problem 418. Is $f(x, y) = y^3(x - \frac{1}{2})^{\frac{2}{3}}$ differentiable at $(1, 2)$?

Problem 419. For each of the following problems, either prove it or give a counterexample by finding functions and variables for which it does not hold. Assume $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ are differentiable. Assume $x, y \in \mathbb{R}^2$ and $c \in \mathbb{R}$. Assume $x, y, x + y$ are in the domain of f and g .

1. $\nabla f(x + y) = \nabla f(x) + \nabla f(y)$
2. $\nabla(f + g)(x) = \nabla f(x) + \nabla g(x)$
3. $\nabla(cf)(x) = c\nabla f(x)$
4. $\nabla f(cx) = c\nabla f(x)$
5. $\nabla f(cx) = \nabla f(c) \nabla f(x)$
6. $\nabla(f \cdot g)(x) = \nabla f(x)g(x) + f(x)\nabla g(x)$
7. $\nabla(\frac{f}{g})(x) = \frac{\nabla f(x)g(x) - f(x)\nabla g(x)}{g^2}$

As in Calculus I, it is very nice to know when and where a function is continuous. The following theorem answers that question in both cases.

Theorem 44. A Continuity Theorem. From Calculus I, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at x then f is continuous at x . In Calculus III, if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable at x , then f is continuous at x .

The next theorem is a very important one. In Calculus I, we first computed derivatives using the definition and then proved rules to help us differentiate more complex functions; we do the same here. We won't prove this theorem, but it gives a very easy way to compute directional derivatives by using the dot product and the gradient of the function.

Theorem 45. Directional Derivative Theorem. $D_{\vec{v}} f(u) = \nabla f(u) \cdot \vec{v}$ for any u , $\vec{v} \in \mathbb{R}^3$ where $|\vec{v}| = 1$.

Problem 420. Redo Problem 416 using Theorem 45.

To date we have studied the derivatives of functions from $\mathbb{R}$ to $\mathbb{R}$, from $\mathbb{R}$ to $\mathbb{R}^2$, and from $\mathbb{R}^2$ to $\mathbb{R}$. Of course there is nothing special about $\mathbb{R}^2$ here. We might as well have studied $\mathbb{R}^n$ as all the derivatives would follow the same rules. Now let's consider the derivative of a function, $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Definition 96. If $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is any vector valued function of two variables defined by $f(x, y) = (u(x, y), v(x, y))$, then the derivative of f is given by

$$Df = \begin{pmatrix} u_x(x, y) & u_y(x, y) \\ v_x(x, y) & v_y(x, y) \end{pmatrix}$$

Table 10.1: Derivative Notation

Function	Derivative
$f : \mathbb{R} \rightarrow \mathbb{R}$	$f' : \mathbb{R} \rightarrow \mathbb{R}$
$f : \mathbb{R}^2 \rightarrow \mathbb{R}$	$\nabla f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$
$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$	$Df : \mathbb{R}^2 \rightarrow L(\mathbb{R}^2, \mathbb{R}^2)$

Problem 421. Compute the derivative of $f(x, y) = (x^2 \sin(xy), \frac{e^y}{\tan(x)})$.

Because of the number of different domains and ranges of functions we are studying, we have several variations of the chain rule. Before we begin, I would like to take this opportunity to apologize for the number of notations used by mathematicians, physicists, and engineers for derivatives, partial derivatives, total derivatives, gradients, Laplacians, etc. There are a number of notations and all are convenient at one time or another. I attempt to adhere for the most part to the functional notation for derivatives (f_1, f_x, f' , etc.), but Leibnitz notation, $\frac{\partial f}{\partial x}$, is a convenient notation as well. Table 10.1 illustrates my preferred notations, where $L(\mathbb{R}^2, \mathbb{R}^2)$ denotes the set of all 2x2 matrices.

The chain rule you learned in Calculus I, which is stated next, can be generalized to take the derivative of the composition of any of the functions we have been studying. That is, given any two functions with domains so that their composition actually makes sense and so that they are differentiable at the appropriate places, we can compute their derivative using the same chain rule that you learned in Calculus I with one minor change. When the domains and ranges of the functions change, the derivatives change. Thus, in the following statement, depending on the domain of f , sometimes f' means the derivative of a parametric curve, but sometimes it means the gradient of f , ∇f , and sometimes it means the matrix, Df , of derivatives of f . The same holds for the derivative of g . And finally, the symbol $\cdot$ might mean multiplication or the dot product or matrix multiplication. You'll know from context which one. The point is to realize that no matter how many different notational ways we have of writing the chain rule, it always boils down to this one.

Theorem 46. Chain Rule. If $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions, then

$$(f \circ g)' = (f' \circ g) \cdot g'$$

or with the independent variable displayed,

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x).$$

Next we state the chain rule for differentiating functions that are the composition of a function of two variables with a planar curve. Notice that this theorem is *exactly* the same as the original theorem, but restated for functions with different domains. Because $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ we replace the f' from the previous theorem with ∇f and because $\vec{g} : \mathbb{R} \rightarrow \mathbb{R}^2$ we replace the g' in the previous theorem with $\vec{g}'$. Thus the theorem still says (in English) that “the derivative of (f composed with g) is the (derivative of f) evaluated at g times (the derivative of g).”

Theorem 47. Chain Rule. If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\vec{g} : \mathbb{R} \rightarrow \mathbb{R}^2$ are both differentiable, then

$$(f \circ \vec{g})' = ((\nabla f) \circ \vec{g}) \cdot \vec{g}'.$$

Writing this with the independent variable t in place we could write:

$$(f \circ \vec{g})'(t) = (\nabla f)(\vec{g}(t)) \cdot \vec{g}'(t).$$

On the right hand side of the last line of the Chain Rule, we have the composition of ∇f with $g(t)$. Because we are multiplying vectors, “ $\cdot$ ” represents dot product and not multiplication.

Problem 422. Let $f(x, y) = x^2 - 3y^2$ and $\vec{g}(t) = (2, 3) + (4, 5)t$. Compute $(f \circ \vec{g})'$ both by direct composition and by using Theorem 47.

Problem 423. Compute $(w \circ \vec{g})'$ where $w(x, y) = e^x \sin(y) - e^y \sin(x)$ and $\vec{g}(t) = (3, 2)t$. Write a complete sentence that says what line $(w \circ \vec{g})'(-1)$ is the slope of.

Problem 424. Redo the following problems using Theorem 47 and paying special attention to the use of unit vectors.

1. Problem 374 and Problem 375.
2. Problem 378.
3. Problem 379.

Next we state the chain rule for differentiating functions that are the composition of a function of several variables with a function from the plane into the plane. Notice that this theorem is *exactly* the same as the original theorem, but restated for functions with different domains. Because $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ we replace the f' from the original theorem with ∇f and because $\vec{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ we replace the g' in the previous theorem with $D\vec{g}$. Thus the theorem still says that “the derivative of $(f \text{ composed with } g)$ is the (derivative of f) evaluated at g times (the derivative of g).” Because $D\vec{g}$ is a matrix, the right hand side of this is now a vector times a matrix.

Theorem 48. Chain Rule. If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\vec{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ are both differentiable, then

$$\nabla(f \circ \vec{g}) = ((\nabla f) \circ \vec{g}) \cdot D\vec{g}.$$

Writing this with the independent variables displayed,

$$\nabla(f(g(s, t))) = (\nabla f)(g(s, t)) \cdot (D\vec{g})(s, t).$$

Problem 425. Let $f(x, y) = 2x^2 - y^2$ and $\vec{g}(s, t) = (2s + 5t, 3st)$. Compute $\nabla(f \circ \vec{g})$ in two ways. First, compute by composing and then taking the derivative. Second, apply Theorem 48.

Problem 426. Let $w(x, y) = \ln(x + y) - \ln(x - y)$ and $g(s, t) = (te^s, -e^{st})$. Compute $\nabla(w \circ \vec{g})$ in two ways. First, compute by composing and then taking the derivative. Second, apply Theorem 48.

Problem 427. Prove or give a counterexample to the statement that $f'(\vec{l}(t)) = (f(\vec{l}(t)))'$ for all differentiable functions $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\vec{l} : \mathbb{R} \rightarrow \mathbb{R}^2$.

The next problem tells us something important. If you are sitting on some function in three-space and you are trying to decide what direction you should travel to go uphill at the steepest possible rate, then the gradient tells us this direction.

Problem 428. Use Problem 367 and Theorem 45 to show that $D_{\vec{v}} f(\vec{u})$ is largest when $\vec{v} = \nabla f(\vec{u})$ and smallest when $\vec{v} = -\nabla f(\vec{u})$.

Surfaces. Earlier we gave a list of the types of functions we have studied so far. Of course, there is only one definition for a function, so we are really talking about functions with different domains and ranges as was illustrated by the need for different chain rules for functions with different domains. In linear algebra, we see functions with domain the set of matrices (the determinant

function) and $T(f) = \int_0^1 f(x) dx$ is a function from the set of all continuous functions into the real numbers.

In earlier courses, you studied not only functions, but *relations* such as $x^2 + y^2 = 1$. What was the difference? Well, a function has a unique y for each x while a relation may have several y values for a given x value. In three dimensions a function will have a unique z for a given coordinate pair, (x, y) . When one has multiple z values for a given (x, y) value, we call it a *surface*. We have thus far studied mostly functions from $\mathbb{R}^2$ to $\mathbb{R}$ such as $f(x, y) = x^2 + y^2$ or $f(x, y) = ye^{x^2}$, but surfaces are equally important. Surfaces are to functions in three-space as relations were to functions in two-space.

Problem 429. Sketch these functions in $\mathbb{R}^3$.

1. $f(x, y) = |x| - |y|$
2. $h(x, y) = \sqrt{xy}$
3. $g(x, y) = \sin(x)$
4. $i(x, y) = 2x^2 - y^2$

Problem 430. Sketch these surfaces in $\mathbb{R}^3$.

1. $x^2 + y^2 + z^2 = 1$
2. $y^2 + z^2 = 4$
3. $x^2 - y + z^2 = 0$
4. $|y| = 1$

Surfaces may be expressed as $F(x, y, z) = k$, where $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function and k is a real number. The first example above represents the set of all points in $\mathbb{R}^3$ that satisfy $F(x, y, z) = 1$ where F is the function defined by $F(x, y, z) = x^2 + y^2 + z^2$. The mathematical convention is to rewrite these as follows:

1. $F(x, y, z) = 1$ where $F(x, y, z) = x^2 + y^2 + z^2$.
2. $F(x, y, z) = 4$ where $F(x, y, z) = y^2 + z^2$.
3. $F(x, y, z) = 0$ where $F(x, y, z) = x^2 - y + z^2$.
4. $F(x, y, z) = 1$ where $F(x, y, z) = |y|$.

Tangent Planes to Functions

Problem 431. Find the equation of the plane tangent to the function $f(x, y) = 25 - x^2 - y^2$ at the point $(3, 1, 15)$. Sketch the graph of the function and the plane.

Problem 432. Find the equation of the plane tangent to the function $f(x, y) = \sqrt{x^2 + y^2}$ at the point $(3, 4, 5)$. Sketch the graph of the function and the plane.

Tangent Planes to Surfaces

Definition 97. If $F(x, y, z) = k$ is a surface, then the **tangent plane** to F at $u = (x, y, z)$ is the plane passing through u with normal vector, $\overrightarrow{\nabla F(u)}$.

Problem 433. Find the equation of the tangent plane to the surface $x^2 - 2y^2 - 3z^2 + xyz = 4$ at the point $(3, -2, -1)$.

Problem 434. Find the equations of two lines perpendicular to the surface in the previous problem at the point $(3, -2, -1)$ on the surface.

Problem 435. Find the equations of two lines perpendicular to the surface $z + 1 = ye^y \cos(z)$ at the point $\vec{p} = (1, 0, 0)$.

Chapter 11

Optimization and Lagrange Multipliers

“The wise man doesn’t give the right answers, he poses the right questions.” - Claude Levi-Strauss

Probably the most applied concept in all of calculus is finding the maxima and minima of functions. In industry, these can be problems from engineering, such as trying to design an airplane wing that yields the maximum lift and stability while at the same time minimizing the drag coefficient. This way, we build a plane that flies easily while using less fuel. Since planes measure fuel consumption in gallons per second, a small change in wing design can result in considerable profit for the company (and a big raise for you). Have you noticed the addition of the upward turned tips at the ends of the airplane wings in recent years?

In the financial markets, mutual funds are sets of stocks. People may buy shares of the fund instead of buying shares of individual stocks. Mutual fund managers (and their clients) want to choose groups of stocks that will increase in value and make them (and their clients) rich. Thus, a fund manager wants to design a mutual fund that will maximize profits for the investors, but because investors fear volatility (large fluctuations in the value of their portfolios), the manager also wants to minimize the volatility of the mutual fund. This is an example of an optimization problem with what is called a *constraint* because you want to maximize profit but are constrained by the customers’ concerns about volatility.

Mathematicians have spent a considerable amount of time in industry working on both of these interesting problems that are representative of “real-world” applications. In mathematics, the difference between being able to understand or apply a formula to such a problem and the ability to derive or create your own formulas for the problems is the difference between working as an engineer, mutual fund manager, or biologist on a team (a wonderful job in it’s own right) and working in a think tank such as Bell Labs or Los Alamos or MSRI (the Mathematical Sciences Research Institute) where you are tackling the problems that no one can solve and creating the mathematics that will be implemented by the teams in industry.

In Calculus I you solved *optimization* or *max-min* problems by setting the derivative of a function to zero to tell you where the rate of change (or slope) of the function was zero. In Calculus III we do exactly the same thing. Except that instead of solving $f' = 0$ we are solving $\nabla f = 0$ and we are seeking a horizontal tangent *plane* instead of a horizontal tangent *line*. The procedures are very similar and both find the point in the domain of the functions where potential maxima and minima are attained.

Definition 98. A **critical point** of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is any point $x \in \mathbb{R}^2$ where $\nabla f(x)$ is zero or undefined.

Since $\nabla f = 0$ translates to $\nabla f(x,y) = (f_x(x,y), f_y(x,y)) = (0,0)$ we are seeking points (x,y) in

the plane where both $f_x(x,y) = 0$ and $f_y(x,y) = 0$, or where at least one is undefined. Review the definitions for “local minimum, local maximum, and inflection point” for functions from $\mathbb{R}$ to $\mathbb{R}$ (i.e. from Calculus I). You may look these up in a book or on the web.

Recall from Definition 95 that an ε - **neighborhood** of the point (s,t) in $\mathbb{R}^2$ is the set of all points in $\mathbb{R}^2$ that are a distance of less than ε away from (s,t) .

Definition 99. If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and x is in the domain of f , then we say $(x, f(x))$ is a **local minimum** if $f(x) < f(y)$ for every y in some ε -neighborhood of x .

Local maximum is defined similarly. A function may have infinitely many local minima and maxima.

Problem 436. Find the minimum of $f(x,y) = x^2 - 2x + y^2 - 4y + 5$ in two different ways.

1. First, set $\nabla f(x,y) = 0$ and solve for (x,y) .
2. Second, complete the square to write f as $f(x,y) = (x-a)^2 + (y-b)^2$ and graph.

Problem 437. Let $f(x,y) = 8y^3 + 12x^2 - 24xy$.

1. Find all critical points of f .
2. Sketch f using any software to verify your answers.
3. Use any software to solve the whole problem. In other words, use software to compute your partial derivatives and solve for the roots of these partial derivatives.

Recall in Calculus I that if x were a critical point for f then $(x, f(x))$ could have been a minimum, maximum, or an inflection point for f . Inflection points were critical points where the function f switched concavity (i.e. where the first derivative is zero and the second derivative switches signs). In Calculus III the analogous points are critical points that are neither maxima nor minima and we call these *saddle points*.

Definition 100. If $f : \mathbb{R} \rightarrow \mathbb{R}^2$ is differentiable at x , then we say $(x, f(x))$ is a **saddle point** if $\nabla f(x) = 0$ and no matter how small an $\varepsilon > 0$ we choose, there are points y and z in the ε -neighborhood of x so that $f(x) < f(y)$ and $f(x) > f(z)$.

This definition of a *saddle point* says that if we are at a saddle point and we decide to walk away from it, then there are paths away from the critical point along which f increases and paths away from the point along which f decreases. Given a critical point, how do we determine if it was a maximum, minimum, or saddle point? That is, how do we *classify* the critical points of f ? How did we do it in Calculus I? We restate the *Second Derivative Test* which is a slick way to classify the critical points of the single-variable, real-valued functions from Calculus I. Notice how nicely it parallels the next theorem for classifying the multi-variable, real-valued functions of Calculus III.

Theorem 49. Second Derivative Test I. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable and $f'(x) = 0$ then,

1. If $f''(x) > 0$, then $(x, f(x))$ is a minimum.
2. If $f''(x) < 0$, then $(x, f(x))$ is a maximum.
3. If f'' switches signs at x , then $(x, f(x))$ is an inflection point.

Here is a sweet theorem for classifying critical points of functions of two variables.

Theorem 50. Second Derivative Test II. If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\nabla f(u) = 0$ and $D = f_{xx}(u)f_{yy}(u) - (f_{xy}(u))^2$, then

1. $(u, f(u))$ is a local min if $D > 0$ and $f_{xx}(u) > 0$.
2. $(u, f(u))$ is a local max if $D > 0$ and $f_{xx}(u) < 0$.
3. $(u, f(u))$ is a saddle point if $D < 0$.
4. No information if $D=0$.

Example 36. Classify all critical points of a function f with gradient $\nabla f(x, y) = (xy - 2x - 3y + 6, yx - y + 4x - 4)$.

Problem 438. Compute the critical points of $f(x, y) = xy^2 - 6x^2 - 3y^2$ and classify these critical points as local maxima, local minima, or saddle points.

Problem 439. Compute and classify the critical points of $f(x, y) = xy + \frac{2}{x} + \frac{4}{y}$.

Problem 440. Compute and classify the critical points of $f(x, y) = e^{-(x^2+y^2-4y)}$.

The next problem reminds you of an important aspect of max/min problems from Calculus I. If you wanted the local maxima and minima of a function like $f(x) = x^2$ on $[-1, 3]$, then you checked not only the places where $f' = 0$ but also the values of f at the endpoints (boundary) of the interval. The same must be done for functions of several variables. When you applied this technique, you were applying the Extreme Value Theorem.

Theorem 51. Extreme Value Theorem for Functions of One Variable. If $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable function, then f has a maximum and a minimum on this interval.

The corresponding theorem for real-valued functions of two variables requires at least an intuitive idea of the notions of what it means for a set to be *closed* and *bounded*. A set S is **bounded** if there is a number M so that $|x| \leq M$ for all $x \in S$. A set S is **closed** if it contains its boundary points. Think of the open and closed intervals. An open interval does not contain its end points, the points on the boundary of the set. A closed interval does contain its boundary points. The set of all points (x, y) satisfying $x^2 + y^2 \leq 9$ is closed, while the set of all points (x, y) satisfying $x^2 + y^2 < 9$ is not closed.

Definition 101. A set S in $\mathbb{R}^2$ is **closed** if it contains all its boundary points.

Theorem 52. Extreme Value Theorem for Functions of Two Variables. If M is a closed and bounded set and $f : M \rightarrow \mathbb{R}$ is a differentiable function, then f has a maximum and a minimum on M .

Example 37. Optimize $f(x, y) = 2x^2 + y^2 + 1$ subject to $g = 0$ where $g(x, y) = x^2 + y^2 - 1$ in four ways!

Problem 441. Let $T(x, y) = 2x^2 + y^2 - y$ be the temperature at the point (x, y) on the circular disk of radius 1 centered at $(0, 0)$.

1. Find the critical points of T .
2. Find the minimum and maximum of T over the circle (perimeter), $x^2 + y^2 = 1$ by parameterizing the circle.
3. Find the minimum and maximum of T over the disk, $x^2 + y^2 \leq 1$.

For the next problem, you will need to parameterize each of the four line segments that form the line and check the maximum and minimum of the function over not only the interior of the square, but also over each of the four lines.

Problem 442. Find the maximum and minimum of $f(x,y) = 2x^2 - 3y^2 + 10$ over the square disk, $S = \{(x,y) | 0 \leq x \leq 3, 2 \leq y \leq 4\}$.

Problem 443. Find all maxima and minima of $f(x,y) = x^2 - y^2 + 4y$ over the rectangle $R = \{(x,y) | -1 \leq x \leq 1, -3 \leq y \leq 3\}$.

Problem 444. Find all maxima and minima of $f(x,y) = 2x^2 + 3y^2 + 2$ over all points on or inside the triangle with vertices, $(-2, 0)$, $(0, 2)$, and $(2, 0)$.

In Calculus I an n^{th} degree polynomial will have at most $n - 1$ critical points. What about in Calculus III? Here is a 6^{th} degree polynomial which has far more than 5 critical points. All of these may be found by hand and easily seen when graphed.

Problem 445. Find all thirteen critical points of $f(x,y) = x^3y^3 - x^3y - 3xy^3 + 3xy + 1$.

Problem 446. Let $f(x,y) = x^3 - y^3$ and $p = (2, 4)$. Find the direction in which f increases the most rapidly.

Problem 447. Ted is riding his mountain bike and is at altitude (in feet) of $A(x,y) = 5000e^{-(3x^2+y^2)/100}$. What is my slope of descent or ascent if I am riding in the direction $(-1, 1)$ starting at the point, $(10, 10, 5000e^{-4})$? In what direction should I travel to ascend the most rapidly? To descend the most rapidly?

Problem 448. Let $T(x,y,z) = \frac{10}{x^2+y^2+z^2}$ and $\vec{r}(t) = (t \cos(\pi t), t \sin(\pi t), t)$. If $T(x,y,z)$ represents the temperature in space at the point (x,y,z) and $\vec{r}(t)$ represents Ted's position at time t , then compute $(T \circ \vec{r})'(3)$ and explain what this number represents in a complete sentence. To check your answer, compute it in two ways. First compose the functions and take the derivative. Second, use the chain rule.

We have tackled optimization problems before when we found maxima and minima of functions like $f(x,y) = x^2 + x^2y + y^2 + 4$ from Problem 436. We have also sought maxima and minima of curves (or paths) on surfaces when we found the maxima and minima of $T(x,y) = 2x^2 + y^2 - y$ over the circle $x^2 + y^2 + 1$ as in Problem 441. Problem 441 is called a **constrained** optimization problem because we want a maxima or a minima of T subject to the constraint that it is above the unit circle. The method of Lagrange multipliers is a slick way to tackle constrained optimization problems.

Theorem 53. Lagrange Multipliers. Suppose that $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$. To maximize (or minimize) the function f subject to the constraint $g = 0$ we solve the two equations,

1. $\nabla f(x) = \lambda \nabla g(x)$
2. $g(x) = 0$

for x and for λ . The variable λ is called the Lagrange multiplier, and x is the point at which f is maximized (or minimized).

Example 38. Minimize the function $f(x,y) = y^2 - x^2$ over the region $g(x,y) \leq 0$ where $g(x,y) = \frac{x^2}{4} + y^2 - 1$.

Problem 449. Find any possible maxima or minima of $f(x,y) = x^2 + y^2$ subject to $xy = 3$ in three ways. First use Lagrange multipliers by putting $g(x,y) = xy - 3$ so that the constraint is $g(x,y) = 0$. Second, substitute $y = 3/x$ into the equation and solve. Third, sketch f and identify the portion of f that is above the equation, $xy = 3$.

Problem 450. Find any minima and maxima of $f(x,y) = 4x^2 + y^2 - 4xy$ subject to $x^2 + y^2 = 1$. It may be helpful to eliminate λ first.

Problem 451. Find the minimum of $f(x,y,z) = 3x + 2y + z$ subject to $9x^2 + 4y^2 - z = 0$ via Lagrange multipliers.

Chapter 12

Integration

“You cannot teach a man anything; you can only help him find it within himself.” - Galileo Galilei

Calculus III continues to parallel Calculus I and here we are at integration. Let's review integration in one variable before we tackle integration in several variables. By now you have been using anti-derivatives (and hence the Fundamental Theorem of Calculus) to compute integrals for so long that it is worth remembering the original definition for the definite integral. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function, then we define $\int_a^b f(x) dx$ as the limit of Riemann sums. If f is positive, then this is the limit of sums of areas of rectangles. Now we will define our integrals once again as limits of sums, but this time we will take limits of sums of volumes.

First, let's formally restate the definition of the definite integral from Calculus I.

Definition 102. If $[a, b]$ is a closed interval, then a partition P of $[a, b]$ is an ordered sequence $a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b$. The **norm** or **mesh** of P is $\max\{x_i - x_{i-1} : i = 1, 2, \dots, n\}$ and is denoted $\|P\|$.

Definition 103. The **integral of f over $[a, b]$** , denoted by $\int_a^b f$ is defined (assuming the limit exists) as

$$\int_a^b f = \lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(\hat{x}_i) \cdot (x_i - x_{i-1})$$

where $\hat{x}_i \in [x_{i-1}, x_i]$ and n is the number of divisions of the partition P . If the limit does not exist, we say that f is **not** integrable.

Of course, this is the limit of the sum of the areas of a collection of rectangles where the width of the rectangles tends toward zero as the mesh of the partition tends toward zero. In Calculus III, we do the same except that we must partition both the x and y -axes and we are summing volumes over rectangles rather than summing areas over intervals.

Definition 104. Given any two sets, A and B , the **Cartesian Product** of A and B is the set

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}.$$

In the case where A and B are closed intervals these are simply rectangles in the plane,

$$[a, b] \times [c, d] = \{(x, y) \mid x \in [a, b], y \in [c, d]\}$$

Definition 105. If R is the rectangle $R = [a, b] \times [c, d]$, then a **partition, U , of R** is a partition of $[a, b]$, $P = \{a = x_0 < x_1 < \dots < x_n = b\}$ along with a partition of $[c, d]$, $Q = \{c = y_0 < y_1 < y_2 < \dots < y_m = d\}$.

$\dots < y_m = d\}$. The **norm** or **mesh** of U is the largest width of any of the rectangles, $[x_{i-1}, x_i] \times [y_{j-1}, y_j]$ where $i = 1, 2, \dots, n$ and $j = 1, 2, \dots, m$. The norm of U is denoted by $\|U\|$.

Definition 106. If $f \in C_R$ (i.e. f is continuous on the rectangle R), then

$$\int_R f \, dA = \lim_{\|U\| \rightarrow 0} \sum_{i=1}^n \sum_{j=1}^m f(\hat{x}_i, \hat{y}_j) (x_i - x_{i-1})(y_j - y_{j-1})$$

where

$$(\hat{x}_i, \hat{y}_j) \in [x_{i-1}, x_i] \times [y_{j-1}, y_j].$$

Thankfully, in Calculus III we won't have to compute any such limits because there is a theorem that states:

$$\int_R f \, dA = \int_c^d \int_a^b f(x, y) \, dx \, dy$$

which gives us a straightforward way to compute integrals using anti-derivatives by using the Fundamental Theorem of Calculus from Calculus I:

Theorem 54. Fundamental Theorem of Calculus I. If f is continuous on $[a, b]$ and F is any anti-derivative of f , then $\int_a^b f = F(b) - F(a)$.

The next problem reminds us that in Calculus I we can integrate with respect to either x or y and obtain the same result. These techniques are especially handy in Calculus III.

Problem 452. Compute the area bounded by $y = x^2$, the x -axis, and $x = 2$ two ways. First, use an integral with respect to x , $\int \dots dx$, then use an integral with respect to y , $\int \dots dy$. Sketch a picture to help explain your endpoints of integration.

Most of the theorems that you proved in Calculus I have an analog in Calculus III. The following theorem provides a list of properties for double integrals that won't surprise you. We won't state them again, but they also hold for any integrable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ where $n > 2$.

When we are integrating over a two dimensional region R , we will write $\int_R f \, dA$, where the A reminds us that we are integrating over a region with *area*. This will be a double integral. When we are integrating over a three dimensional region, we will write $\int_R f \, dV$, where the V reminds us that we are integrating over a region with *volume*. This will be a triple integral.

Theorem 55. Properties of the Integral. Suppose f is integrable on a closed and bounded rectangular region R . Then

1. $\int_R [f(x, y) + g(x, y)] \, dA = \int_R f(x, y) \, dA + \int_R g(x, y) \, dA$
2. $\int_R c[f(x, y)] \, dA = c \int_R f(x, y) \, dA$ where c is a real number
3. $\int_R f(x, y) \, dA \geq \int_R g(x, y) \, dA$ if $f(x, y) \geq g(x, y)$ for all $(x, y) \in R$
4. $\int_R f(x, y) \, dA = \int_{R_1} f(x, y) \, dA + \int_{R_2} f(x, y) \, dA$, where R_1 and R_2 are rectangular regions such that R_1 and R_2 have no points in common except for points on parts of their boundaries and $R = R_1 \cup R_2$

Problem 453. Compute $\int_1^2 \int_0^3 2x + 3y \, dy \, dx$ and $\int_0^3 \int_1^2 2x + 3y \, dx \, dy$. This is called “reversing the order of integration.” Sketch the solid that you found the volume of.

Problem 454. Compute and compare these three integrals:

1. $\int_1^2 \int_{x-2}^0 2x + 3y \, dy \, dx$
2. $\int_{x-2}^0 \int_1^2 2x + 3y \, dx \, dy$
3. $\int_{-1}^0 \int_1^{y+2} 2x + 3y \, dx \, dy$

As you can see from the previous two problems, when the limits of integration contain variables care must be taken in reversing the order of integration. When the limits of integration are numbers, we can reverse the order of the integrals and obtain the same result. This is known as Fubini’s Theorem.

Theorem 56. Fubini’s Theorem. Suppose f is a continuous function of two variables, x and y , defined on the rectangle $R = [a, b] \times [c, d]$. Then

$$\int_R f(x, y) \, dA = \int_a^b \left[\int_c^d f(x, y) \, dy \right] dx = \int_c^d \left[\int_a^b f(x, y) \, dx \right] dy.$$

Problem 455. Evaluate $\int_B 4x \ln(y) z \, dV$ over the box $B = [0, 2] \times [1, 4] \times [-2, 5]$. Now change the order of integration and verify the result. How many choices are there for orders of integration?

Problem 456. Let $f(x, y) = 4 - x^2 - y^2$ over the region $R = [0, 1] \times [0, 1]$. Compute $\int_0^1 \int_0^1 f(x, y) \, dy \, dx$. Sketch the solid that this integral represents the volume of.

Problem 457. Let $f(x, y) = 4 - x^2 - y^2$. Write an integral for the volume of the solid that is bounded by f and by the planes $z = 0, x = 0, x = 2, y = 0$, and $y = 2$. Compute this integral.

Problem 458. Compute $\int_0^1 \int_0^1 \frac{y}{(xy + 1)^2} \, dx \, dy$.

Problem 459. Compute $\int_0^{\ln(3)} \int_0^1 xye^{xy^2} \, dy \, dx$.

Problem 460. Compute $\int_R \sin(x + y) \, dA$ where $R = \left[0, \frac{\pi}{2}\right] \times \left[0, \frac{\pi}{2}\right]$.

Problem 461. Compute $\int_R xy\sqrt{1 + x^2} \, dA$ where $R = [0, \sqrt{3}] \times [1, 2]$.

Thus far we have integrated primarily over domains that were rectangles, but we can also integrate over more general domains.

Problem 462. Let $f(x, y) = 4x + 2y$.

1. Sketch the region in the xy -plane bounded by $x = 2, x = 4, y = -x, y = x^2$.
2. Compute the volume of the solid below f and above this region.

Problem 463. Evaluate $\int_S xy \, dA$ where S is the region bounded by $y = x^2$ and $y = 1$.

Problem 464. Evaluate $\int_S \frac{2}{1+x^2} \, dA$ over the region S determined by the triangle with vertices $(0, 0)$, $(2, 2)$, and $(0, 2)$ in the x - y plane. Sketch the solid that you found the volume of.

Example 39. Find an integral expression for the volume of one-eighth of the sphere of radius 2 in three ways:

$$\int_{-}^{-} \int_{-}^{-} \text{----} \, dx dy = \int_{-}^{-} \int_{-}^{-} \text{----} \, dy dz = \int_{-}^{-} \int_{-}^{-} \int_{-}^{-} 1 \, dx dy dz$$

Problem 465. Fill in the blanks in order to change the order of integration.

$$\int_0^1 \int_0^{\frac{1-x}{2}} \int_0^{1-x-2y} f(x, y, z) \, dz \, dy \, dx = \int_{-}^{-} \int_{-}^{-} \int_{-}^{-} f(x, y, z) \, dx \, dz \, dy$$

Problem 466. Sketch and compute the volume of the solid bounded by $x^2 = 4y$, $z = 0$, and $5y + 9z - 45 = 0$. Write the integral both as $\int_{-}^{-} \int_{-}^{-} \text{---} \, dx \, dy$ and $\int_{-}^{-} \int_{-}^{-} \text{---} \, dy \, dx$.

Problem 467. Fill in the blanks:

$$\int_0^1 \int_{-y}^y f(x, y) \, dx \, dy = \int_{-}^{-} \int_{-}^{-} f(x, y) \, dy \, dx$$

$$\int_0^2 \int_{y^2}^{2y} f(x, y) \, dx \, dy = \int_{-}^{-} \int_{-}^{-} f(x, y) \, dy \, dx$$

Problem 468. Compute $\int_0^1 \int_{-x}^x e^{x+y} \, dy \, dx$ directly and by reversing the order of integration.

Coordinate Transformations

Loosely speaking, a *coordinate transformation* is a transformation from one coordinate system to another coordinate system and is also called *change of variables*. A cleverly chosen coordinate transformation can make a difficult integral easy. There are infinitely many, but the three most common are: the conversion from rectangular coordinates to polar coordinates (used in double integrals), from rectangular coordinates to spherical coordinates (used in triple integrals), and from rectangular coordinates to cylindrical coordinates (used in triple integrals).

In Calculus I when you did a trigonometric substitution, you did a change of variable. Written as a theorem, it would look like this.

Theorem 57. Change of Variable Theorem. If f is an integrable function over $[a, b]$ and $u : [a, b] \rightarrow [c, d]$ is a differentiable function, then

$$\int_a^b f(x) \, dx = \int_c^d f(u(t)) u'(t) \, dt$$

where we are making the substitution $x = u(t)$ and $u(c) = a$ and $u(d) = b$.

Work the following problem, using the given substitution, but keeping all the independent variables in tact, i.e. don't toss out the "t" when you replace x by $u(t)$.

Problem 469. Apply Theorem 57 to compute:

1. $\int_3^4 \frac{1}{x^2+9} dx$ using the substitution $u(t) = 3\tan(t)$
2. $\int_3^4 \frac{1}{\sqrt{x^2+9}} dx$ using the substitution $u(t) = 3\tan(t)$

For transformation of two variables the theorem is similar.

Theorem 58. Change of Variable Theorem. If f is an integrable function over the domain B and x and y are differentiable functions transforming the region B to the region B' , then

$$\int_B \int f(x,y) dx dy = \int_{B'} \int f(x(u,v), y(u,v)) J(u,v) du dv$$

where we are making the substitution $x = x(u,v)$ and $y = y(u,v)$ and

$$J(u,v) = \det \begin{pmatrix} x_u(u,v) & x_v(u,v) \\ y_u(u,v) & y_v(u,v) \end{pmatrix} = x_u(u,v)y_v(u,v) - y_u(u,v)x_v(u,v)$$

Problem 470. Integrate $\int_0^4 \int_{\frac{y}{2}}^{\frac{y}{2}+1} \frac{2x-y}{2} dx dy$. Now make the change of variable, $u = \frac{2x-y}{2}$ and $v = \frac{y}{2}$ and integrate the result.

Polar Coordinates Refresher

In polar coordinates, the point in the plane $P = (x,y)$ may be denoted by (r, θ) where r is the signed distance from $(0,0)$ to P , and θ is the angle between the vector $\vec{P}$ and the positive x -axis. We restrict θ to the interval $[0, 2\pi]$. We say r is the signed distance to allow equations such as $r = 4\sin(\theta)$ where if $\theta = \frac{11\pi}{6}$ then $r = -2$. The corresponding point would be $(-2, \frac{11\pi}{6})$ which is the same point as $(2, \frac{5\pi}{6})$. It follows that x, y, r , and θ are related by the equations:

1. $x = r\cos\theta$,
2. $y = r\sin\theta$,
3. $r = \sqrt{x^2+y^2}$, and
4. $\theta = \arctan\left(\frac{y}{x}\right)$.

Problem 471. Sketch each of the following pairs of polar functions on the same graphs.

1. $r = 5\cos(\theta)$ and $\theta = 2\pi/3$
2. $r = 2\sin(\theta)$ and $r = 2\cos(\theta)$
3. $r = 2 + 2\cos(\theta)$ and $r = 1$

Problem 472. To make a change of variables from rectangular to polar coordinates, we let x and y be the functions, $x(r, \theta) = r\cos\theta$ and $y(r, \theta) = r\sin\theta$. Show that $J(r, \theta) = r$ using Theorem 58 by filling in the missing computation:

$$J(r, \theta) = \det \begin{pmatrix} x_r & x_\theta \\ y_r & y_\theta \end{pmatrix} = \dots = r.$$

Problem 473. To make a change of variables from polar to rectangular coordinates, we let r and θ be the functions, $r(x,y) = \sqrt{x^2 + y^2}$ and $\theta(x,y) = \arctan(y/x)$. Compute and simplify $J(x,y)$.

Problem 474. Compute $\int_0^{\frac{\pi}{2}} \int_0^{\cos(\theta)} r^2 \sin(\theta) \, dr \, d\theta$.

Problem 475. Convert the previous integral to rectangular coordinates and recompute to verify your answer.

Using the previous problem and Theorem 58, we now see that when making a change of variable from rectangular to polar coordinates we have,

$$\int_B f(x,y) \, dx \, dy = \int_{B'} f(r \cos(\theta), r \sin(\theta)) \, r \, dr \, d\theta$$

where B' is the region B represented in polar coordinates. The next problem demonstrates an intuitive geometric argument supporting this result.

Problem 476. Recall that the area of the sector of a circle with radius r spanning θ radians is $A = \frac{1}{2}r^2\theta$. Let $0 < r_1 < r_2$ and $0 < \theta_1 < \theta_2 < \frac{\pi}{2}$. Sketch the region in the first quadrant bounded by the two circles $r = r_1$, $r = r_2$, and the two lines $\theta = \theta_1$, and $\theta = \theta_2$. Show that the area of the bounded region is $\left(\frac{r_1 + r_2}{2}\right)(r_2 - r_1)(\theta_2 - \theta_1)$.

Example 40. Write an integral with respect to θ that represents the area inside the graph of $r = \sin(2\theta)$ using limits and demonstrate why $\left(\frac{r_i + r_{i+1}}{2}\right)(r_{i+1} - r_i)(\theta_{i+1} - \theta_i)$ appears in the sum. Now write the same area as a double integral with respect to r and θ . Compute the volume of the cylinder of height 4 above $r = \cos(2\theta)$. Hopefully, the theme of looking at the areas or volumes of small pieces comes through.

Definition 107. Two circles are **concentric** if they share a common center, yet have distinct radii.

Definition 108. An **annulus** is a region in the plane trapped between two concentric circles.

Problem 477. Let r_1 and r_2 be real numbers with $0 < r_1 < r_2$. Let R be the portion of the annulus centered at the origin, between the two circles of radii r_1 and r_2 , and above the x -axis. Convert $\int_R e^{x^2+y^2} \, dA$ over the region R to polar coordinates and compute. Write down the endpoints of integration in rectangular coordinates.

Problem 478. Convert to polar and compute $\int_0^1 \int_0^{\sqrt{1-y^2}} \sin(x^2 + y^2) \, dx \, dy$.

Problem 479. Sketch the region bounded by $r = 2$ and $r = 2(1 + \cos(\theta))$ from $\theta = 0$ to $\theta = \pi$. Compute $\int_R y \, dA$ via polar coordinates.

Problem 480. Sketch $\theta = \frac{\pi}{6}$ and $r = 4 \sin \theta$. Compute the area of the smaller of the two regions bounded by the curves in two ways:

1. First compute $\int_R \int r \, dr \, d\theta$.
2. Check your answer by using the formula for the area inside a polar graph, $\frac{1}{2} \int_{\alpha}^{\beta} (f(\theta))^2 \, d\theta$.

Problem 481. Convert to polar and compute $\int_1^2 \int_0^{\sqrt{2x-x^2}} (x^2 + y^2)^{-\frac{1}{2}} dy dx$.

Problem 482. Find the volume of the solid under $z = 3xy$, above $z = 0$, and within $x^2 + y^2 = 2x$.

Problem 483. Find the volume of the solid in the 1st octant (i.e. where $x > 0, y > 0, z > 0$) under $z = x^2 + y^2$ and inside the surface $x^2 + y^2 = 9$.

Cylindrical and Spherical Coordinates

You already know how to represent points in the plane in two different ways, rectangular and polar coordinate systems. We wish to represent points in three-space in two new ways referred to as *cylindrical* and *spherical* coordinates.

Definition 109. The **Cylindrical Coordinate Representation** for a point $P = (x, y, z)$ is denoted by (r, θ, z) where (r, θ) are the polar coordinates of the point (x, y) in the x - y -plane and z remains unchanged (i.e. z is the height of the point above or below the x - y -plane).

Definition 110. The **Spherical Coordinate Representation** for a point $P = (x, y, z)$ is denoted by (ρ, ϕ, θ) where ρ is the distance from the point to the origin, ϕ is the angle between $\vec{P}$ and positive z -axis, and θ is the angle between the x -axis and the projection of $\vec{P}$ onto the x - y -plane. To avoid multiple representations of a single point in space we restrict $\rho \geq 0$, $0 \leq \phi \leq \pi$, and $0 \leq \theta \leq 2\pi$.

Example 41. Write an integral using spherical coordinates that represents the volume of an ice cream cone – that is, the volume inside the cone $\phi = \pi/6$ and inside the sphere $\rho = 6$. Of course, we must insert the Jacobian $\rho^2 \sin(\phi)$.

Problem 484. Sketch and write a sentence describing each of the following regions in spherical coordinates.

1. the region between $\rho = 2$ and $\rho = 3$
2. the region between $\phi = \pi/2$ and $\pi/4$
3. the region below $\rho = 10$ and above $\phi = \pi/3$

Example 42. Convert

$$\int_0^2 \int_0^{\sqrt{4-x^2}} \int_0^{\sqrt{4-x^2-y^2}} \sqrt{4-x^2-y^2} dz dy dx$$

to both spherical and cylindrical coordinates.

From Problem 473, we know that the Jacobian for polar coordinates is

$$J(r, \theta) = \det \begin{pmatrix} \cos(\theta) & -r \sin(\theta) \\ \sin(\theta) & r \cos(\theta) \end{pmatrix} = r.$$

Two of the next three problems generate the Jacobian for cylindrical and spherical coordinate systems. **Warning.** If you haven't had linear algebra yet, the next two problems require computing the determinant of 3×3 matrices. You might skip them, ask me how to compute the determinant of a 3×3 matrix, or Google how to do it.

Problem 485. To make a change of variables from rectangular to cylindrical coordinates, we let $x = x(r, \theta) = r \cos \theta$, $y = y(r, \theta) = r \sin \theta$, and $z = z$. Show that $J(r, \theta, z) = r$ using Theorem 58 by filling in the missing computation:

$$J(r, \theta, z) = \det \begin{pmatrix} x_r & x_\theta & x_z \\ y_r & y_\theta & y_z \\ z_r & z_\theta & z_z \end{pmatrix} = \cdots = r.$$

Problem 486. Consider the solid trapped inside the cylinder $x^2 + y^2 = 9$, above the function $f(x, y) = 4 - x^2 - y^2$ and below the plane $z = 10$.

1. Write a triple integral in rectangular coordinates for the volume, but don't compute.
2. Write a triple integral in cylindrical coordinates for the volume and compute it.
3. Write the volume as the volume of the cylinder minus the volume under the upside down paraboloid.

Problem 487. To make a change of variables from rectangular to spherical coordinates, we let $x(\rho, \phi, \theta) = \rho \sin \phi \cos \theta$, $y(\rho, \phi, \theta) = \rho \sin \phi \sin \theta$, and $z(\rho, \phi, \theta) = \rho \cos \phi$. Show that $J(\rho, \phi, \theta) = \rho^2 \sin(\phi)$ using Theorem 58 by filling in the missing computation:

$$J(\rho, \phi, \theta) = \det \begin{pmatrix} x_\rho & x_\phi & x_\theta \\ y_\rho & y_\phi & y_\theta \\ z_\rho & z_\phi & z_\theta \end{pmatrix} = \cdots = \rho^2 \sin(\phi).$$

Summarizing, when integrating with respect to polar, cylindrical, or spherical coordinates, we will always use the appropriate Jacobian as illustrated below.

1. Polar:

$$\int \int \dots r \, dr \, d\theta, \text{ where } x = r \cos(\theta), y = r \sin(\theta)$$

2. Cylindrical:

$$\int \int \int \dots r \, dr \, d\theta \, dz, \text{ where } x = r \cos(\theta), y = r \sin(\theta), \& z = z$$

3. Spherical:

$$\int \int \int \dots \rho^2 \sin(\phi) \, d\rho \, d\phi \, d\theta, \text{ where } x = \rho \sin \phi \cos \theta, y = \rho \sin \phi \sin \theta, \& z = \rho \cos \phi$$

The order of integration could change and you will find times when the choice of the order of integration transforms an apparently hard problem into an easy one.

Problem 488. Compute the volume of the region bounded by the two spheres $\rho = 4$ and $\rho = 6$ using spherical coordinates. Verify using the formula for the volume of a sphere, $V = \frac{4\pi}{3}r^3$.

Problem 489. Sketch and find the volume of the solid outside the cone $\phi = \frac{\pi}{3}$, inside the sphere $\rho = 6$, and above the plane $z = 0$.

Problem 490. Convert to spherical coordinates then evaluate:

$$\int_{-3}^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_{-\sqrt{9-x^2-z^2}}^{\sqrt{9-x^2-z^2}} (x^2 + y^2 + z^2)^{\frac{3}{2}} dy dz dx$$

Problem 491. Convert to spherical coordinates and compute: $\int_R e^{(x^2+y^2+z^2)^{\frac{3}{2}}} dV$ where R is the region $x^2 + y^2 + z^2 \leq 1$.

Problem 492. Find the volume of the given ellipsoids by first writing a triple integral for the volume in rectangular coordinates and then using the recommended change of variables to convert the triple integral into the new coordinate system. If all goes well, the converted integral will be easier to compute because you are transforming the ellipsoid into a sphere.

1. $\frac{x^2}{4} + \frac{y^2}{9} + \frac{z^2}{36} \leq 1$ via the change of variables $x = 2a, y = 3b, z = 6c$
2. $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} \leq 1$ via the change of variables $x = ua, y = vb, z = wc$

An Integration Application

The **mass** of an object can be found by integrating the density function, δ , over the entire object. If δ is the density function, then the **center of mass** is given by $(\vec{x}, \vec{y}, \vec{z}) = \left(\frac{M_{yz}}{m}, \frac{M_{xz}}{m}, \frac{M_{xy}}{m} \right)$ where

1. $m = \text{mass} = \int \delta(x, y, z) dx dy dz$,
2. $M_{xy} = \int \int \int z \delta(x, y, z) dx dy dz$,
3. $M_{xz} = \int \int \int y \delta(x, y, z) dx dy dz$, and
4. $M_{yz} = \int \int \int x \delta(x, y, z) dx dy dz$.

The numbers, M_{xy} , M_{xz} , and M_{yz} are called the **moments** of the object with respect to z , y , and x respectively.

Problem 493. Find the center of the mass of the solid inside $x^2 + y^2 = 4$, outside $x^2 + y^2 = 1$, below $z = 12 - x^2 - y^2$, and above $z = 0$, assuming the constant density function $\rho(x, y, z) = k$. Graph the solid and mark the center of mass to see if your answer makes sense!

Problem 494. Find the mass of the solid bounded by $z = 2 - \frac{1}{2}x^2$, $z = 0$, $y = x$ and $y = 0$, assuming the density is $\delta(x, y, z) = kz$ where $k > 0$. Challenge: Write this integral with the various orders of integration, $dz dy dx$, $dy dx dz$, $dz dx dy$, and $dx dz dy$.

Problem 495. Find the center of the mass of the solid from the previous problem.

Chapter 13

Line Integrals, Flux, Divergence, Gauss' and Green's Theorem

“The important thing is not to stop questioning.” - Albert Einstein

The phrases *scalar field* and *vector field* are new to us, but the concept is not. A scalar field is simply a function whose range consists of real numbers (*a real-valued function*) and a vector field is a function whose range consists of vectors (*a vector-valued function*).

Example 43. $f(x,y) = x^2 - 2xy$ is a scalar field. $f(x, y) = (2x, 3y)$ is a vector field on $\mathbb{R}^2$ and $f(x, y, z) = (2x - y, y\cos(z^2), x - yz)$ is a vector field on $\mathbb{R}^3$. For the point in the ocean at location (x,y,z) , the vector field $\vec{f}(x,y,z) = (p(x,y,z), q(x,y,z), r(x,y,z))$ might represent the direction of the current at this point where $p, q, r : \mathbb{R}^n \rightarrow \mathbb{R}$.

Definition 111. Let n and m be integers greater than or equal to 2. A **scalar field** on $\mathbb{R}^n$ is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. A **vector field** on $\mathbb{R}^n$ is a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$.

We now modify our definition for ∇ . If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, then we previously defined $\nabla f = (f_x, f_y) = (\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y})$. We now redefine ∇ as an *operator* that acts on the set of differentiable functions and write

$$\nabla = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right).$$

This encompasses our previous notation as we may now write

$$\nabla f = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right)(f) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (f_x, f_y).$$

We previously defined the dot product only between vectors (and points) in $\mathbb{R}^n$ and now redefine that notation as well. If $\vec{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $\vec{f}(x,y) = (p(x,y), q(x,y))$, then we will write

$$\nabla \cdot \vec{f} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right) \cdot (p(x,y), q(x,y)) = \frac{\partial p}{\partial x} + \frac{\partial q}{\partial y} = p_x(x,y) + q_y(x,y).$$

The next definition formalizes what we just wrote, stating it for $\mathbb{R}^3$.

Definition 112. If $\vec{f}(x,y,z) = (p(x,y,z), q(x,y,z), r(x,y,z)) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a differentiable vector field, then the **divergence of $\vec{f}$** is the scalar field from $\mathbb{R}^3 \rightarrow \mathbb{R}$ defined by: $\nabla \cdot \vec{f} = p_x + q_y + r_z$.

The divergence of a function at a point is the tendency for fluid to move away from or toward that point, where a positive divergence indicates fluid is moving away from the point and a negative divergence indicates that fluid is moving toward the point.

Example 44. Sketch the vector fields, $F(x, y) = (x, y)$ and $F(x, y) = (-x, y)$ and compute the divergence of each.

The next definition extends our definition of the cross product, $\times$.

Definition 113. If $\vec{f}(x, y, z) = (p(x, y, z), q(x, y, z), r(x, y, z)) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, then the **curl** of $\vec{f}$ denoted by $\nabla \times \vec{f}$ is defined by the vector valued function

$$\nabla \times \vec{f} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) \times (p, q, r) = \left(\frac{\partial r}{\partial y} - \frac{\partial q}{\partial z}, \frac{\partial p}{\partial z} - \frac{\partial r}{\partial x}, \frac{\partial q}{\partial x} - \frac{\partial p}{\partial y} \right).$$

The divergence of $\vec{f}$, denoted by $\nabla \cdot \vec{f}$, is a scalar field while the curl of $\vec{f}$, denoted by $\nabla \times \vec{f}$, is a vector field. If our vector field represents a fluid flow, then the curl of a function at a point is a the direction from that point about which the fluid rotates most rapidly. The points on the ray with base at the point and pointing in the direction of the curl tend not to move, but to spin as the fluid rotates (curls) around that ray.

Problem 496. Compute the divergence and curl of $\vec{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by

1. $\vec{f}(x, y, z) = (x^2yz, x^2 + y + \sqrt{z}, \pi x^2/yz)$
2. $\vec{f}(x, y, z) = (e^x, \ln(xyz), \sin(x^2y^z))$

Definition 114. Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, $\vec{g} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $\vec{h} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Assume f , $\vec{g}$, $\vec{h}$ are differentiable. Then

1. $(f\vec{g})(x, y, z) = f(x, y, z)\vec{g}(x, y, z)$
2. $(\vec{g} \cdot \vec{h})(x, y, z) = \vec{g}(x, y, z) \cdot \vec{h}(x, y, z)$
3. $(\vec{g} \times \vec{h})(x, y, z) = \vec{g}(x, y, z) \times \vec{h}(x, y, z)$

Problem 497. Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$, $\vec{g} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $\vec{h} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$. Prove or give a counter example:

1. $\nabla \cdot (\vec{g} + \vec{h}) = \nabla \cdot \vec{g} + \nabla \cdot \vec{h}$
2. $\nabla \cdot (f\vec{g}) = f \nabla \cdot \vec{g}$ (Note: $f \nabla \cdot \vec{g}$ means f times $\nabla \cdot \vec{g}$.)
3. $\nabla \times (\vec{g} + \vec{h}) = \nabla \times \vec{g} + \nabla \times \vec{h}$
4. $\nabla \cdot (f\vec{g}) = f \nabla \cdot \vec{g} + \vec{g} \cdot \nabla f$
5. $\nabla \times (f\vec{g}) = f \nabla \times \vec{g} + \vec{g} \times \nabla f$
6. $\nabla \cdot (\nabla \times \vec{g}) = 0$

Arc Length and Line Integrals over Scalar Fields

Problem 498. Use Definition 82 to write the integrals that would represent the arc length of $f(x) = x^2$ from $(-1, 1)$ to $(1, 1)$ and the arc length of $\vec{f}(t) = (t, t^2)$ for $-1 \leq t \leq 1$. Demonstrate your powers of integration from Calculus II by computing this integral – don't use a formula – use a trig substitution and then integration by parts.

Example 45. Compute the area of a wall that has as its base the curve $c(t) = (t, t^2)$, $0 \leq t \leq 1$ and has a height of t^3 at the point (t, t^2) .

Problem 499. For the wall example worked in class, suppose we replace the base $c(t) = (t, t^2)$ with the curve $c(t) = (u(t), v(t))$ where u and v are some real-valued functions and we replace the height $f(t) = t^3$ with some real valued function h . What would the area of the wall be now? Your answer will be an integral in terms of u , v , and h .

Problem 500. Two scalar line integrals:

1. A wall over $\vec{c}(t) = (3 \sin(t), 3 \cos(t))$ from $t = 0$ to $t = \frac{\pi}{2}$ has height $h(x, y) = x^2y$. Graph the wall and determine its area.
2. A wall over $\vec{c}(t) = (\sqrt{9-t^2}, t)$ from $t = 0$ to $t = 3$ has height $h(x, y) = x^2y$. Determine its area.

What you have just been computing are called line integrals over scalar fields.

Definition 115. The **line integral of a scalar field** $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ over the curve $\vec{c}(t) = (x(t), y(t))$ is defined by $L = \int_c g(\vec{c}(t)) |\vec{c}'(t)| dt$.

The line integral over a scalar field is computed with respect to *arc length* and we use the notation,

$$\int_c g ds \text{ to mean } \int_c g(\vec{c}(t)) \cdot |\vec{c}'(t)| dt.$$

We may use line integrals to compute the mass of a curved piece of wire in a similar manner. Suppose we have a piece of wire in the plane that is bent into the shape of a curve, $\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$. If $\delta : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\delta(x, y)$ represents the density of the wire at (x, y) , then the line integral of the density with respect to the arc length is the mass of the wire.

Problem 501. Mass is the integral of density. Find the mass of a wire in the plane with density at (x, y) of $\delta(x, y) = 2xy$ and shape $\vec{c}(t) = (3 \cos(t), 4 \sin(t))$ from $t = 0$ to $t = 2\pi$. Could such a wire exist?

Problem 502. Find the mass of the wire in three-space with density at (x, y, z) of $\delta(x, y, z) = 3z$ and in the shape $\vec{c}(t) = (2 \cos(t), 2 \sin(t), 5t)$, $t \in [0, 4\pi]$. Find its center of mass as well. Center of mass was defined at the end of the last chapter.

Definition 116. A vector field is said to be **conservative** if it is the gradient of some function. Conservative fields are also called **gradient** fields. If $\vec{f}$ is a conservative field, then any function g satisfying $\nabla g = \vec{f}$ is called a *potential* for $\vec{f}$.

Example 46. $\vec{f}(x, y) = (-2xy + 2e^y, -x^2 + 2xe^y)$ is **conservative** since $\vec{f} = \nabla g$ where g is given by $g(x, y) = -x^2y + 2xe^y$.

Problem 503. Is $\vec{f}(x, y) = (xy, x - y)$ conservative? If $\vec{f}$ is conservative, then there must be a function g so that $\nabla g = \vec{f}$. Is there a function g so that $g_x(x, y) = xy$ and $g_y(x, y) = x - y$? Why or why not?

Problem 504. Is $\vec{f}(x, y) = (x^2 + y^2, 2xy)$ conservative?

Problem 505. Show that if $\vec{f}(x, y) = (p(x, y), q(x, y))$ and $p_y(x, y) = q_x(x, y)$ then $\vec{f}$ is a gradient field.

This is part of a larger theory that we don't have time to cover, but is worth at least discussing. Problem 505 and its converse are valid in three dimensions as well. Suppose $f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a

vector field. If f is conservative, then $\nabla \times f = 0$. Conversely, if f has continuous first partial derivatives and $\nabla \times f = 0$ then f is conservative. If $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $f = (p, q)$, then Problem 505 showed that if $p_y = q_x$ then f was conservative. If $p_y = q_x$, then $q_x - p_y = 0$ which is the same statement as $\nabla \times f \cdot \vec{k} = 0$. In summary, understanding the notation $\nabla \times f$ is enough to determine when a vector field is conservative.

Problem 506. Is $\vec{f}(x, y) = \left(\frac{y}{x^2 + y^2}, \frac{-x}{x^2 + y^2} \right)$ a gradient field?

The next two examples remind us that work may be computed by integrating force.

Example 47. From physics we know that $\text{Work} = \text{Force} \cdot \text{Distance}$. So lifting a 5 lb book 2 feet requires 10 ft · lbs of work. Suppose we have a spring that has force f proportional to the square of the distance it is stretched from equilibrium so that $f(x) = kx^2$ (where $k > 0$ is the spring's coefficient). If $0 = x_0 \leq x_1 \leq \dots \leq x_N = 4$, then the work done to stretch it 4 units would be

$$\begin{aligned} W &= \lim_{N \rightarrow \infty} \sum_{i=1}^N k\hat{x}_i^2 (x_{i+1} - x_i) \text{ where } \hat{x}_i \in [x_i, x_{i+1}] \\ &= \int_0^4 kx^2 dx \text{ by the definition of the integral.} \end{aligned}$$

Thus in one dimension, work is the integral of force: $W = \int f(x) dx$.

Example 48. Suppose we have a cone full of water with radius 10 ft, height 15 ft, its point on the ground, and standing so that it holds the water. How much work is done in pumping all the water out of the cone? Hints: Force = Mass · Acceleration (due to gravity) and Mass of one slice = volume of slice · Density of H_2O . Add (integrate) the amount of work done in moving each horizontal slice of water out of the tank. The density of water is 62.5 lbs/ft³ and acceleration due to gravity is 32 ft/sec².

We now move to the study of vector (force) fields because we would like to be able to compute the total work done by an object as it moves through some field that acts on the object by either aiding or hindering its motion. As examples, think of a metal ball passing through a magnetic field, an electron passing through an electric field, a submarine passing through a fluid field, or a person passing through a gravitational field. Suppose we have an object passing along a curve through a vector field. Let $\vec{f}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be our (differentiable) vector field and $c: \mathbb{R} \rightarrow \mathbb{R}^2$ be our (differentiable) planar curve or path of the object. Therefore $\vec{f}$ is of the form, $\vec{f}(x, y) = (p(x, y), q(x, y))$ for some $p, q: \mathbb{R}^2 \rightarrow \mathbb{R}$ and c is of the form, $c(t) = (a(t), b(t))$ for some $a, b: \mathbb{R} \rightarrow \mathbb{R}$. Thus we may compute the work done as the particle moves through the vector field by integrating over the field evaluated at the particle, dotted with the speed of the particle. This gives the component of the force that is acting in the direction of travel of the particle.

$$\begin{aligned} W &= \int \vec{f}(\vec{c}(t)) \cdot \vec{c}'(t) dt \\ &= \int (p(a(t), b(t)), q(a(t), b(t))) \cdot (a'(t), b'(t)) dt \\ &= \int p(a(t), b(t)) a'(t) + q(a(t), b(t)) b'(t) dt \end{aligned}$$

There are two additional ways in which vector line integrals are often written. First, the independent variable t is often omitted:

$$W = \int_{\vec{c}} \vec{f}(\vec{c}(t)) \cdot \vec{c}'(t) dt = \int_{\vec{c}} \vec{f}(\vec{c}) \cdot d\vec{c}.$$

Second, if we $f(x, y) = (p(x, y), q(x, y))$ and $\vec{c}(t) = (x(t), y(t))$ then

$$\begin{aligned} W &= \int f(\vec{c}) \cdot d\vec{c} \\ &= \int (p(x, y), q(x, y)) \cdot (x'(t), y'(t)) dt \\ &= \int p(x(t), y(t)) x'(t) + q(x(t), y(t)) y'(t) dt \\ &= \int p dx + q dy \end{aligned}$$

We now have many different types of integrals to compute. Here is a table of notations to help you determine which integral we are computing. Some of these notations we have already seen, some are coming soon.

- dx or dt means the usual, for example:

$$\int_1^2 x^2 dx = \int_1^2 t^2 dt$$

- ds means we integrate with respect to arc length, for example:

$$\int_{\vec{c}} f ds = \int_{\vec{c}} f(\vec{c}(t)) |\vec{c}'(t)| dt$$

- $d\vec{c}$ means a line integral over a vector field along some curve, for example:

$$\int_{\vec{c}} \vec{f} \cdot d\vec{c} = \int_{\vec{c}} \vec{f}(\vec{c}(t)) \cdot \frac{\vec{c}'(t)}{|\vec{c}'(t)|} ds = \int_{\vec{c}} \vec{f}(\vec{c}(t)) \cdot \frac{\vec{c}'(t)}{|\vec{c}'(t)|} |\vec{c}'(t)| dt = \int_{\vec{c}} \vec{f}(\vec{c}(t)) \cdot \vec{c}'(t) dt$$

- dA means we integrate over a two dimensional domain, for example the double integral:

$$\int_D f dA = \iint f(x, y) dx dy$$

- dV means we integrate over a three dimensional domain, for example the triple integral:

$$\int_D f dV = \iiint f(x, y, z) dx dy dz$$

Problem 507. Suppose we have a particle traveling through the force field, $\vec{f}(x, y) = (xy, 2x - y)$.

1. Compute the work done as the particle travels through the force field f along the curve $\vec{c}(t) = (t, t^2)$ from $t = 0$ to $t = 1$.
2. Compute the work done as the particle travels through the force field f along the curve $\vec{c}(t) = (2t, 4t^2)$ from $t = 0$ to $t = \frac{1}{2}$.

What you have just computed is called a *line integral over a vector field*.

Definition 117. The **line integral of the vector field** $\vec{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $\vec{f}(x, y) = (p(x, y), q(x, y))$ over the curve $\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ given by $\vec{c}(t) = (x(t), y(t))$ is defined by

$$L = \int \vec{f}(\vec{c}) \cdot d\vec{c} = \int (p(x(t), y(t)), q(x(t), y(t))) \cdot (x'(t), y'(t)) dt$$

Problem 508. Compute the line integral of $\vec{f}(x, y) = (4x + y, x + 2y)$ over $\vec{c}$ where $\vec{c}$ is the curve bordering the rectangle with corners at $(0, 0)$, $(1, 0)$, $(1, 2)$, $(0, 2)$. Start your path at the origin and work counter-clockwise around the rectangle. You will need to compute and sum four integrals along four parameterized lines.

You have been using the Fundamental Theorem of Calculus for three semesters now. This powerful result in one dimension is all that is needed to prove the version we need for three dimensions.

Theorem 59. The Fundamental Theorem of Calculus. If F is any anti-derivative of f , then
$$\int_a^b f(x) dx = F(b) - F(a).$$

Theorem 60. The Fundamental Theorem of Calculus for Line Integrals. If $\vec{f}$ is a gradient field and g is any anti-derivative of $\vec{f}$ (i.e. $\nabla g = \vec{f}$), then

$$\int_{\vec{c}} \vec{f}(\vec{c}) \cdot d\vec{c} = g(\vec{c}(b)) - g(\vec{c}(a)).$$

Problem 509. Prove Theorem 60.

Problem 510. Compute the line integral from Problem 508 using Theorem 60.

The next two theorems are important results associated with line integrals. We don't assert all the necessary hypothesis – curves are smooth, functions are assumed to be integrable, etc.

Theorem 61 states that if we reverse the path along which we compute a line integral, then we change the sign of the result. Thinking of the line integral as work, this makes sense, because the work done by traveling one direction along a path should be the opposite of the work done traveling the other way. When we have a curve $\vec{c}$ and we write $-\vec{c}$ we mean the same set of points in the plane, but we simply reverse the direction. If we have the curve $\vec{c}(t) = (x(t), y(t))$ from $t = a$ to $t = b$, then $-\vec{c}$ is simply the same curve where t goes from $t = b$ to $t = a$.

Theorem 61. Path Reversal Theorem for Line Integrals. If $\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ is a parametric curve and $\vec{f}$ is a vector field, then

$$\int_c \vec{f}(\vec{c}) \cdot d\vec{c} = - \int_{-c} \vec{f}(\vec{c}) \cdot d\vec{c}.$$

Problem 511. Show that $\int_{\vec{c}} \vec{f}(\vec{c}) \cdot d\vec{c} = - \int_{-\vec{c}} \vec{f}(\vec{c}) \cdot d\vec{c}$ where $\vec{f}(x, y) = (xy, y - x)$, $\vec{c}(t) = (t^2, t)$, $t \in [0, 2]$. Since c is the path from $(0, 0)$ to $(4, 2)$, then $-c$ is the same path but from $(4, 2)$ to $(0, 0)$.

Theorem 62 states that line integrals over conservative fields are independent of path. Suppose you and I start at the same point at the bottom of a mountain and we walk to the top following different paths. Did we do the same amount of work? Yes. Because gravity is a conservative (gradient) field it does not matter what path we follow as long as we both start at the same place and end at the same place. By the same logic, if we start at a point on the mountain, walk around, and return to the same spot, then the total work is zero.

Theorem 62. Independence of Path Theorem for Line Integrals. If $\vec{c}_1$ and $\vec{c}_2 : \mathbb{R} \rightarrow \mathbb{R}^2$ are two (distinct) paths beginning at the point (x_1, y_1) and ending at the point (x_2, y_2) in the plane and $\vec{f}$ is a gradient field, then

$$\int_{c_1} \vec{f}(\vec{c}) \cdot d\vec{c} = \int_{c_2} \vec{f}(\vec{c}) \cdot d\vec{c}$$

Problem 512. Let f be the vector field $\vec{f}(x, y) = (3x^2y + x, x^3)$ and c be the planar curve $\vec{c}(t) = (\cos(t), \sin(t))$ from $t = 0$ to $t = \pi$.

1. Write out and simplify, but don't compute, the line integral of f over $\vec{c}$.
2. Compute this line integral by choosing the simpler path $\vec{p}(t) = (-t, 0)$ from $t = -1$ to $t = 1$ and applying Theorem 62.
3. Recompute this line integral by finding a function g so that $\nabla g = (3x^2y + x, x^3)$ and applying Theorem 60.

Problem 513. Let $\vec{f}(x, y) = (x - y, x + y)$.

1. Compute $\int_{\vec{c}_1} \vec{f}(\vec{c}) d\vec{c}$ where $\vec{c}_1(t) = (t, t^2)$, $t \in [0, 1]$.
2. Compute $\int_{\vec{c}_2} \vec{f}(\vec{c}) d\vec{c}$ where $\vec{c}_2(t) = (\sin(t), \sin^2(t))$ $t \in [2\pi, \frac{5\pi}{2}]$.
3. Explain your answer.

When we compute line integrals, we often integrate along a curve that is the boundary of some region. Here are a few of the buzz words about curves that we will use.

Definition 118. Let $\vec{c} : [a, b] \rightarrow \mathbb{R}^2$. We say that $\vec{c}$ is a **simple closed curve** if $\vec{c}$ starts and ends at the same point (i.e. $\vec{c}(a) = \vec{c}(b)$) and never crosses itself.

When we are integrating around a simple closed curve $\vec{c}$ with respect to the arc length of the curve, we will use the notation $\oint_{\vec{c}} \cdots ds$. The circle indicates that the curve is closed, starting and ending at the same point in the plane.

Definition 119. Let $\vec{c} : [a, b] \rightarrow \mathbb{R}^2$ be a simple closed curve on $[a, b]$. We say that $\vec{c}$ is **positively oriented** if as t increases from a to b and we traverse the curve $\vec{c}$ then the enclosed region is on our left.

As we will show in a forthcoming lecture, Gauss' Divergence Theorem is equivalent to Green's Theorem. They are simply the same theorem stated in two different ways. Gauss' Divergence Theorem says that if we have a certain simple closed curve representing the boundary of a region over which we have a vector field (a flow), then the flow across boundary (the flux) must equal the integral of the divergence of the fluid (or the electricity or whatever) over the region enclosed by the boundary.

For both Green's and Gauss' Theorems, when integrating along the boundary of the region the simple closed curve must be positively oriented. That is, you must integrate in a counter-clockwise direction so that the region is on your left as you travel along the parametric curve.

Theorem 63. Gauss' Divergence Theorem for the Plane. If F is a vector field and $\vec{c}$ is a simple closed curve and $\vec{n}$ is the unit normal to $\vec{c}$, then $\oint_{\vec{c}} \vec{F} \cdot \vec{n} ds = \int_D \nabla \cdot \vec{F} dA$

Example 49. Verify Gauss' Divergence theorem for $\vec{f}(x, y) = (-2x, -2y)$ over the triangular domain with vertices $(0, 0)$, $(0, 1)$ and $(1, 1)$.

Problem 514. Verify Gauss' Divergence Theorem by computing both integrals (the flux integral and the divergence integral) with $\vec{f}(x, y) = -4(x, y)$ and $\vec{c}(t) = (\cos(t), \sin(t))$ assuming $t \in [0, 2\pi]$.

Problem 515. Verify Gauss' Divergence Theorem for $\vec{f}(x,y) = (xy, 2x - y)$ over the region D in the first quadrant bounded by $y = 0$, $x = 0$ and the line $y = 1 - x$.

Problem 516. Verify Gauss' Divergence Theorem for the flow $f(x,y) = (0,y)$ over the rectangular region, $\{(x,y) : 0 \leq x \leq 2, 0 \leq y \leq 3\}$.

Theorem 64. Green's Theorem If $\vec{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $\vec{f}(x, y) = (p(x, y), q(x, y))$, is a vector field and $\vec{c} : \mathbb{R} \rightarrow \mathbb{R}^2$ is a positively oriented simple closed curve so that $\vec{c}(t) = (x(t), y(t))$ and D is the region enclosed by $\vec{c}$, then

$$\oint_{\vec{c}} \vec{f}(\vec{c}) \cdot d\vec{c} = \int_D q_x(x,y) - p_y(x,y) dA$$

Example 50. Verify Green's theorem for $\vec{f}(x,y) = (4y, 3x + y)$ over the region bounded by the x -axis and $y = \cos(x)$. Compute both sides of Green's theorem for $\vec{f}(x,y) = (x - y, x + y)$ over the circle $\vec{c}(t) = (2 \sin(t), 2 \cos(t))$ and explain why they are not equal.

Problem 517. Verify Green's Theorem for the flow $\vec{f}(x,y) = (-y^2, xy)$ over the region bounded by $x = 0, y = 0, x = 3$, and $y = 3$.

Problem 518. Verify Green's Theorem for the vector field $\vec{f}(x,y) = (-y, x)$ over the region bounded by the curve $\frac{x^2}{4} + \frac{y^2}{9} = 1$.

Problem 519. Verify Green's Theorem for the vector field $\vec{f}(x,y) = (xy, 3x)$ over the region formed by the three points, $(-1, 0)$, $(1, 0)$, and $(0, 4)$.

Theorem 65. An Area Theorem. If we have a simple, closed, positively oriented parametric curve $\vec{c}(t) = (x(t), y(t))$, then the area enclosed by the curve may be computed by: $A = \frac{1}{2} \int_a^b x dy - y dx$.

Problem 520. Find the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ via the formula $A = \frac{1}{2} \int x dy - y dx$ and the parametrization $\vec{c}(t) = (a \cos(t), b \sin(t))$, $t \in [0, 2\pi]$.

Gauss' Divergence Theorem is valid in higher dimensions as well, although it is often the case that integrating over certain parts of the boundary is challenging.

Theorem 66. Gauss' Divergence Theorem for Three Dimensions. If S is a solid in three dimensional space, we write

$$\int_{\delta S} f \cdot n dS = \int_S \nabla \cdot f dV$$

where δS denotes the boundary of S , n denotes the unit outward normal and dV indicates that we are integrating over the entire volume of the solid.

Problem 521. Verify Gauss' Divergence Theorem for the vector field $\vec{f}(x,y,z) = (x^2y, 2xz, yz^3)$ over the three dimensional box, $0 \leq x \leq 1, 0 \leq y \leq 2$, and $0 \leq z \leq 3$.

Problem 522. Let S be the cylinder of radius 2 with base centered at the origin and height 3. Let $\vec{f}(x,y,z) = (x^3 + \tan(yz), y^3 - e^{xz}, 3z + x^3)$. Use the divergence theorem to compute the flux across the side of the cylinder.

Problem 523. Let S be the solid bounded by $2x + 2y + z = 6$ in the first octant. Let $\vec{f}(x,y,z) = (x, y^2, z)$. Sketch the solid and verify Gauss' Divergence Theorem.

Congratulations, for some of you this note constitutes having independently worked through three semesters of Calculus, which is quite an accomplishment.

Chapter 14

Practice Problems

14.1 Functions, Limits, Velocity, and Tangents Drill

Graphing

1. Graph $f(x) = x^2 - 4$, listing x - and y -intercepts.
2. Graph $g(x) = \frac{x-1}{x+2}$, listing both vertical and horizontal asymptotes.
3. Graph $h(x) = \sin(x)$, listing the domain and range.
4. Graph $r(x) = \sqrt{3x-2}$, listing the domain and x -intercept.
5. Graph $f(x) = \begin{cases} x-5 & \text{if } x > 1 \\ x^2+1 & \text{if } x \leq 1 \end{cases}$

Evaluate the following limits or state why they do not exist.

1. $\lim_{x \rightarrow 2} (x^3 + 2x^2 - 5x + 7)$
2. $\lim_{x \rightarrow \frac{\pi}{2}} \cos(2x)$
3. $\lim_{x \rightarrow -2} \frac{x+2}{x^2-4}$ and $\lim_{x \rightarrow -2} \frac{x^2-4}{x+2}$
4. $\lim_{x \rightarrow 1} f(x)$, where $f(x) = \begin{cases} x-5 & \text{if } x > 1 \\ x^2+1 & \text{if } x \leq 1 \end{cases}$
5. $\lim_{x \rightarrow 3} \sqrt{x-3}$
6. $\lim_{h \rightarrow 2} \frac{3xh+h}{h}$
7. $\lim_{x \rightarrow -3} \frac{x^2+5x+6}{x^2-x-12}$
8. $\lim_{t \rightarrow 1} \frac{t^3-1}{t^2-1}$
9. $\lim_{x \rightarrow -1} \frac{1+2x}{1+x}$

10. $\lim_{x \rightarrow 0} \frac{\sqrt{3-x} - \sqrt{3}}{x}$
11. $\lim_{x \rightarrow \frac{\pi}{8}^+} \cos(2x)$, $\lim_{x \rightarrow \frac{\pi}{8}^-} \cos(2x)$, and $\lim_{x \rightarrow \frac{\pi}{8}} \cos(2x)$
12. $\lim_{x \rightarrow 1^-} f(x)$, $\lim_{x \rightarrow 1^+} f(x)$, and $\lim_{x \rightarrow 1} f(x)$
 where $f(x) = \begin{cases} x+3 & \text{if } x > 1 \\ 9x^2 - 5 & \text{if } x \leq 1 \end{cases}$
13. Let $f(x) = 1 - 2x^2$. Compute $\lim_{t \rightarrow x} \frac{f(t) - f(x)}{t - x}$.

List the interval(s) on which each of the following is continuous.

- $f(x) = \frac{3x+2}{6x^2-x-1}$
- $G(x) = \sqrt{x+5}$
- $f(x) = x^4 + 2x^3 - 6x + 1$
- $h(x) = \sqrt{2-4x}$
- $g(t) = \frac{t}{t-2}$
- $f(\beta) = 3\tan(\beta)$
- $f(x) = \begin{cases} x^2 + 2 & \text{if } x \geq 1 \\ 5x - 2 & \text{if } x < 1 \end{cases}$
- $f(x) = \begin{cases} 3 - x^2 & \text{if } x \geq \pi \\ \cos(2x) & \text{if } x < \pi \end{cases}$

Velocity

- Given position function $f(t) = 3t^2 + t - 1$, where t is in seconds and $f(t)$ is in feet, find the average velocity between 1 and 4 seconds. Find the average velocity between p and q seconds.
- Suppose that $y(t) = -5t^2 + 100t + 5$ gives the height t (in hundreds of feet) of a rocket fired straight up as a function of time t (in seconds). Determine the instantaneous velocity at times $t = 2$ and at $t = 4$.

Secant and Tangent Lines

- Find the equation of the secant line of $g(x) = 1 + \sin(x)$ containing the points $(0, 1)$ and $(\frac{\pi}{2}, 2)$.
- Find the slope of the tangent line to $f(t) = 5t - 2t^2$ at $t = -1$.
- Find the equation of the tangent line to $F(x) = \frac{x}{x+1}$ at $x = -3$. Graph both F and the tangent line.
- Find the equation of the tangent line to $H(x) = \sqrt{1+x}$ at $x = 8$ and sketch the graph of both H and the tangent line.

Functions, Limits, and Velocity Solutions

Graphing

1. parabola with vertex at $(0, -4)$
2. rational function with horizontal asymptote $y = 1$ and vertical asymptote $x = -2$
3. be sure you can graph all your trig functions
4. root functions look like half a parabola sideways
5. half parabola and half straight line

Evaluate the following limits or state why they do not exist.

1. 13
2. -1
3. $-\frac{1}{4}$ and -4
4. DNE since $\text{LHL} = 2$ doesn't equal $\text{RHL} = -4$
5. DNE since LHL DNE (note that $\text{RHL} = 0$)
6. $3x + 1$
7. $\frac{1}{7}$
8. $\frac{3}{2}$
9. limit does not exist; $x = -1$ is a horizontal asymptote, so neither left nor right limits exist
10. $\frac{-1}{2\sqrt{3}}$
11. $\sqrt{2}/2$ for all three answers
12. 4 for all three answers
13. $-4x$

List the interval(s) on which each of the following is continuous.

1. $(-\infty, -\frac{1}{3}), (-\frac{1}{3}, \frac{1}{2}), (\frac{1}{2}, \infty)$
2. $[-5, \infty)$
3. $(-\infty, \infty)$
4. $(-\infty, \frac{1}{2}]$
5. $(-\infty, 2), (2, \infty)$
6. continuous everywhere except $\beta = \frac{(2k-1)\pi}{2}$ where k is an integer
7. $(-\infty, \infty)$

8. all x except $x = \pi$

Velocity

1. 16 ft/sec, $3(p + q) + 1$ ft/sec
2. 8000 and 6000 ft/sec

Secant and Tangent Lines

1. $y = \frac{2}{\pi}x + 1$
2. $y = 9x + 2$
3. $y = \frac{1}{4}x + \frac{9}{4}$
4. $y = \frac{1}{6}x + \frac{5}{3}$

14.2 Differentiation Drill

1. Graph these functions and their derivatives.

(a) $F(x) = -x^2 + 5$

(b) $h(x) = x^3 - 9x$

(c) $f(x) = \cos(2x)$

(d) $g(t) = |2 + 3t|$

2. Compute the derivative of each function.

(a) $g(t) = \sqrt{3t} + \frac{3}{t} - \frac{5}{t^3}$

(b) $z(t) = (x^2 + 1)(x^3 + 2x)$

(c) $y = \frac{2x^3 + x}{2 - x^2}$

(d) $g(t) = (2t^2 - 5t^8 + 4t)^{12}$

(e) $F(z) = \sqrt[3]{2z + 7z^3}$

3. Compute and simplify the derivatives of these exponential and logarithmic functions.

(a) $a(x) = 12 - e^{x^2}$

(b) $b(x) = \ln(x^2)$

(c) $c(x) = e^{\sqrt{x^2-1}}$

(d) $d(x) = x \ln(x^3)$

(e) $g(t) = \frac{t + e^t}{3t^2 - 10t + 5}$

(f) $f(t) = \frac{1+t}{e^t}$

(g) $n(x) = \ln(xe^{x^2})$

4. Compute and simplify the derivatives of these trigonometric functions.

(a) $C(\beta) = 3\beta \cos(5\beta)$

(b) $H(x) = (5x^2 + x)(4x^4 + \tan(x))$

(c) $o(x) = \csc(x) \cot(x)$

(d) $p(x) = \sin(\cos(x))$

(e) $r(x) = (\sin(x)/x)^3$

(f) $i(x) = \sin^3(x)/x^3$

5. Compute and simplify the derivatives of these mixed functions.

(a) $g(t) = t \sin(t)e^t$

(b) $h(x) = \tan(x^3 - 3x)$

(c) $k(t) = \cos^{-1}(x^2 + 3x)$

(d) $m(x) = \sin^{-1}(5x)$

- (e) $q(t) = e^{\sec(t^2-2t)}$
- (f) $s(x) = 3^x + x^3$
- (g) $t(y) = 2^{y^2-3y}$
- (h) $u(x) = x3^{\sin(x)}$
- (i) $G(x) = (x+1)^{\sin(x)}$

6. Each of the hyperbolic trigonometric functions is defined – verify that the stated derivatives are correct.

- (a) $\sinh(x) = \frac{e^x - e^{-x}}{2}$ Show: $(\sinh(x))' = \cosh(x)$
- (b) $\cosh(x) = \frac{e^x + e^{-x}}{2}$ Show: $(\cosh(x))' = \sinh(x)$
- (c) $\tanh(x) = \frac{\sinh(x)}{\cosh(x)}$ Show: $(\tanh(x))' = \operatorname{sech}^2(x)$
- (d) $\operatorname{csch}(x) = \frac{1}{\sinh(x)}$ Show: $(\operatorname{csch}(x))' = -\operatorname{csch}(x)\coth(x)$
- (e) $\operatorname{sech}(x) = \frac{1}{\cosh(x)}$ Show: $(\operatorname{sech}(x))' = -\operatorname{sech}(x)\tanh(x)$
- (f) $\coth(x) = \frac{1}{\tanh(x)}$ Show: $(\coth(x))' = -\operatorname{csch}^2(x)$

7. Show that $\sinh^{-1}(x) = \ln(x + \sqrt{x^2 + 1})$.

8. Show that $(\sinh^{-1}(x))' = \frac{1}{\sqrt{x^2 + 1}}$.

9. Compute the slope of the tangent line to $f(x) = \frac{2x}{2^x}$ when $x = 0$.

10. Compute the equation of the tangent line to $g(x) = x \tan(x)$ when $x = \frac{\pi}{3}$.

Differentiation Drill Solutions

1. Graph these functions and their derivatives.

- (a) $F'(x) = -2x$
- (b) $h'(x) = 3x^2 - 9$
- (c) $f'(x) = -2\sin(2x)$
- (d) $g'(t) = 3(2 + 3t)/|2 + 3t|$

2. Compute the derivative of each function.

- (a) $g'(t) = \frac{\sqrt{3}}{2\sqrt{t}} - \frac{3}{t^2} + \frac{15}{t^4}$

- (b) $z'(t) = 5x^4 + 9x^2 + 2$
 (c) $y' = -\frac{2x^4 - 13x^2 - 2}{(x^2 - 2)^2}$
 (d) $g'(t) = 48t^{11}(5t^7 - 2t - 4)^{11}(10t^7 - t - 1)$
 (e) $F'(z) = \frac{2 + 21z^2}{3\sqrt[3]{(2z + 7z^3)^2}}$

3. Compute and simplify the derivatives of these exponential and logarithmic functions.

- (a) $a'(x) = -2xe^{x^2}$
 (b) $b'(x) = 2/x$
 (c) $c'(x) = \frac{xe^{\sqrt{x^2-1}}}{\sqrt{x^2-1}}$
 (d) $d'(x) = 3 + \ln(x^3)$
 (e) $g'(t) = \frac{(3t^2 - 16t + 15)e^t - 3t^2 + 5}{(3t^2 - 10t + 5)^2}$
 (f) $f'(t) = \frac{-t}{e^t}$
 (g) $n'(x) = 2x + 1/x$

4. Compute and simplify the derivatives of these trigonometric functions.

- (a) $C'(\beta) = 3(\cos(5\beta) - 5\beta\sin(5\beta))$
 (b) $H'(x) = (10x + 1)(4x^4 + \tan(x)) + (5x^2 + x)(16x^3 + \sec^2(x))$
 (c) $o'(x) = -\frac{\cos^2(x) + 1}{\sin^3(x)}$
 (d) $p'(x) = -\sin(x)\cos(\cos(x))$
 (e) $r'(x) = 3\sin^2(x)\frac{x\cos(x) - \sin(x)}{x^4}$
 (f) $i'(x) =$ same answer as last one!

5. Compute and simplify the derivatives of these mixed functions.

- (a) $g'(t) = te^t \cos(t) + (t + 1)e^t \sin(t)$
 (b) $h'(x) = (3x^2 - 3)\sec^2(x^3 - 3x)$
 (c) $k'(t) = \frac{2x + 3}{\sqrt{1 - (x^2 + 3x)^2}}$
 (d) $m'(x) = \frac{5}{\sqrt{1 - 25x^2}}$
 (e) $q'(t) = (2t - 2)e^{\sec(t^2 - 2t)} \sec(t^2 - 2t) \tan(t^2 - 2t)$
 (f) $s'(x) = 3x^2 + 3^x \ln(3)$
 (g) $t'(y) = \ln(2)(2y - 3)2^{y^2 - 3y}$
 (h) $u'(x) = 3^{\sin(x)} + \ln(3)x\cos(x)3^{\sin(x)}$

(i) $G'(x) = (x+1)^{\sin(x)} (\cos(x)\ln(x+1) + \sin(x)/(x+1))$

6. No solution.

7. No solution.

8. No solution.

9. $f'(0) = 2$.

10. $g'(\pi/3) = \sqrt{3} + 4\pi/3$

14.3 Derivative Applications Drill

Linear Approximations

1. Sketch $f(x) = 1 + \sin(2x)$. Compute and sketch the linear approximation to f at $x = \pi/2$.
2. Ted's hot air balloon has an outer diameter 24 feet and the material is .25 inches thick. Assuming it is spherical approximate the volume using linear approximates and compare it to the exact volume.

Limits and Infinity

1. Compute $\lim_{x \rightarrow \infty} \frac{2x - 4}{3x - 2}$.
2. Compute $\lim_{x \rightarrow 3^+} \frac{2x - 4}{x - 3}$ and $\lim_{x \rightarrow 3^-} \frac{2x - 4}{x - 3}$.
3. Compute $\lim_{x \rightarrow \infty} \frac{2x - 4}{3x^2 - 2}$.
4. Compute $\lim_{x \rightarrow \infty} \frac{\sqrt{x^2 - 3}}{x - 1}$ and $\lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 - 3}}{x - 1}$.

Graphing

1. For each of the following, find the greatest lower bound and the least upper bound for the function on the given interval.
 - (a) $f(x) = x^3 - 3x + 10$ on $[0, 2]$
 - (b) $H(x) = x + \frac{1}{x}$ on $[-2, -0.5]$
 - (c) $g(t) = \frac{5}{4 - t^2}$ on $[-6, 4]$
 - (d) $T(x) = \tan(x)$ on $[0, \frac{\pi}{2}]$
2. For each of the following graph f , f' , and f'' on the same set of axes.
 - (a) $f(x) = 6x^2 + x - 1$
 - (b) $f(x) = 1 - x^3$
3. Sketch an accurate graph of the following functions. List as much of the following information as you can: intercepts, critical points, asymptotes, intervals where f is increasing or decreasing, intervals where f is concave up or concave down, local maxima and minima, inflection points, and least upper bound and greatest lower bound.
 - (a) $F(t) = 4 - t^2$
 - (b) $f(x) = x^3 - 4x + 2$
 - (c) $G(t) = t^3 + 6t^2 + 9t + 3$
 - (d) $E(x) = e^{x^2}$
 - (e) $F(z) = \frac{1}{1 + z^4}$

- (f) $y = 3x^4 + 4x^3 - 12x^2$
- (g) $f(x) = \frac{4}{5}x^5 - \frac{13}{3}x^3 + 3x + 4$
- (h) $k(x) = x^{1/3}$ Watch for where k' is undefined.
- (i) $g(x) = x(x-2)^{2/3}$
- (j) $\frac{e^x + e^{-x}}{2}$
- (k) $p(x) = x^3 - 3x^2 + 3$
- (l) $T(x) = \sin(x) + \cos(x)$
- (m) $h(x) = \frac{2x^2}{x^2 + x - 2}$
- (n) $r(t) = \frac{\sqrt{3}}{2}t - \sin(t)$
- (o) $z = \frac{x^2 - x}{x + 1}$ Find 1 vertical and 1 slant asymptote.
- (p) $y = x^{1/3}(4 - x)$ Watch for where y' is undefined.
- (q) $y = \frac{x}{1 - x^2}$ Find 2 vertical and 1 horizontal asymptotes.
- (r) $y = \frac{\sqrt{x^2 - 3}}{x - 1}$ Watch for 2 horizontal and 1 vertical asymptotes.
- (s) $z = \frac{x^2 + 1}{x}$ What function does this graph approximate for large values of x ?
- (t) $z = \frac{x^4 - 1}{x^2}$ What function does this graph approximate for large values of x ?

Rolle's Theorem, Mean Value Theorem, and L'Hôpital's Rule

1. Verify that $f(x) = \sin(2\pi x)$ satisfies the hypothesis of the Rolle's Theorem on $[-1, 4]$ and find all values c satisfying the conclusion to the Rolle's Theorem.
2. Verify that $f(x) = x^3 - x^2 - x + 1$ satisfies the hypothesis of the Mean Value Theorem on $[-1, 2]$ and find all values c satisfying the conclusion to the MVT.
3. Evaluate each of the following limits using L'Hôpital's Rule, if L'Hôpital's rule applies.

- (a) $\lim_{x \rightarrow \infty} \frac{4x + 3}{10x + 5}$
- (b) $\lim_{x \rightarrow \infty} \frac{2x^2 - x + 3}{3x^2 + 5x - 1}$
- (c) $\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t^2 + 3}$
- (d) $\lim_{x \rightarrow \infty} \frac{(\ln(x))^2}{x}$
- (e) $\lim_{x \rightarrow 0^+} 3x \ln(x)$
- (f) $\lim_{x \rightarrow 0^+} \frac{4^{3x} - \cos(2x)}{\tan(x)}$

- (g) $\lim_{x \rightarrow 2^+} \frac{\log(x-2)}{\csc(x^2-4)}$
- (h) $\lim_{x \rightarrow 0^+} x^{\tan(x)}$
- (i) $\lim_{x \rightarrow 0^-} (x+1)^{\csc(x)}$
- (j) $\lim_{x \rightarrow \infty} \left(\frac{x}{x+3} \right)$
- (k) $\lim_{x \rightarrow \infty} (3x)^{\frac{1}{x}}$
- (l) $\lim_{x \rightarrow 0^+} (\sec(x))^{\frac{1}{x^2}}$
- (m) $\lim_{x \rightarrow 0^+} (e^x + 3x)^{\frac{1}{x}}$ (Actuarial Exam)
- (n) $\lim_{x \rightarrow \frac{\pi}{2}^+} \left(\frac{1}{\cos(x)} - \tan(x) \right)$
- (o) $\lim_{x \rightarrow 0^+} x^{\tan(x)}$
- (p) $\lim_{x \rightarrow 0^-} (1-x)^{\frac{1}{x}}$
- (q) $\lim_{x \rightarrow \frac{\pi}{2}^+} \left(x - \frac{\pi}{2} \right)^{1-\sin x}$

Max/Min Problems

1. What is the minimum perimeter possible for a rectangle with area 16 square inches?
2. A right triangle with (constant) hypotenuse h is rotated about one of its legs to sweep out a right circular cone. Find the radius that will create the cone of maximum volume.
3. Find the point (x, y) on the graph of $x = \frac{1}{y^2}$ that is nearest the origin $(0, 0)$ and is in Quadrant IV.
4. The cost per hour in dollars for fuel to operate a certain airplane is $0.021v^2$, where v is the speed in miles per hour. Additional costs are 4000 per hour. What is the speed for a 1500 mile trip that will minimize the total cost?
5. A rice silo is to be constructed in the rice fields of southeast Texas so that the silo is in the form of a cylinder topped by a hemisphere. The cost of construction per square foot of the surface area of the hemisphere is exactly twice that of the cylinder. Determine the radius and height of the silo if the volume of the silo must be 2000 cubic feet and the cost is to be minimized.

Related Rate Problems

1. Suppose that a spherical balloon is inflated at a rate of 10 cubic feet per second. At the instant that the balloon's volume is 972π cubic feet, at what rate is the surface area increasing?
2. An airplane is flying at a constant altitude of 6 miles and will fly directly over your house, where you have radar equipment. Using radar, you note that the distance between the plane is decreasing at 400 miles/hour. What is the speed of the plane when the distance from your house to the plane is 10 miles?

3. A man 6 feet tall walks at a rate of 5 feet per second toward a streetlight that is 16 feet above the ground. At what rate is the tip of his shadow moving?
4. Your snow cone has height 6 inches and radius 2 inches and is draining out through a hole in the bottom at a rate of one half cubic inch per second. You naturally wonder, what will the rate of change of the depth be when the liquid is 3 inches deep?

Derivative Applications Solutions

Linear Approximations

1. Linear approximation is $y = -2x + 1 + \pi$.
2. Exact volume = 6795.2, approximate volume = 6785.83.

Limits and Infinity

1. $\frac{2}{3}$
2. $\infty, -\infty$
3. 0
4. $1, -1$

Graphing

1. For each of the following, find the greatest lower bound and the least upper bound for the function on the given interval.
 - (a) 8, 12
 - (b) $-2.5, -2$
 - (c) no greatest lower bound or least upper bound, although there is a local min at $(0, 5/4)$
 - (d) greatest lower bound is 0, no least upper bound
2. For each of the following graph f , f' , and f'' on the same set of axes.
 - (a) no graphs, but think about the relationships between the three functions
3. Sketch an accurate graph of the following functions. List as much of the following information as you can: intercepts, critical points, asymptotes, intervals where f is increasing or decreasing, intervals where f is concave up or concave down, local maxima and minima, inflection points, and least upper bound and greatest lower bound.
 - (a) no graphs!

Rolle's Theorem, Mean Value Theorem, and L'Hôpital's Rule

1. Since all trig functions are differentiable everywhere and $f(-1) = 0 = f(4)$, f satisfies the hypothesis. $c = -\frac{3}{4}, -\frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \frac{7}{4}, \frac{9}{4}, \frac{11}{4}, \frac{13}{4}, \frac{15}{4}$ satisfy the conclusion.

2. Since all polys are differentiable everywhere, f satisfies the hypothesis. $c = \frac{1 \pm \sqrt{7}}{3}$ both satisfy the conclusion.
3. Evaluate each of the following limits using L'Hôpital's Rule if L'Hôpital's Rule applies.
- (a) $2/5$
 - (b) $2/3$
 - (c) 0
 - (d) 0
 - (e) 0
 - (f) $3\ln(4)$
 - (g) 0
 - (h) 1
 - (i) e
 - (j) 1
 - (k) 1
 - (l) e
 - (m) $e^{4/3}$
 - (n) no solution
 - (o) no solution
 - (p) no solution
 - (q) no solution

Max/Min Problems

- 1. A 4 x 4 square
- 2. Radius = $\sqrt{2/3} h$
- 3. $(\sqrt[6]{2}, -\frac{1}{\sqrt[3]{2}})$
- 4. approximately 523 miles/hour
- 5. Radius = $\sqrt[3]{750/\pi}$

Related Rate Problems

- 1. 20/9 square feet/sec
- 2. 500 mi/hr
- 3. 8 ft/sec
- 4. $\frac{3}{8\pi}$ inches per second

14.4 Integration Drill

Riemann Sums

1. Sketch $f(x) = x^2 + 2$ on $[0, 3]$. Subdivide the interval into 6 subintervals of equal length and compute the upper and lower Riemann sums of f over this partition.
2. Use the definition of the integral (as the limit of Riemann sums) to compute the area under $f(x) = x^2 + 2$ on $[0, 3]$.
3. Approximate the area under $f(x) = -\sqrt{5-x}$ on $[1, 2]$ with $n = 4$ using upper and lower sums.

Definite and Indefinite Integrals

1. Compute the following indefinite integrals.

(a) $\int 2x + \frac{3}{4}x^2 dx$

(b) $\int 1 + e^s ds$

(c) $\int \sin(x) dx$

(d) $\int \cos(y) dy$

(e) $\int \pi \cos(x) + \pi dx$

(f) $\int \sin(2t) dt$

(g) $\int 5\sqrt{x} - \frac{1}{x} dx$

(h) $\int (x^2 + 1) dx$

(i) $\int (x^2 + 1)^2 dx$

(j) $\int x(x^2 + 1)^2 dx$

(k) $\int 2x \cos(x^2) dx$

(l) $\int \frac{5}{t} + \frac{1}{5t} + 5^t dt$

(m) $\int \sin^4(t) \cos(t) dx$

(n) $\int (5x^3 - 3x^2)^8 (15x^2 - 6x) dx$

(o) $\int (y^2 - 2y)^2 (5y^4 - 2) dy$

(p) $\int t^2(t^3 - 3t) dt$

(q) $\int -\pi x + e dx$

2. Compute the following definite integrals.

(a) $\int_0^2 6 - 2x dx$

(b) $\int_2^5 x^2 - 6x + 10 dx$

(c) $\int_{-1}^3 4x^3 - x + 2 dx$

3. Evaluate each integral.

(a) $\int 2x + 1 dx$

- (b) $\int \frac{1}{3}t^3 - \sqrt{t} \, dt$
- (c) $\int 4(\sec^2(x) - \cos(x)) \, dx$
- (d) $\int z\sqrt{z} + \frac{4}{z^2} + {}^3\sqrt{4z} \, dz$
- (e) $\int \frac{x+1}{\sqrt{2x}} \, dx$
- (f) $\int \sin(t) + \sqrt{\frac{3}{t}} \, dt$
- (g) $\int x^2 \sin(x^3) \, dx$
- (h) $\int \cos^5(x) \sin(x) \, dx$
- (i) $\frac{1}{2} \int 2x\sqrt{1-x^2} \, dx$
- (j) $\int \frac{z+1}{{}^3\sqrt{3z^2+6z+5}} \, dz$
- (k) $\int_{-1}^0 (\cos(2t) + t) \, dt$
- (l) $\int_0^{\frac{\pi}{4}} \sin(\theta) - \cos(\theta) \, d\theta$
- (m) $\int_{-2}^1 (5-x)^3 \, dx$
- (n) $\int_{\pi/4}^{\pi/2} \frac{\cos(x)}{\sin^2(x)} \, dx$
- (o) $\int_{-\sqrt{3}}^2 4x\sqrt{x^2+1} \, dx$

4. Evaluate each of the following integrals, utilizing the suggested substitution.

- (a) $\int \frac{x}{\sqrt{x+5}} \, dx; \quad u(x) = \sqrt{x+5}.$
- (b) $\int x\sqrt[3]{x-1} \, dx; \quad u(x) = \sqrt[3]{x-1}.$
- (c) $\int \frac{x^3}{\sqrt{4-x^2}} \, dx; \quad u(x) = \sqrt{4-x^2}.$
- (d) $\int \frac{x^3}{\sqrt{x^2+1}} \, dx; \quad u(x) = \sqrt{x^2+1}.$
- (e) $\int \sqrt{9-x^2} \, dx; \quad x = 3\sin(u).$
- (f) $\int \frac{1}{4-x^2} \, dx; \quad x = 2\sin(u).$
- (g) $\int \frac{1}{1+x^2} \, dx; \quad x = \tan(u).$

Average Value, Mean Value, Fundamental Theorem and Arclength

1. Compute the average value of $f(x) = x^3$ over $[-3, 1]$.
2. Compute the average value $f(x) = |x|$ over $[-1, 1]$.
3. Let $f(x) = x^2 + 2x - 4$ be the continuous function on $[0, 4]$. Find c in $[0, 4]$ as stated in the Mean Value Theorem for integrals.
4. Let $F(x) = \int_{\frac{1}{2}}^x \cos(t) dt$. Compute the integral and then take the derivative of your result.
5. Compute $\frac{d}{dx} \int_{\frac{1}{2}}^x t^3 - 4t + e^t dt$
6. Use the chain rule to compute $\frac{d}{dx} \int_9^{\sqrt{x}} e^{t^2} dt$
7. Find the arc length of $y^2 = 4x^3$ from $(0, 0)$ to $(4, 16)$.

Area and Volume

1. Find the area of the region bounded by $y = x^2$, $x = -2$, $x = -1$, and the x -axis.
2. Find the area of the region bounded by $y = 2|x|$, $x = -3$, $x = 1$, and the x -axis.
3. Find the area of the region bounded by $y = x^2 - 4$ and the x -axis.
4. Find the area of the region bounded by $y = x^2$ and $y = -x^2 + 6x$.
5. Find the area of the region bounded by $y = x^3$ and $y = 4x$.
6. Find the area of the region bounded by $x = y^2$ and $x - 2y = 3$.
7. Is there a region bounded by $x - 3 = y^2 - 4y$ and $x = -y^2 + 2x + 3$?
8. Find the area of the region bounded by $x = 2y - y^2$ and $x = -y$.
9. Find the area of the region bounded by $y = x^3 + 3x^2 - x + 1$ and $y = 5x^2 + 2x + 1$.
10. Find the volume of the solid generated by revolving the region bounded by the curve $y = \sqrt{x}$, the x -axis, and the line $x = 4$ revolving about the x -axis using (a) the disk method and (b) the shell method.
11. Find the volume of the solid generated by revolving about the y -axis the region bounded by the line $y = 2x$ and the curve $y = x^2$.
12. Find the volume of the solid generated by revolving about the line $x = 6$ the region bounded by the line $y = 2$ and the curve $y = x^2$.
13. Compute the volume of the solid whose base is the disk $x^2 + y^2 \leq 4$ and whose cross sections are equilateral triangles perpendicular to the base.
14. Verify the formula for the volume of a sphere by revolving the region bounded by the circle $x^2 + y^2 = R^2$ about the y -axis with the disk method.
15. Find the volume of the solid generated by revolving the region bounded by the curve $y = (4x - 1)^{1/3}$, the x -axis, and the line $x = 4$, revolving about the x -axis using the (a) cylindrical shell method and (b) the disk method.

16. Find the volume of the solid generated by revolving the region bounded by the curve $y^2 - y = x$ and the y -axis, about x -axis.
17. Find the volume of the solid generated by revolving the region bounded by the curve $y = 8 - x^2$, $y = x^2$, about the y -axis.
18. Find the volume of the solid generated by revolving the region by the curves $y = |x| + 1$ and $y = 2$, about the x -axis.

Center of Mass

1. Let the length of a rod be 10 meters and the linear density of the rod $\rho(x)$ be written in the form $\rho(x) = ax + b$ with $x = 0$ representing the left end of the rod and $x = 10$ representing the right end of the rod. If the density of the rod is 2kg/m at the left end and 17 kg/m at the right end, find the mass and the center of mass of the rod.
2. Find the centroid (center of mass with of an object with constant density) of the region bounded by $y = x^3$ and $y = x$ in the first quadrant. What would the center of mass be if we considered only the third quadrant? What if we considered both the first and third quadrant?

Integration Solutions

Riemann Sums

1. Lower = 12.875, Upper = 17.375, Average = 15.125
2. 15
3. Upper ≈ -1.836 , Lower ≈ -1.903 , Average ≈ -1.869

Definite and Indefinite Integrals

1. Compute the following indefinite integrals.

- (a) $x^2 + \frac{1}{4}x^3 + c$
- (b) $s + e^s + c$
- (c) $-\cos(x) + c$
- (d) $\sin(y) + c$
- (e) $\pi \sin(x) + \pi x + c$
- (f) $-\frac{1}{2} \cos(2t) + c$
- (g) $\frac{10}{3}x^{3/2} - \ln(x) + c$
- (h) $\frac{1}{3}x^3 + x + c$
- (i) $\frac{1}{5}x^5 + \frac{2}{3}x^3 + x + c$

- (j) $\frac{1}{6}(x^2 + 1)^3 + c$
- (k) $\sin(x^2) + c$
- (l) $5\ln(t) + \frac{1}{5}\ln(t) + \frac{1}{\ln(5)}5^t + c$
- (m) $\frac{1}{5}\sin^5(t) + c$
- (n) $\frac{1}{9}(5x^3 - 3x^2)^9 + c$
- (o) $\frac{5}{9}y^9 - \frac{5}{2}y^8 + \frac{20}{7}y^7 - \frac{2}{5}y^5 + 2y^4 - \frac{8}{3}y^3 + c$
- (p) $\frac{1}{6}t^6 - \frac{3}{4}t^4 + c$
- (q) $-\frac{\pi}{2}x^2 + ex + c$

2. Compute the following definite integrals.

- (a) 8
- (b) 6
- (c) 84

3. Evaluate each integral.

- (a) $x + x^2 + k$
- (b) $\frac{1}{12}t^4 - \frac{2}{3}t^{3/2} + k$
- (c) $4(\tan(x) - \sin(x)) + k$
- (d) $\frac{2}{5}\sqrt{z^5} - 4z^{-1} + \frac{3}{16}\sqrt[3]{4z + k}$
- (e) $\frac{1}{3}\sqrt{x}(3 + x)\sqrt{2} + k$
- (f) $-\cos(t) + 2\sqrt{(3t)} + k$
- (g) $-\frac{1}{3}\cos(x^3) + k$
- (h) $-\frac{1}{6}(\cos(x))^6 + k$
- (i) $-\frac{1}{3}(1 - x^2)^{3/2} + k$
- (j) $\frac{1}{4}(3z^2 + 6z + 5)^{2/3} + k$
- (k) $\frac{1}{2}\sin(2) - \frac{1}{2} + k$
- (l) $1 - \sqrt{2} + k$
- (m) $\frac{2145}{4} + k$
- (n) $\sqrt{2} - 1 + k$

$$(o) \frac{20}{3} \sqrt{5} - \frac{32}{3} + k$$

4. Evaluate each of the following integrals, utilizing the suggested substitution.

$$(a) \frac{2}{3} \sqrt{x+5} (-10+x) + k$$

$$(b) \frac{3}{28} (x-1)^{4/3} (3+4x) + k$$

$$(c) -\frac{1}{3} \sqrt{4-x^2} (x^2+8) + k$$

$$(d) \frac{1}{3} \sqrt{x^2+1} (-2+x^2) + k$$

$$(e) \int \frac{1}{2} x \sqrt{9-x^2} + \frac{9}{2} \arcsin(1/3x) + k$$

$$(f) \frac{1}{4} \ln(x+2) - \frac{1}{4} \ln(x-2) + k$$

$$(g) \arctan(x) + k$$

Average Value, Mean Value, Fundamental Theorem and Arclength

1. -5
2. $\frac{1}{2}$
3. $\frac{\sqrt{93}-3}{3} \approx 2.215$
4. $F'(x) = \cos(x)$
5. $F'(x) = x^3 - 4x + e^x \, dt$
6. $F'(x) = \frac{e^x}{2\sqrt{x}}$
7. $\frac{74\sqrt{37}-2}{27} \approx 16.6$

Area and Volume

1. $\frac{7}{3}$
2. 10
3. $\frac{32}{3}$
4. 9
5. 8
6. $\frac{32}{3}$
7. *no*
8. $\frac{9}{2}$
9. $\frac{71}{6}$
10. 8π
11. $\frac{8\pi}{3}$
12. $32\sqrt{2}\pi$
13. $\frac{32\sqrt{3}}{3}$
14. $\frac{4\pi r^3}{3}$
15. no answer – attempt to estimate whether your answer makes sense
16. no answer – attempt to estimate whether your answer makes sense

17. no answer – attempt to estimate whether your answer makes sense
18. no answer – attempt to estimate whether your answer makes sense

Center of Mass

1. 95 and approximately 6.3
2. $(\frac{16}{105}, \frac{4}{15})$

14.5 Integration Techniques Drill

Here's a strategy for tackling indefinite integrals.

- Is it easy? I.e. can we just guess a function f so that f' is the integrand?
- Is there a function f in the integrand along with the derivative f' so that this is a reverse chain rule/substitution problem, for example, $\int x^2 \sin(4 - x^3) dx$?
- Is there a substitution such as $u = x + 5$ for $\int \frac{x}{\sqrt{x+5}} dx$ or $x = \tan(t)$ for $\int \frac{x^2}{\sqrt{x^2+4}} dx$?
- Can we rewrite the integrand using algebra, a trigonometric identity, long division, or partial fractions to make an easier problem?
- Is the integrand of the form $u \cdot v'$ for some functions u and v so that I can apply integration by parts?
- Is there a power or a product of powers of trigonometric functions that I know to apply certain identities to?
- Is there some clever trick that I just *know* to apply to this integral such as when I see this particular problem? For example:

$$- \int \cos^2(x) dx \text{ (trick = half-angle formula)}$$

$$- \int \sec(x) dx \text{ (trick = multiply by } \frac{\sec(x) + \tan(x)}{\sec(x) + \tan(x)})$$

1. Integrate by substitution.

$$(a) \int x^2 \sqrt{x^3 - 3} dx$$

$$(b) \int \frac{(\sqrt{x} + \pi)^4}{\sqrt{x}} dx$$

$$(c) \int x \sqrt{x+5} dx$$

$$(d) \int \sin(42x^5 - 6) 37x^4 dx$$

$$(e) \text{ Evaluate } \int x \sin(x^2) \cos(x^2) dx \text{ using } u(x) = \sin(x^2). \text{ Now evaluate } \int x \sin(x^2) \cos(x^2) dx \text{ using } u(x) = \cos(x^2). \text{ Are your answers the same? Why or why not?}$$

2. Integrate by parts.

$$(a) \int x e^{-x} dx$$

$$(b) \int x \sin(2x) dx$$

$$(c) \int x^2 \sinh(x) dx \text{ where } \sinh(x) = \frac{e^x - e^{-x}}{2} \text{ and } \cosh(x) = \frac{e^x + e^{-x}}{2}$$

3. Integrate by partial fractions.

- (a) $\int \frac{2}{x^2 - 1} dx$
- (b) $\int \frac{4x^2 + 13x - 9}{x^3 + 2x^2 - 3x} dx$
- (c) $\int \frac{6x - 11}{(x - 1)^2} dx$
- (d) $\int \frac{-19x^2 + 50x - 25}{x^2(3x - 5)} dx$

4. Integrate by trigonometric identities.

- (a) $\int \sin^3(x) dx$
- (b) $\int \cos^4(x) dx$
- (c) $\int \cos^3(x) \sin^6(x) dx$
- (d) $\int \tan^3(x) \sec^5(x) dx$
- (e) $\int \sin^2(x) \cos^2(x) dx$

5. Integrate by trigonometric substitution.

- (a) $\int \frac{\sqrt{x^2 - 9}}{x} dx$
- (b) $\int \frac{1}{4 + x^2} dx$ ($x = 2 \tan(t)$ works as does $x = 2 \sinh(t)$).
- (c) $\int \frac{x^2}{4 - x^2} dx$
- (d) $\int \frac{1}{x^3 \sqrt{x^2 - 25}} dx$

6. Integrate these Improper Integrals

- (a) $\int_0^\infty \frac{3x}{4 + x^2} dx$
- (b) $\int_0^\infty \frac{7x}{x^4 + 9} dx$
- (c) $\int_{-1}^\infty \frac{1}{x^2 + 2x + 5} dx$
- (d) $\int_0^1 \frac{-2}{x^3 - 5x^2} dx$
- (e) $\int_{-\infty}^\infty \frac{2^x}{3 + 2^x} dx$
- (f) $\int_0^e \frac{\ln(x)}{3x} dx$
- (g) Use the arc length formula to verify that the circumference the circle $x^2 + y^2 = R^2$ is $2\pi R$.
Where did you use the concept of improper integrals in this problem?

7. Evaluate the indefinite integrals using any method.

(a) $\int \sqrt{x-7} \, dx \quad \int x\sqrt{x-7} \, dx \quad \int x^2\sqrt{x-7} \, dx$

(b) $\int \frac{4t^2}{\sqrt{8-t^3}} \, dt$

(c) $\int \frac{3 \cot(t) + 2 \csc(t)}{\sin(t)} \, dt$

(d) $\int \frac{\sec^2 \sqrt{x+5}}{\sqrt{x+5}} \, dx$

(e) $\int_4^5 \frac{1}{\sqrt{x-4}} \, dx$

(f) $\int \cos(ax+b) \, dx$ (Assume a and b are constants.)

(g) $\int 3^{4x} \, dx$

(h) $\int b^{kx} \, dx$ (Assume k is a constant.)

(i) $\int (\sec(e^{2x}) + \csc(e^{2x}))^2 e^{2x} \, dx$

(j) $\int \frac{2 - e^{3x}}{e^x} \, dx$

(k) $\int \frac{e^{2x}}{e^x - 5} \, dx$

(l) $\int \frac{1}{3\sqrt{x}(4 + \sqrt{x})^4} \, dx$

(m) $\int_0^\infty x e^{-2x} \, dx$

(n) $\int \frac{\ln^3(4x)}{5x} \, dx$

(o) $\int x\sqrt{2x+1} \, dx$

8. Evaluate the indefinite integrals using any method.

(a) $\int \frac{2x-1}{x^3-5x^2-6x} \, dx$

(b) $\int \sin^3(x) \, dx$

(c) $\int \frac{x^2-x-6}{x^3-1} \, dx$

(d) $\int (16-x^2)^{3/2} \, dx$

(e) $\int \frac{3x+1}{x^2-4} \, dx$

(f) $\int \csc^6(x) \, dx$

$$(g) \int \frac{2x^3 - 4x^2 + x + \frac{21}{8}}{2x^2 - x - 2} dx$$

$$(h) \int \frac{7x^2 - 3x - 1}{x^3 - x^2 - x - 2} dx$$

$$(i) \int \cos^2(x) dx$$

$$(j) \int \frac{4}{x\sqrt{x^4 + 9}} dx$$

$$(k) \int \sin^2(x) \cos^2(x) dx$$

$$(l) \int \sin^3(x) \cos^4(x) dx$$

$$(m) \int \frac{2x - 1}{x^3 - 4x^2} dx$$

$$(n) \int \cot^5(x) \csc^2(x) dx$$

$$(o) \int \tan^3(x) dx$$

$$(p) \int \tan^4(x) \sec^4(x) dx$$

$$(q) \int \frac{2}{x^2\sqrt{x^2 - 4}} dx$$

9. Evaluate the indefinite integrals using any method.

$$(a) \int \frac{3}{\sqrt{2x - x^2}} dx$$

$$(b) \int \frac{2x^3 - 4x^2 + 3x + 1}{2x^2 - x - 2} dx$$

$$(c) \int \frac{7x^2 - 3x - 1}{x^3 - x^2 - x - 2} dx$$

$$(d) \int x \sin(3x) dx$$

$$(e) \int x^2 4^x dx$$

$$(f) \int x^5 \sqrt{x^2 + 4} dx \text{ (Question: Does } u = \sqrt{x^2 + 4} \text{ or } u = x^2 + 4 \text{ work better?)}$$

$$(g) \int \frac{\sqrt{x}}{1 + x^{1/3}} dx \text{ (Hint: Let } x = t^6 \text{)}$$

$$(h) \int \log(x) dx$$

$$(i) \int \cos^{-1}(x) dx$$

$$(j) \int \sqrt{x^2 - x - 6} dx$$

$$(k) \int x^3 \ln(x) dx$$

- (l) $\int \frac{6}{\sqrt{6x+x^2}} dx$
 (m) $\int e^x \cos(2x) dx$
 (n) $\int \cos(x) \ln(\sin(x)) dx$

Integration Techniques Solutions

1. Integrate by substitution.

- (a) $2/9 (x^3 - 3)^{3/2}$
 (b) $2/5(\sqrt{x} + \pi)^5$
 (c) $2/15 (x+5)^{3/2} (-10 + 3x)$
 (d) $-\frac{37}{210} \cos(42x^5 - 6)$
 (e) $1/4 (\sin(x^2))^2, -1/4 (\cos(x^2))^2$

2. Integrate by parts.

- (a) $-(1+x)e^{-x}$
 (b) $1/4 \sin(2x) - 1/2x \cos(2x)$
 (c) $x^2 \cosh(x) - 2x \sinh(x) + 2 \cosh(x)$

3. Integrate by partial fractions.

- (a) $-\ln|x-1| + \ln|x+1|$
 (b) $2 \ln(x-1) - \ln(x+3) + 3 \ln(x)$
 (c) $5(x-1)^{-1} + 6 \ln(x-1)$
 (d) $2/3 \ln(3x-5) - 5x^{-1} - 7 \ln(x)$

4. Integrate by trigonometric identities. Warning: Answers generated by Maple, not necessarily in simplest form...

- (a) $-\cos(x) + \frac{1}{3} \cos^3(x)$
 (b) $1/4 (\cos(x))^3 \sin(x) + 3/8 \cos(x) \sin(x) + 3/8x$
 (c) $-1/9 (\cos(x))^4 (\sin(x))^5 - \frac{5}{63} (\sin(x))^3 (\cos(x))^4 - 1/21 \sin(x) (\cos(x))^4 + \dots$
 (d) $\int 1/7 \frac{(\sin(x))^4}{(\cos(x))^7} + \frac{3}{35} \frac{(\sin(x))^4}{(\cos(x))^5} + 1/35 \frac{(\sin(x))^4}{(\cos(x))^3} - 1/35 \frac{(\sin(x))^4}{\cos(x)} - \dots$
 (e) $-1/4 (\cos(x))^3 \sin(x) + 1/8 \cos(x) \sin(x) + 1/8x$

5. Integrate by trigonometric substitution.

- (a) $\sqrt{x^2 - 9} + 3 \arctan\left(3 \frac{1}{\sqrt{x^2 - 9}}\right)$
- (b) $\frac{1}{2} \arctan(x/2)$
- (c) $-x - \ln(x - 2) + \ln(x + 2)$
- (d) $\frac{1}{50} \frac{\sqrt{x^2 - 25}}{x^2} - \frac{1}{250} \arctan\left(5 \frac{1}{\sqrt{x^2 - 25}}\right)$

6. Integrate these Improper Integrals

- (a) diverges
- (b) $\frac{7\pi}{12}$
- (c) $\frac{\pi}{4}$
- (d) diverges
- (e) diverges
- (f) diverges
- (g) the integrand is undefined at $\pm R$

7. Evaluate the indefinite integrals using any method.

- (a) $\frac{2}{3} (x - 7)^{3/2}, \frac{2}{15} (x - 7)^{3/2}, (14 + 3x) \frac{2}{105} (x - 7)^{3/2} (392 + 84x + 15x^2)$
- (b) $-\frac{8}{3} \sqrt{8 - t^3}$
- (c) $-3 (\sin(t))^{-1} - 2 \cot(t)$
- (d) $2 \tan(\sqrt{x + 5})$
- (e) 2
- (f) $\frac{\sin(ax + b)}{a}$
- (g) $\frac{1}{4} \frac{3^{4x}}{\ln(3)}$
- (h) $\frac{b^{kx}}{k \ln(b)}$
- (i) $\frac{1}{2} \tan(e^{2x}) + \ln(\tan(e^{2x})) - \frac{1}{2} \cot(e^{2x})$
- (j) $-(2e^{-x} + e^{2x}/2)$
- (k) $e^x + \ln(e^x - 5)$
- (l) $-\frac{2}{9} (4 + \sqrt{x})^{-3}$
- (m) $\frac{1}{4}$
- (n) $\frac{1}{20} (\ln(4x))^4$
- (o) $\frac{1}{15} (2x + 1)^{3/2} (-1 + 3x)$

8. No more solutions – find an integrator on the web or use a symbolic calculator.

14.6 Sequences and Series Drill

Sequences

1. Write the first five terms of the sequence in simplest form.

(a) $\{5 - \frac{2n}{3}\}_{n=2}^{\infty}$

(b) $\{\cos(\frac{n\pi}{4})\}_{n=0}^{\infty}$

(c) $\{\frac{3^n + 2}{2^n}\}_{n=1}^{\infty}$

2. Write a formula for each sequence. I.e. $x_n = \text{---}$ for $n = 1, 2, 3, \dots$

(a) $\frac{3}{5}, \frac{6}{25}, \frac{9}{125}, \frac{12}{625}, \dots$

(b) $-1, 1, -1, 1, -1, \dots$

(c) $\frac{1}{2}, -\frac{2}{3}, \frac{3}{4}, -\frac{4}{5}, \frac{5}{6}, \dots$

3. What is the least upper bound and the greatest lower bound of each given sequence? Prove it.

(a) $\{\frac{2n-2}{n}\}_{n=1}^{\infty}$

(b) $\{\frac{2n+5}{3n+1}\}_{n=2}^{\infty}$

4. Is the sequence monotonic or not? Prove it.

(a) $\{\frac{n}{n^2+1}\}_{n=1}^{\infty}$

(b) $\{\sin(\frac{2n\pi}{3})\}_{n=1}^{\infty}$

(c) $\{\frac{n-2}{n}\}_{n=1}^{\infty}$

5. Use limits to find whether or not the sequence converges.

(a) $x_n = \frac{n+2}{n^2+3n+2}$

(b) $x_n = n \sin\left(\frac{1}{n}\right)$

(c) $x_n = \left(\frac{1}{n}\right)^n$

(d) $x_n = \left(1 - \frac{1}{n}\right)^n$ Remember, $\ln$ is your friend.

6. Use the definition of limit to find the smallest N so that $\left|\frac{3}{n} - 0\right| < .001$ for $n \geq N$.

7. Assume ε is a small positive number and use the definition of limit to find the smallest N in terms of ε so that $\left|\frac{n+3}{2n} - \frac{1}{2}\right| < \varepsilon$ for $n \geq N$.

Series

1. List, in simplest form, the first three partial sums for the series and write a formula for the N^{th} partial sum.

(a) $1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \frac{1}{81} + \dots$

(b) $3 + \frac{3}{5} + \frac{3}{25} + \frac{3}{125} + \dots$

2. What does the geometric series converge to?

(a) $\sum_{n=1}^{\infty} \frac{3}{4^n}$

(b) $\sum_{n=1}^{\infty} \left(\frac{9}{10}\right)^n$

3. Use the Harmonic Series to argue that $\sum_{n=3}^{\infty} \frac{5}{3n}$ diverges.

4. Apply the integral test to determine if the series converges.

(a) $\sum_{n=2}^{\infty} \frac{2n}{(n^2 - 3)^2}$

(b) $\sum_{n=2}^{\infty} \frac{\ln(n)}{n}$

5. Determine if these series converge via one of the n^{th} term test, integral test, p-series test, comparison test, or limit comparison test.

(a) $\sum_{n=2}^{\infty} \frac{2}{1 - n^2}$

(b) $\sum_{n=1}^{\infty} \frac{1}{4n^2 + 2n}$

(c) $\sum_{n=0}^{\infty} \left(\frac{n+1}{n+3}\right)$

(d) $\sum_{n=1}^{\infty} \left(\sin\left(\frac{\pi}{4}\right)\right)^n$

(e) $\sum_{n=0}^{\infty} \frac{n+2}{3n+4}$

(f) $\sum_{n=0}^{\infty} \frac{1}{3n} + \frac{1}{2^n}$

(g) $\sum_{n=1}^{\infty} \frac{3}{2^n} + \frac{4}{3^n}$

(h) $\sum_{n=2}^{\infty} 4 + \frac{1}{n^2}$

$$(i) \sum_{n=1}^{\infty} \frac{1}{n} - \frac{1}{n+2}$$

$$(j) \sum_{n=0}^{\infty} e^{-5n}$$

$$(k) \sum_{n=1}^{\infty} \frac{3n}{n^4 + 16}$$

$$(l) \sum_{n=3}^{\infty} \sqrt{\frac{1}{3n+4}}$$

$$(m) \sum_{n=2}^{\infty} \frac{2}{n^2 - 1}$$

$$(n) \sum_{n=1}^{\infty} \frac{1}{n^n}$$

$$(o) \sum_{n=1}^{\infty} \frac{2}{\sqrt{n^2 + 2n}}$$

$$(p) \sum_{n=1}^{\infty} \frac{2}{n3^n}$$

$$(q) \sum_{n=2}^{\infty} \frac{1}{n^2 + 5n + 6}$$

$$(r) \sum_{n=2}^{\infty} \frac{1}{3^n - 1}$$

$$(s) \sum_{n=2}^{\infty} \frac{\ln(n)}{n^3}$$

$$(t) \sum_{n=2}^{\infty} \frac{\ln(n^2)}{n^2}$$

6. Determine whether the alternating series is convergent or divergent.

$$(a) \sum_{n=2}^{\infty} (-1)^n \frac{2}{n^2 - 1}$$

$$(b) \sum_{n=0}^{\infty} (-1)^n \frac{3n}{4^n}$$

$$(c) \sum_{n=1}^{\infty} (-1)^n \frac{4^n}{n^2}$$

$$(d) \sum_{n=3}^{\infty} \frac{(-5)^{n-2}}{3^{n+1}}$$

$$(e) \sum_{n=2}^{\infty} (-1)^n \left(\frac{1}{\log(n)} \right)$$

$$(f) \sum_{n=0}^{\infty} (-1)^n \left(\frac{3}{n+3} \right)$$

- (g) $\sum_{n=0}^{\infty} \cos(n\pi)$
- (h) $\sum_{n=1}^{\infty} (-1)^n \frac{3^n}{3n}$
- (i) $\sum_{n=1}^{\infty} (-1)^n \left(\frac{1}{\sqrt{n}} \right)$
- (j) $\sum_{n=1}^{\infty} (-1)^n \frac{3}{n^{10}}$
- (k) $\sum_{n=2}^{\infty} (-1)^n \left(4 + \frac{1}{n^2} \right)$
- (l) $\sum_{n=1}^{\infty} \frac{\sec(n\pi)}{n}$

7. Determine if it converges absolutely, converges conditionally, or diverges.

- (a) $\sum_{n=2}^{\infty} \frac{(-4)^n}{3^{n+1}}$
- (b) $\sum_{n=2}^{\infty} (-1)^n \frac{2}{n-1}$
- (c) $\sum_{n=2}^{\infty} (-1)^n \frac{1}{n(\log_4 n)^2}$
- (d) $\sum_{n=2}^{\infty} \left(\frac{-5}{8} \right)^n$
- (e) $\sum_{n=1}^{\infty} (-1)^n \frac{n^n}{n!}$
- (f) $\sum_{n=2}^{\infty} (-1)^n \frac{n!}{n^n}$
- (g) $\sum_{n=1}^{\infty} \cos(n\pi)$
- (h) $\sum_{n=2}^{\infty} (-1)^n \frac{3n}{n^2 - 1}$
- (i) $\sum_{n=1}^{\infty} \frac{3n}{(-3)^n}$
- (j) $\sum_{n=1}^{\infty} (-1)^n \frac{\sqrt{n}}{n+4}$
- (k) $\sum_{n=1}^{\infty} \frac{n^2}{(-2)^n}$
- (l) $\sum_{n=1}^{\infty} \frac{n^e}{(-e)^n}$

8. For each of the following power series, determine its interval of convergence.

- (a) $\sum_{n=1}^{\infty} \frac{x^n}{n+3}$
- (b) $\sum_{n=0}^{\infty} (-x)^n \frac{1}{4^n}$
- (c) $\sum_{n=1}^{\infty} (-1)^n \frac{3^n}{n^2} x^n$
- (d) $\sum_{n=0}^{\infty} \frac{(x-5)^n}{3^{n+1}}$
- (e) $\sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n^2+1}}$
- (f) $\sum_{n=1}^{\infty} \frac{x^n}{\log(n+1)}$
- (g) $\sum_{n=0}^{\infty} \frac{(x+3)^n}{n^n}$
- (h) $\sum_{n=0}^{\infty} \frac{\cos(n\pi)x^n}{n!}$
- (i) $\sum_{n=1}^{\infty} \frac{(x+4)^n}{(n+4)4^n}$
- (j) $\sum_{n=1}^{\infty} \left(1 + \frac{1}{n}\right)^{2n} x^n$
- (k) $\sum_{n=0}^{\infty} n^{2n} x^n$
- (l) $\sum_{n=1}^{\infty} \frac{(x-2)^n}{2n+2}$

9. For each of the following functions, find its Taylor or MacLaurin series and state the radius of convergence.

- (a) $f(x) = -\frac{1}{x^2}; c = 1$
- (b) $f(x) = \sin(2x); c = 0$
- (c) $f(x) = 3^x; c = 1$
- (d) $f(x) = e^x; c = 2$
- (e) $f(x) = \cos(x); c = \frac{\pi}{2}$
- (f) $f(x) = \ln(1+x); c = 0$

10. Use multiplication, substitution, differentiation, and integration whenever possible to find a series representation of each of the following functions.

- (a) $f(x) = x^2 \cos(x)$
- (b) $f(x) = \frac{\sin^2(x)}{x}$
- (c) $f(x) = 5xe^{3x}$

$$(d) f(x) = \frac{2}{(1-x)^2}$$

Sequences and Series Solutions

Sequences

1. Write the first five terms of the sequence in simplest form.

$$(a) \frac{11}{3}, \frac{9}{3}, \frac{7}{3}, \frac{5}{3}, \dots$$

$$(b) 1, \frac{\sqrt{2}}{2}, 0, \frac{-\sqrt{2}}{2}, -1, \dots$$

$$(c) \frac{5}{2}, \frac{11}{4}, \frac{29}{8}, \dots$$

2. Write a formula for each sequence. I.e. $x_n = \dots$ for $n = 1, 2, 3, \dots$

$$(a) x_n = \frac{3n}{5^n} \dots$$

$$(b) (-1)^n \dots$$

$$(c) (-1)^{n+1} \frac{n}{n+1} \dots$$

3. What is the least upper bound and the greatest lower bound? Prove it.

$$(a) \text{GLB} = 0, \text{LUB} = 2$$

$$(b) \text{GLB} = 2/3, \text{LUB} = 9/7$$

4. Is the sequence monotonic or not? Prove it.

(a) decreasing

(b) not monotonic

(c) increasing

5. Use limits to find whether or not the sequence converges.

(a) converges to 0

(b) converges to 1

(c) converges to 0

(d) converges to e^{-1}

6. $N = 3001$

7. $N =$ the first positive integer larger than $\frac{3}{2\varepsilon}$

Series

1. List, in simplest form, the first three partial sums for the series and write a formula for the N^{th} partial sum.
 - (a) $1, \frac{4}{3}, \frac{13}{9}, \frac{40}{27}, \dots, S_n = \frac{1-3^n}{1-3} \frac{1}{3^{n-1}}$
 - (b) $3, 3\frac{6}{5}, 3\frac{31}{25}, \dots, S_n = 3 \frac{1-5^n}{(1-5)5^{n-1}}$
2. What does the geometric series converge to?
 - (a) 1
 - (b) 9
3. $\sum_{n=3}^{\infty} \frac{5}{3n} = \frac{5}{3} \sum_{n=3}^{\infty} \frac{1}{n} > \sum_{n=3}^{\infty} \frac{1}{n}$ which diverges
4. Apply the integral test to determine if the series converges.
 - (a) converges
 - (b) diverges
5. Determine if these series converge via one of the n^{th} term test, integral test, p-series test, comparison test, or limit comparison test.
 - (a) converges by limit comparison with $\frac{1}{n^2}$
 - (b) converges by limit comparison with $\frac{1}{n^2}$
 - (c) diverges, n^{th} term test
 - (d) converges, geometric series
 - (e) diverges, n^{th} term test
 - (f) separate the two series, one harmonic (diverges), one geometric (converges)
 - (g) separate the two series, both geometric (converges)
 - (h) separate the two series, one constant (diverges by n^{th} term test), one converges by integral or p-series test
 - (i) limit comparison with $\frac{1}{n^2}$
 - (j) converges, geometric series
 - (k) limit comparison with $\frac{1}{n^3}$
 - (l) limit comparison with $\frac{1}{\sqrt{n}}$
 - (m) converges by limit comparison with $\frac{1}{n^2}$
 - (n) converges by comparison to $\frac{1}{n^2}$

- (o) converges by limit comparison with $\frac{1}{n}$
 - (p) converges by comparison to $\frac{1}{n^2}$
 - (q) converges by comparison to $\frac{1}{n^2}$
 - (r) converges by limit comparison with $\frac{1}{3^n}$
 - (s) converges by comparison to $\frac{1}{n^2}$
 - (t) converges, integral test
6. Determine whether the alternating series is convergent or divergent.
- (a) converges
 - (b) converges
 - (c) diverges, terms don't tend to zero
 - (d) diverges, rewrite it so that you can see that $a_n = (5/3)^n$
 - (e) converges
 - (f) converges
 - (g) diverges
 - (h) diverges
 - (i) converges
 - (j) converges
 - (k) diverges, terms don't tend to zero
 - (l) converges, rewrite it without a trig function
7. Conditionally Convergent, Absolutely Convergent, or Divergent?
- (a) D
 - (b) CC
 - (c) AC, to get absolutely convergent, convert $\log_4$ to $\ln$ and use integral test
 - (d) AC
 - (e) D
 - (f) AC, ratio test + that clever natural log trick for limits
 - (g) D
 - (h) CC
 - (i) AC
 - (j) CC
 - (k) this is a neat problem
 - (l) this is a really neat problem
8. Intervals of convergence.
- (a) $(-1, 1)$

- (b) $(-4, 4)$
- (c) $(-1/3, 1/3)$
- (d) $(2, 8)$
- (e) $(-1, 1)$
- (f) $(-1, 1)$
- (g) no interval, diverges for all $x \neq -3$
- (h) no interval, diverges for all $x \neq 0$
- (i) $(-8, 0)$
- (j) $(-1, 1)$
- (k) this is a neat problem
- (l) $(1, 3)$

9. Taylor Series (just the first few terms)

- (a) $-1 + 2(x-1) - 3(x-1)^2 + 4(x-1)^3 - 5(x-1)^4 + 6(x-1)^5$
- (b) $2x - 8x^3/3! + 32x^5/5! - 128x^7/7! \dots$
- (c) $3 + 3 \ln(3)(x-1) + \frac{3}{2}(\ln(3))^2(x-1)^2 + \frac{1}{2}(\ln(3))^3(x-1)^3 + \frac{1}{8}(\ln(3))^4(x-1)^4 + \frac{1}{40}(\ln(3))^5(x-1)^5$
- (d) $e^2 + e^2(x-2) + \frac{1}{2}e^2(x-2)^2 + \frac{1}{6}e^2(x-2)^3 + \frac{1}{24}e^2(x-2)^4 + \frac{1}{120}e^2(x-2)^5$
- (e) $-x + \frac{\pi}{2} + \frac{1}{6}\left(x - \frac{\pi}{2}\right)^3 - \frac{1}{120}\left(x - \frac{\pi}{2}\right)^5$
- (f) $x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5$

10. (a) $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2(n+1)}}{(2n)!}$
- (b) $\sum_{n=0}^{\infty} \frac{x^{2n}}{(2n+1)!}$
- (c) $5 \sum_{n=0}^{\infty} \frac{3^n}{n!} x^{n+1}$
- (d) $2 \sum_{n=0}^{\infty} (n-1)r^n$ Hint: You know the series for $\frac{1}{1-r}$.

14.7 Conic Sections, Parametric Equations, and Polar Coordinates Drill

Conics

- Complete the square for the parabola, $y = 4x^2 - 8x + 7$. Sketch the parabola, labeling the vertex, focus, and directrix. Compute the area of the region bounded by the parabola, the x-axis, the vertical line through the focus, and $x = \frac{3}{2}$.
- For the ellipse $3x^2 + 2y^2 - 12x + 4y + 2 = 0$, find
 - its center, vertices, foci, and the extremities of the minor axis;
 - an equation of the tangent to the ellipse at $(0, -1)$.
- For the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$,
 - set up a definite integral to evaluate the length of the arc of the ellipse from $(3, 0)$ to $(-3, 0)$;
 - find an approximation of the arc length in part 1.
- For the hyperbola $9y^2 - 4x^2 + 16x + 36y - 16 = 0$,
 - find its center, vertices, foci, and equations of asymptotes;
 - draw a sketch of the hyperbola and its asymptotes.
- For each of the following equations, identify whether the curve is a parabola, circle, ellipse, or hyperbola by removing the xy term from the equation by rotation of the axes.
 - $x^2 + 2xy + y^2 - 4x + 4y = 0$
 - $x^2 + 2xy + y^2 + x - y - 1 = 0$
 - $24xy - 7y^2 - 1 = 0$
 - $8x^2 - 12xy + 17y^2 + 24x - 68y - 32 = 0$

Parametrics

- Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ for each of the following parametric curves.
 - $x = a \cos(t), y = b \sin(t)$
 - $x = \sin^{-1}(t), y = \tan^{-1}(t)$
 - $x = t^e e^t, y = t \ln(t)$ (only compute dy/dx for this one)
- Let $\mathcal{C}$ be the curve defined by $x = \sin(t) + t, y = \cos(t) - t, t$ is in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. Find an equation of the tangent to $\mathcal{C}$ at the point $(0, 1)$.
- Suppose we have an object traveling in the plane with position at time t of $x = 6 - t^2$ and $y = t^2 + 4$. When is the object stopped? When is it moving left? Right? Rewrite the parametric equations as a function by eliminating time.
- Find the length:
 - $x = e^t \cos(t)$ and $y = e^t \sin(t)$ from $t = 1$ to $t = 4$

- (b) $x = 4\cos^3(t)$ and $y = 4\sin^3(t)$ from $t = \pi/2$ to $t = \pi$
5. Find the points of intersection of the curves $\mathcal{C}_1$ with parametric equations $x = t + 1$, $y = 2t^2$ and $\mathcal{C}_2$ with parametric equations $x = 2t + 1$, $y = 2t^2 + 7$. (Actuarial Exam)
6. Let $\mathcal{C}$ be the curve defined by $x = 2t^2 + t - 1$, $y = t^2 - 3t + 1$, $-2 < t < 2$. Find an equation of the tangent to $\mathcal{C}$ at the point $(0, 5)$. (Actuarial Exam)

Polars

1. For each of the following problems, find a polar equation of the graph having the given equation in the rectangular coordinate system.
- (a) $4x - 3y = 7$
- (b) $xy = 1$
- (c) $x^3 + y^3 = 3xy$
2. For each of the following problems, find an equation in the rectangular coordinate system having the given polar equation.
- (a) $r = \frac{2}{1 - 3\cos(\theta)}$
- (b) $r = 3\sin(\theta)$
- (c) $r^2 = 6\cos(2\theta)$
- (d) $r = 5\sec(\theta)$
3. For each of the following problems, find the points of intersection of the graphs of the given pairs of polar curves.
- (a) $r = 4\cos(2\theta)$; $r = 2$
- (b) $r = 1 - \cos(\theta)$; $r = \cos(\theta)$
4. For each of the following problems, find the area of the region enclosed by the graph of the given polar equation.
- (a) $r = 4 + 4\cos(\theta)$
- (b) $r = 4 - \sin(\theta)$
- (c) $r = 2\sin(3\theta)$
5. For each of the following problems, find the area of the intersection of the region enclosed by the graphs of the two given polar equations.
- (a) $r = 4 - 2\cos(\theta)$; $r = 2$
- (b) $r = 2\sin(2\theta)$; $r = 2\cos(\theta)$
6. For each of the following problems, find the area of the inner loop of the limaçon.
- (a) $r = 1 - 2\cos(\theta)$
- (b) $r = 3 - 4\sin(\theta)$

Conic Sections, Parametric Equations, and Polar Coordinates Solutions

Conics

1. $y = 4(x-1)^2 + 3$, directrix $y = 2\frac{15}{16}$, focus $= (1, 3\frac{1}{16})$, vertex $= (1, 3)$, and area $= 1\frac{2}{3}$
2. center $= (2, -1)$, vertices $= (2, -1 \pm \sqrt{6})$, foci $= (2, -1 \pm \sqrt{2})$, minor extremities $= (0, -1)$ and $(4, -1)$, equation of tangent at $(0, -1)$ is $x = 0$
3. $\frac{2}{3} \int_{-3}^3 \sqrt{1 + \frac{4}{9}(9-x^2)} dx = 5 \sin^{-1}(\frac{2\sqrt{5}}{5}) \approx 7.54$
4. center $= (2, -2)$, vertices $= (2, 0)$ and $(2, -4)$, asymptotes $y = \pm \frac{2}{3}(x-2) - 2$, foci $= (2, -2 \pm \sqrt{13})$
5. no solutions

Parametrics

1. (a) $\frac{dy}{dx} = \frac{b \cos(t)}{-a \sin(t)}, \frac{d^2y}{dx^2} = \frac{b}{a^2 \sin^3(t)}$
 (b) $\frac{dy}{dx} = \frac{1+t^2}{\sqrt{1-t^2}}, \frac{d^2y}{dx^2} = (1+t^2) \frac{2t(1-t^2) + (1+t^2)t}{\sqrt{1-t^2}}$
 (c) $\frac{dy}{dx} = \frac{\ln(t)+1}{t^{e-1}e^{t+1} + t^e e^t}$
2. $y = -\frac{1}{2}x + 1$
3. stopped at $t = 0$, moving left at all positive times, right at all negative times, $y = -x + 10$ for $x \leq 6$
4. arc lengths
 (a) $\sqrt{2}(e^4 - e^1)$
 (b) -6
5. $x = 1 \pm \sqrt{14/3}$
6. $y = \frac{5}{3}x + 5$

Polars

1. (a) $r = \frac{7}{4\cos(\theta) - 3\sin(\theta)}$
 (b) $r = \frac{1}{\pm \sqrt{\cos(\theta)\sin(\theta)}}$
 (c) $r = \frac{3\cos(\theta)\sin(\theta)}{\cos^3(\theta) + \sin^3(\theta)}$
2. (a) $y = 2\sqrt{2x^2 + 3x + 1}$

- (b) $x^2 + y^2 = 3y$
 - (c) $x^4 + 2x^2y^2 + y^4 = 6x^2 - 6y^2$
 - (d) $x = 5$
3. (a) $0 \pm \pi/12, \pi/2 \pm \pi/12, \pi \pm \pi/12, 3\pi/2 \pm \pi/12$
(b) $\pm\pi/6$
4. (a) 24π
(b) $33\pi/2$
(c) 2π
5. (a) 6π
(b) $2(3\pi - 8)/3$
6. $\frac{\pi - 3\sqrt{3}}{2}$

14.8 Vectors, Lines, and Functions Drill

Vectors

1. Let $\vec{x} = (-1, 3)$ and $\vec{y} = (4, 1)$ and graph $\vec{x}$, $\vec{y}$, $x\vec{y}$, and $\overrightarrow{x-\vec{y}}$.
2. Find the norm of each vector in the previous problem.
3. Graph $\overrightarrow{(1, -2, 1)}$ and $\overrightarrow{(-2, 4, 3)}$ and find the angle between them.

Lines

1. Sketch the vectors $x = (1, 3, 5)$ and $y = (2, 4, -3)$ and the line through these points.
2. Find a parametric equation, $\vec{l}$, for the line in the previous problem so that $\vec{l}(0) = x$ and $\vec{l}(1) = y$.
3. Plot $\vec{l}(t) = (2, -1, 4) + (1, 0, 0)t$. Assuming this represents the position of an object, compute the speed of the object.
4. Find an equation for the position of the object that has: the same speed as the object in the previous problem, the same position at time 0, and is travelling in the opposite direction.

Functions

1. Sketch a graph of $f(x, y) = 2x^2 + 3y^2$.
2. Sketch the intersection of f from the previous problem with the plane, $z = 4$.
3. Sketch the function $g(x, y) = x^3 + y^2$. Be sure and label a few points.
4. Compute the composition $g \circ \vec{l}$ where g is from the previous problem and $\vec{l}(t) = (2, -1) + (1, 0)t$.
5. Compute $(g \circ l)'(3)$ and indicate its meaning.
6. Sketch $l(t) = (0, 1) + (1, 0)t$ for $t \geq 0$.
7. Sketch $c(t) = (3\cos(t), 3\sin(t))$ for $0 \leq t \leq 2\pi$.
8. Sketch $e(t) = (4\cos(t), 2\sin(t))$ for $0 \leq t \leq 2\pi$.
9. Graph $\vec{r}(t) = (\cos(t), 2t, \sin(t))$ for $t \in [0, 6\pi]$; if $\vec{r}(t)$ is the position of an object at time t then show that the speed of the object is constant.

Vectors, Lines, and Functions Solutions

Vectors

1. a parallelogram with diagonals
2. $\|x\| = \sqrt{10}$ and $\|y\| = \sqrt{17}$

3. approximately 122 degrees

Lines

- 1.
2. $\vec{l}(t) = (1, 3, 5) + (1, 1, -8)t$ (∞ possible solns)
3. $\|\vec{l}'(t)\| = 1$
4. $\vec{p}(t) = (2, -1, 4) - (1, 0, 0)t$ (∞ possible solns)

Functions

1. a squished paraboloid
2. an ellipse
3. a water slide
4. $(t + 2)^3 + 1$
5. 75, the slope of the line (in three-space) that lies above the line $\vec{l}$ and is tangent to the graph of g at the point $(5, -1, 126)$
6. l = line
7. c = circle, radius = 3
8. e = ellipse
9. s = spiral; a spring wrapped around the y-axis in three-space; speed is $\sqrt{5}$

14.9 Cross Products, Planes, and Limits Drill

Cross Products

1. Compute the cross product of $\vec{u} = (3, 2, -1)$ and $\vec{v} = (2, \pi, -3)$ and state the significance of the resulting vector.
2. Find two vectors of unit length orthogonal to both $\vec{a} = (4, -3, 2)$ and $\vec{b} = (2, 5, -3)$.

Planes

1. Find the equation of a plane parallel to $3x + 2z = 4 - y$ and containing $(2, 4, -2)$.
2. Find the equation of the line that represents the intersection of $3z - 4x + 2y = 4$ and $2z - x = 6 - y$.
3. Find an equation for the plane containing $(1, 2, 3)$, $(2, 3, 4)$, and $(-1, 1, 1)$. Find another equation for the same plane.
4. Find the distance between the two planes: $2x - 3y + 5z = 5$ and $2x - 3y + 5z = 15$.

Limits

1. Consider the function defined by $f(x, y) = \begin{cases} \frac{x^2}{x^2 + y} & x^2 + y \neq 0 \\ 0 & (x, y) = (0, 0) \end{cases}$

Does the limit exist as $(x, y) \rightarrow (0, 0)$ along $y = kx$ exist for every $k \in \mathbb{R}$? Does the limit exist as $(x, y) \rightarrow (0, 0)$ along $y = kx^2$ exist for every $k \in \mathbb{R}$? Is the function f continuous at $(x, y) = (0, 0)$?

Cross Products, Planes, and Limits Solutions

Cross Products

1. $(\pi - 6, 7, 3\pi - 4)$
2. $\pm \frac{1}{\sqrt{933}}(-1, 16, 26)$

Planes

1. $3x + y + 2z = 6$
2. $(4, 10, 0) + (-2, -10, 4)t$ (infinitely many other ways to write this one line; check yours by substituting it into each plane)
3. $z - x = 2$
4. $\frac{10}{\sqrt{38}}$

Limits

1. No, for a chosen value of k , the limit along the path $y = kx^2$ would be $\frac{1}{1+k}$. Therefore the limit along different parabolic paths (different values of k) would yield different results.

14.10 Domains, Graphing, and Derivatives Drill

Domains of Functions

1. Find the domain of $f(x, y) = \ln(x^2 + y^2 - 1)$.
2. Find the domain of $g(x, y) = \tan^{-1}\left(\frac{x}{y}\right)$.
3. Find the domain of $h(x, y) = \frac{4}{|x| - |y|}$.
4. Find the domain of $k(x, y, z) = \frac{2xy}{z^2 + z - 1}$.
5. Find the domain of $\ell(x, y, z) = \frac{xz}{\sqrt{1 - y^2}}$.
6. Determine the domain over which $f(x, y) = \ln(x^3 y^4)$ is continuous.
7. Determine the domain over which $g(x, y) = \frac{x}{|x||y|}$ is continuous.
8. Determine the domain over which $h(x, y, z) = \frac{x^2 - 1}{z\sqrt{y^2 - 1}}$ is continuous.

Graphing Functions

1. Graph $f(x, y) = 1 - x^2 + y^2$.
2. Graph $g(x, y) = x + y$.
3. Graph $h(x, y) = \frac{x^2}{9} + \frac{y^2}{4}$.
4. Graph $k(x, y) = |x| - |y|$.
5. Graph $r(x, y) = \sin(x)$.

Partial Derivatives and Gradients

1. Let $g(x, y) = x^3 - 4\log_7 x^2 + \sin^{-1}(xy)$ and compute g_x .
2. Let $g(x, y) = \frac{x^2}{\sin(xy)}$ and compute g_y .
3. Let $g(x, y) = e^{xy^2}$ and compute ∇g .
4. Let $g(x, y) = x \ln(y) - 4xy + x$ and compute $g_x(1, 1)$ and $g_y(1, 1)$.
5. Let $g(x, y) = 5x^2 y \sin(x - y)$ and compute $g_y(\pi, 2\pi)$.
6. Let $h(x, y, z) = \sqrt{2xz - 5y} + \cos^3(z \sin(x))$ and compute ∇h .
7. Let $h(x, y, z) = z^{xy}$ and compute $h_x(2, 3, 4)$.
8. Find f_{xx} , f_{zy} , and f_{xzy} for $f(x, y, z) = \frac{y}{x^3} - \sin(zy) - 3z^3$.

Directional Derivatives

Remember: Directional vectors should be unit vectors.

1. Using the definition of directional derivative, compute the derivative of $f(x,y) = x - y^2$ at $(1,2)$ in the direction, $(1,1)$.
2. Find $D_{\vec{u}} f(p)$ for $f(x,y) = e^{xy} + 2x^2y^3$; $p = (3,1)$; $\vec{u}$ is a unit vector parallel to $\vec{v} = (-5,12)$.
3. Find $D_{\vec{u}} f(p)$ for $f(x,y) = e^{xy} + \ln(xy)$; $p = (1,2)$; $\vec{u}$ is a unit vector which makes an angle $\frac{-\pi}{3}$ from the positive x axis.
4. Find $D_{\vec{u}} f(p)$ for $f(x,y,z) = x^2y - 4y^2z + xyz^2$; $p = (-2,1,-1)$; $\vec{u}$ is a unit vector parallel to the vector $\vec{AB}$ where $A = (2,1,-5)$ and $B = (-2,4,3)$.

Derivatives

For each of the following functions, compute the indicated “derivative” of the function. Because of the differing domains of the functions, the derivative could be a function (a partial derivative), a vector of functions (a gradient), or a matrix of functions (the ‘total’ derivative)!

1. $f(x,y) = x^2y^3$
2. $g(x,y) = \left(\frac{x^2}{y} - 3xy, \sin\left(\frac{x}{y}\right)\right)$
3. $h(x,y) = x^2 - e^{xy\sqrt{z}} + \sinh(yz)$
4. $r(s,t,u) = (st, s^2tu, \sqrt{stu}, \ln(st^2u))$

Chain Rule

1. Let $f(x,y) = 2x^2 - 7y$ and $\vec{g}(t) = (\sin(t), \cos(t))$. Compute the derivative of $f \circ g$ in two ways. First compose f and g and take the derivative. Second, apply the chain rule. Verify that your solutions are the same.
2. Let $f(x,y) = 4x^3y + e^{3y} + \frac{2}{x}$, $x(t) = t^2$, $y(t) = 4t - 3$, and $g(t) = (x(t), y(t))$. Compute $(f \circ g)'(-1)$.
3. For each of the following problems, find $\frac{dg}{dt}$ and evaluate at the given value of t .
 - (a) $g(x,y) = 3xy + e^xy^2$ where $x(t) = 4t^2 + t$, $y(t) = 6 + 5t$, and $t = 0$
 - (b) $g(x,y,z) = x^3y + xz + \frac{x}{y-z}$ where $x(t) = t^2 + 3$, $y(t) = 4t - t^2$, $z(t) = \cos(t - 3\pi)$, and $t = 0$
4. Let $f(x,y) = xy \ln(x)$ and $\vec{g}(s,t) = (2st, t - s^3)$. State the domain f , g , and $f \circ g$. Compute the gradient of $(f \circ g)$.
5. For each of the following problems, find f_s and f_t (i.e. $\frac{\partial f}{\partial s}$ and $\frac{\partial f}{\partial t}$).
 - (a) $f(x,y) = 2x - y^2$, $x(s,t) = s \cos(t)$, and $y(s,t) = (s+t)e^t$

(b) $f(x, y, z) = (x + 2y + 3z)^4$ and $x(s, t) = s + t$, $y(s, t) = s - t$, $z(s, t) = st$

6. For each of the following problems, find $\frac{\partial g}{\partial u}$, $\frac{\partial g}{\partial v}$, and $\frac{\partial g}{\partial w}$ (i.e. g_u , g_v , and g_w).

(a) $g(x, y) = (x + y) \ln(xy)$, $x(u, v, w) = u + v - 3w$, and $y(u, v, w) = uv + 3w$

(b) $g(x, y, z) = yz + xz + xy$, $x(u, v, w) = u + v - 3vw$, $y(u, v, w) = v + w + 4uw$, and $z(u, v, w) = u + w - 5uv$

7. For each of the following three problems, find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ at the indicated point.

(a) $z^3 - xy + 2yz + y^3 - 3 = 0$; $(1, 1, 1)$

(b) $\frac{1}{x^2} + \frac{1}{y^2} + \frac{1}{z^2} = \frac{49}{36}$; $(-1, 2, 3)$

(c) $ye^{xyz} \cos(2xz) = 1$; $(\pi, 1, 4)$

Tangent Lines and Planes

- Find the (shortest) distance from the point $(0, 1, 0)$ to the plane $x + 2y + 3z = 4$.
- Find the equation of the tangent plane to the function $z = x^2 - 4y^2$ at $(3, 1, 5)$.
- Find the equation of the tangent plane to $z + 1 = xe^y \cos(z)$ at the point $(1, 0, 0)$.
- Find the equation of the line normal (perpendicular) to $x^2 + 2y^2 + 3z^2 = 6$ and passing through $(1, -1, 1)$.
- Find the tangent plane approximation of $h(x, y) = x + x \ln(xy)$ when $x = e$ and $y = 1$.

Domains, Graphing, and Derivative Solutions

Domains of Functions

- $\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 > 1\}$
- $\{(x, y) \in \mathbb{R}^2 : y \neq 0\}$
- $\{(x, y) \in \mathbb{R}^2 : |x| \neq |y|\}$
- all of $\mathbb{R}^3$ except where z equals the Pisot numbers
- $\{(x, y) \in \mathbb{R}^2 : |y| < 1\}$

Graphing Functions

- a saddle centered at $(0, 0, 1)$
- a plane
- squished paraboloid
- planar saddle with a 'point' at $(0, 0, 0)$

5. a three dimensional sine wave

Partial Derivatives and Gradients

$$1. g_x(x, y) = 3x^2 - \frac{8}{x \ln(7)} + \frac{y}{\sqrt{1 - (xy)^2}}$$

$$2. g_y(x, y) = \frac{-x^3 \cos(xy)}{\sin^2(xy)}$$

$$3. \nabla g(x, y) = (y^2 e^{xy^2}, 2xy e^{xy^2})$$

$$4. g_x(1, 1) = g_y(1, 1) = -3$$

$$5. g_y(\pi, 2\pi) = 2\pi 5\pi^2$$

$$6. \nabla h(x, y, z) = \left(\frac{z}{\sqrt{2xz - 5y}} - 3z \cos(x) \cos^2(z \sin(x)) \sin(z \sin(x)), \frac{-5}{\sqrt{2xz - 5y}}, \right. \\ \left. \frac{x}{\sqrt{2xz - 5y}} - 3 \sin(x) \cos^2(z \sin(x)) \sin(z \sin(x)) \right)$$

$$7. h_x(2, 3, 4) \approx 17,035$$

$$8. f_{xx} = \frac{12y}{x^5}, f_{zy} = -\cos(zy) + zy \sin(zy), f_{zxzy} = 0$$

Directional Derivatives

Remember: Directions vectors should be unit vectors.

1. $\frac{-3}{\sqrt{2}}$
2. $\frac{1}{13}(588 + 31e^3)$
3. $(1 - \frac{\sqrt{3}}{2})(e^2 + \frac{1}{2})$
4. $\frac{42}{\sqrt{89}}$

Derivatives

1. $\nabla f(x, y) = (2xy^3, 3x^2y^2)$
2. $Dg(x, y) = \begin{pmatrix} \frac{2x}{y} - 3y & -\frac{x^2}{y^2} - 3x \\ \frac{1}{y} \cos(\frac{x}{y}) & -\frac{x}{y^2} \cos(\frac{x}{y}) \end{pmatrix}$
3. $\nabla h(x, y) = (2x - y\sqrt{z}e^{xy\sqrt{z}}, -x\sqrt{z}e^{xy\sqrt{z}} + z \cosh(yz))$
4. $Dr(s, t, u) = \begin{pmatrix} t & s & 0 \\ \frac{2stu}{2\sqrt{stu}} & \frac{s^2u}{2\sqrt{stu}} & \frac{s^2t}{2\sqrt{stu}} \\ \frac{t^2u}{st^2u} & \frac{2stu}{st^2u} & \frac{st^2}{st^2u} \end{pmatrix}$

Chain Rule

1. $(f \circ g)'(t) = \sin(t)(4\cos(t) + 7)$
2. $(f \circ g)'(-1) \approx 188$
3. (a) 84
4. $\nabla(f \circ g) = ((2t^2 - 8s^3t)\ln(2st) + (2st^2 - 2s^4t)\frac{1}{s}, (4st - 2s^4)\ln(2st) + (2st^2 - 2s^4t)\frac{1}{t})$
5. (a) $f_s(s, t) = 2\cos(t) - 2(s+t)e^{2t}$ and $f_t(s, t) = -2s\sin(t) - 2(s+t)e^{2t} - 2(s+t)^2e^{2t}$

Tangent Lines and Planes

1. $\frac{\sqrt{14}}{7}$
2. $6x - 8y - z = 5$
3. $x + y - z = 1$
4. $L(t) = (2, -4, 6)t + (1, -1, 1)$
5. $3x + ey - z = 2e$

14.11 Optimization and Lagrange Multipliers Drill

Critical Points

- For each of the following functions, find the critical points and determine if they are maxima, minima, or saddle points.
 - $f(x, y) = 1 - x^2 - y^2$
 - $g(x, y) = e^{-x} \sin y$
 - $F(x, y) = 2x^2 + 2xy + y^2 - 2x - 2y + 5$
 - $g(x, y) = x^2 + xy + y^2$
 - $z = 8x^3 - 24xy + y^3$
- For each of the following functions, find the absolute extrema of the function on the given closed and bounded set R in $\mathbb{R}^2$.
 - $f(x, y) = 2x^2 - y^2$; $R = \{(x, y) : x^2 + y^2 \leq 1\}$
 - $g(x, y) = x^2 + 3y^2 - 4x + 2y - 3$; $R = \{(x, y) : 0 \leq x \leq 3, -3 \leq y \leq 0\}$
- Find the direction at which the maximum rate of change of $g(x, y) = \ln(xy) - 3x + 2y$ at $p = (3, 2)$ will occur and find the maximum rate of change.
- Find the direction in which the function $f(x, y) = x^3 - y^5$ increases the fastest at the point $(2, 4)$.
- For each of the following four problems, find all critical points of f and classify these critical points as relative maxima, relative minima, or saddle points using the second derivative test whenever possible.
 - $f(x, y) = xy^2 - 6x^2 - 3y^2$
 - $f(x, y) = \frac{9x}{x^2 + y^2 + 1}$
 - $g(x, y) = x^2 + y^3 + \frac{768}{x + y}$

Optimization and Lagrange Multipliers

- Find all extrema of $f(x, y) = 1 - x^2 - y^2$ subject to $x + y = 1$, $x \geq 0$, and $y \geq 0$.
- Find all extrema of $f(x, y) = 1 - x^2 - y^2$ subject to $x + y \leq 1$, $x \geq 0$, and $y \geq 0$.
- Find the absolute extrema of the function $g(x, y) = 2\sin(x) + 5\cos(y)$ on the rectangular region $R = \{(x, y) : 0 \leq x \leq 2, 0 \leq y \leq 5\}$.
- Find the minimum value of $z = x^2 + y^2$ subject to $x + y = 24$.
- Find the extreme values of $f(x, y) = 2x^2 + y^2 - y$ subject to $x^2 + y^2 = 4$ using Lagrange multipliers.
- Find three positive numbers whose sum is 123 such that their product is as large as possible.
- A container in $\mathbb{R}^3$ has the shape of a cube with each edge length 1. A (triangular) plate is placed in the container so that it intersects the cube in the plane $x + y + z = 1$. If the container is heated so that the temperature at each point is given by $T(x, y, z) = 4 - 2x^2 - y^2 - z^2$ in hundreds of degrees, what are the hottest and coldest points on the plate?

8. A company has three production plants, each manufacturing the same product. If plant A produces x units at the cost of $\$(x^2 + 2,000)$, plant B produces y units at the cost of $\$(2y^2 + 3,000)$, and plant C produces z units at the cost of $\$(z^2 + 4,000)$. If there is an order for 11,000 units to be filled, determine how the production should be arranged among these three plants so that the total production cost can be minimized.

Optimization and Lagrange Multipliers Solutions

Critical Points

- $(0, 0, 1)$ is a max
 - none, why?
 - $(0, 1, 4)$ is a min
 - $(0, 0, 0)$ is a min
- $(\pm 1, 0, 2)$ are local minima and $(0, \pm 1, -1)$ are local maxima
 - all points to consider should be: $(2, -1/3)$, $(0, 0)$, $(3, 0)$, $(3, -3)$, $(0, -3)$, $(2, 0)$, $(0, -1/3)$, $(3, -1/3)$, and $(2, -3)$; $(2, -1/3, -22/3)$ is a min; $(0, -3, 18)$ is a max
- $(-8/3, 5/2)$ and $\sqrt{481}/6$
- $(3, -320)$
 - $(3, \pm 6, -54)$ are saddles and $(0, 0, 0)$ are all maxima
 - $(0, 1, 9/2)$ (D is long – use software!)
 - icky algebra – use software!

Optimization and Lagrange Multipliers

- $(1/2, 1/2, 1/2)$ is the max, each of $(1, 0, 0)$ and $(0, 1, 0)$ is a min
- $(0, 0, 1)$ is the max, each of $(1, 0, 0)$ and $(0, 1, 0)$ is a min
- there is one saddle on the interior, but no extrema on the interior; the extrema occur on the boundaries at $(0, 0)$, $(\pi/2, 5)$, $(2, 0)$, and $(\pi/2, 0)$
- $(12, 12, 288)$
- $(0, \pm 2)$ and $(\pm \frac{\sqrt{15}}{2}, 1/2)$
- $x = 41, y = 41, z = 41$

14.12 Integration Drill

Basics

1. Evaluate $\int x^3y - 3x \, dx$ and $\int_1^3 x^3y - 3x \, dy$
2. Evaluate $\int xye^{xy} \, dx$ and $\int_{-1}^1 xye^{xy} \, dy$
3. Compute the area of the region bounded by the parabola $y = x^2 - 2$ and the line $y = x$ by first computing a single integral with respect to x and then computing a single integral with respect to y .
4. Compute the area bounded by $y = x^3$ and $y = 5x$ in four ways. (a) Single integral with respect to x , (b) single integral w.r.t. y , (c) double integral, $dx \, dy$, and (d) double integral, $dy \, dx$.

Regions

1. Sketch the region $\phi = \pi/6$ in spherical coordinates.
2. Sketch the region bounded by $r = 1$ and $r = 2\sin(\theta)$ in polar coordinates.
3. Sketch the region bounded by $x^2 + y^2 \leq 9$ and write in polar coordinates.
4. Sketch the region between $x^2 + y^2 = 25$, $x^2 + y^2 = 4$, and $x \geq 0$ and write in polar coordinates.
5. Find the area of the region D bounded by $y = \cos(x)$ and $y = \sin(x)$ on the interval $\left[0, \frac{\pi}{4}\right]$.

Integration

1. Compute $\int_0^2 \int_0^3 x^2y^2 - 3xy^5 \, dy \, dx$ and $\int_0^3 \int_0^2 x^2y^2 - 3xy^5 \, dx \, dy$. Are they equal? What theorem is this an example of?
2. Evaluate $\int_0^1 \int_0^y e^{y^2} \, dx \, dy$.
3. Compute $\int_R x^2 e^{xy} dA$ where $R = \{(x, y) : 0 \leq x \leq 3, 0 \leq y \leq 2\}$.
4. Compute $\int_0^3 \int_0^2 \sqrt{2x+y} \, dy \, dx$.
5. Compute $\int_R \frac{\ln \sqrt{y}}{xy} dA$ where $R = \{(x, y) : 1 \leq x \leq 4, 1 \leq y \leq e\}$.
6. Evaluate $\int_D 160xy^3 \, dA$ where D is the region bounded by $y = x^2$ and $y = \sqrt{x}$.
7. Sketch and find the volume of the solid bounded above by the plane $z = y$ and below in the xy -plane by the part of the disk $x^2 + y^2 \leq 1$.
8. Sketch the region, D , that is bounded by $x = y^2$ and $x = 3 - 2y^2$ and evaluate $\int_D (y^2 - x) dA$
9. Compute $\int_0^2 \int_0^{\sin(x)} y \cos(x) \, dy \, dx$.

10. Determine the endpoints of integration for $\int \int_S e^{xy} dA$ where S is the region bounded by $y = \sqrt{x}$ and $y = \frac{x}{9}$. Don't integrate.
11. Determine the endpoints of integration for $\int \int_S dA$ where S is bounded by the $x = y^2 + 4y$ and $x = 3y + 2$.
12. Determine the endpoints of integration for $\int \int_S 2x dA$ where S is the region bounded by $yx^2 = 1$, $y = x$, $x = 2$, and $y = 0$.
13. Evaluate $\int \int_D dA$ where D is the region bounded by the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$

Polar and Spherical Coordinate Integration

1. Sketch the region D between $r = \cos\left(\frac{\theta}{2}\right)$ and $x^2 + y^2 = 1$ with $0 \leq \theta \leq \pi$. Evaluate $\int_D 1 dA$.
2. Consider $\int_R x^2 + y^2 + 1 dA$ where $R = \{(x, y) : x \geq 0, 9 \leq x^2 + y^2 \leq 16\}$. Write the integral in both rectangular and polar coordinates. Compute each to verify your answer.
3. Find the volume of the solid bounded by the cone $\phi = \frac{\pi}{6}$ and the sphere $\rho = 4$.

Coordinate Transformations

1. Find the Jacobian for the transformation: $x = u^2 + v^2 + w$, $y = uw - v$, and $z = \ln(w) - \frac{v}{u}$
2. Evaluate $\int_0^4 \int_{y/2}^{(y/2)+1} \frac{2x-y}{2} dx dy$ using $u = \frac{2x-y}{2}$ and $v = \frac{y}{2}$.
3. Use the transformation $x = \frac{u}{v}$ and $y = v$ to rewrite (but not evaluate) the double integral $\int \int \sqrt{xy^3} dx dy$ over the region in the plane bounded by the x -axis, the y -axis, and the lines $y = -2x + 2$ and $x + y = 7$.
4. Compute $\int_0^1 \int_0^{y^2} (1-y) \sin\left(\frac{x}{y}\right) dx dy$ using $u = \frac{x}{y}$ and $v = 1 - y$.
5. Write an integral in rectangular coordinates that represents the area enclosed by the ellipse $\frac{x^2}{16} + \frac{y^2}{49} = 1$. Now, compute this integral by using the transformation $x = 4u$ and $y = 7v$.
6. Find the volume of the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ using the transformation $x = au$, $y = bv$, $z = cw$.
7. Evaluate the triple integral $\int_0^6 \int_0^8 \int_{\frac{y}{2}}^{\frac{y}{2}+4} \frac{2x-y}{2} + \frac{y}{6} + \frac{z}{3} dx dy dz$ using the transformation $u = \frac{2x-y}{2}$, $v = \frac{y}{2}$, and $w = \frac{z}{3}$.

Triple Integrals, Cylindrical, and Spherical Coordinates

1. Sketch and find the volume of the solid formed by $f(x,y) = 4x + 2y$ above the region in the xy -plane bounded by $x = 2$, $x = 4$, $y = -x$, $y = x^2$.
2. Fill in the blanks: $\int_0^1 \int_0^{\frac{1-x}{2}} \int_0^{1-x-2y} f(x,y,z) \, dz \, dy \, dx$
 $= \int_-^- \int_-^- \int_-^- f(x,y,z) \, dy \, dz \, dx$
 $= \int_-^- \int_-^- \int_-^- f(x,y,z) \, dy \, dx \, dz$
 $= \int_-^- \int_-^- \int_-^- f(x,y,z) \, dx \, dy \, dz$
3. $\int_0^6 \int_0^{\frac{12-2x}{3}} \int_0^{\frac{12-2x-3y}{4}} f \, dz \, dy \, dx = \int_-^- \int_-^- \int_-^- f \, dy \, dx \, dz = \int_-^- \int_-^- \int_-^- f \, dx \, dz \, dy$
4. $\int_0^{\frac{1}{6}} \int_0^{\frac{\sqrt{1-36x^2}}{3}} \int_0^{\frac{\sqrt{1-36x^2-9y^2}}{2}} f \, dz \, dy \, dx = \int_-^- \int_-^- \int_-^- f \, dx \, dz \, dy$
5. Find the volume of the solid bounded by $x^2 + y^2 + z = 8$ and $z = 4$.
6. Evaluate $\int_0^2 \int_0^{\sqrt{4-x^2}} \int_0^{\sqrt{4-x^2-y^2}} z \sqrt{4-x^2-y^2} \, dz \, dy \, dx$ by converting to (a) cylindrical coordinates; (b) spherical coordinates.

Application - Center of Mass

1. Suppose $\delta(x,y) = x + y$ is the density function of a thin sheet of material bounded by the curve $x^2 = 4y$ and $x + y = 8$. Find its total mass. Find its first moments. Find its center of mass. Find its second moments.

Integration Solutions

Basics

1. $\frac{1}{4}x^4y - \frac{3}{2}x^2$ and $4x^3 - 6x$
2. $e^{xy}(x - \frac{1}{y})$ and $e^x(1 - \frac{1}{x}) + e^{-x}(1 + \frac{1}{x})$
3. $9/2$
4. $25/4$

Regions

1. it's a cone
2. it's the intersection of two circles
3. $r \leq 3$

4. $2 \leq r \leq 5$ and $\theta \in [-\pi/2, \pi/2]$

5. $\sqrt{2} - 1$

Integration

1. -705 , Fubini's Theorem

2. $\frac{1}{2}(e - 1)$

3. $-\frac{1}{4}(17 - 5e^6)$

4. ≈ 11.62

5. $\approx .346$

6. 6

7. $\frac{2}{3}$

8. $\frac{24}{5}$

9. $\frac{1}{6} \sin^3(2)$

10. $\int_0^{81} \int_{\frac{x}{9}}^{\sqrt{x}} e^{xy} dA$

11. 4.5

12. $\frac{2}{3} + 2\ln(2)$

13. 6π

Polar and Spherical Coordinate Integration

1. $\frac{\pi}{4}$

2. $\frac{189\pi}{4}$

3. $\frac{-64(\sqrt{3} - 2)\pi}{3}$

Coordinate Transformations

1. $-2u(\frac{1}{w} - 1) - 2v + \frac{1}{u}(2v^2 - w) + v/u^2$

2. r

3. $\rho^2 \sin(\phi)$

4. 2

5. $\int_0^7 \int_{7v-v^2}^{v-v^2/2} \sqrt{u} du dv$

6. area of ellipse is πab in this case 28π (just like circle)

7. volume of ellipsoid is $4/3\pi abc$ (just like sphere!)

Triple Integrals, Cylindrical, and Spherical Coordinates

1. $2472/5$

$$2. \int_0^1 \int_0^{1-x} \int_0^{\frac{1-x-z}{2}} f(x,y,z) \, dy \, dz \, dx$$
$$\int_0^1 \int_0^{1-z} \int_0^{\frac{1-x-z}{2}} f(x,y,z) \, dx \, dz \, dy$$
$$\int_0^1 \int_0^{\frac{1-z}{2}} \int_0^{1-z-2y} f(x,y,z) \, dx \, dy \, dz$$

$$3. \int_0^3 \int_0^{-2x+6} \int_0^{\frac{-4x-2x+12}{3}} f(x,y,z) \, dy \, dx \, dz$$
$$\int_0^4 \int_0^{-\frac{3}{4}y+3} \int_0^{\frac{-4z-3y+12}{2}} f(x,y,z) \, dx \, dz \, dy$$

$$4. \int_0^{\frac{1}{3}} \int_0^{\sqrt{\frac{1-9y^2}{2}}} \int_0^{\frac{\sqrt{1-9y^2-4z^2}}{6}} f(x,y,z) \, dy \, dx \, dz$$

5. 8π

14.13 Line Integrals, Flux, Divergence, and Gauss' Theorem Drill

Note. In this section, we write the functions and integrals using many different notations. If you are unsure about the meaning of a notation, please ask!

Vector Fields, Curl, and Divergence

1. Sketch the vector field, $\vec{f}(x,y) = x^2\vec{i} + \vec{j}$.
2. Sketch the vector field, $\vec{g}(x,y) = (x, -y)$.
3. Sketch the vector field, $\vec{h}(x,y,z) = y\vec{j}$.
4. Compute the divergence and curl of the vector field, $\vec{f}(x,y,z) = (y^2z, x^3 + z + y, \cos(xyz))$.
5. Find g satisfying $\nabla g = F$ if it exists. $F(x, y) = (ye^{xy} + 2x)\vec{i} + (xe^{xy} - 2y)\vec{j}$
6. Find g satisfying $\nabla g = F$ if it exists. $F(x, y) = (e^x \sin(y), e^x \cos(y))$
7. Is $\vec{f}(x,y) = (2xy, x^2)$ a conservative vector field? If so, find a potential for it.
8. Is $\vec{h}(x,y) = (y \cos(x), \sin(x))$ a conservative vector field? If so, find a potential for it.
9. Is $\vec{s}(x,y,z) = (e^x \sin z + yz)\vec{i} + (xz + y)\vec{j} + (e^x \cos z + xy + z^2)\vec{k}$ a conservative vector field? If so, find a potential for it.
10. Is $\vec{g}(x,y) = 2x\vec{i} + y\vec{j}$ a conservative vector field? If so, find a potential for it.
11. Show that $\mathbf{F}(x,y,z) = x\vec{i} + y\vec{j} + 2z\vec{k}$ is conservative and find a function f such that $\mathbf{F} = \nabla f$.

Line Integrals over Scalar Fields

1. Let $f(x,y) = x + y$ and $\vec{c}$ be the unit circle in $\mathbb{R}^2$. Evaluate $\int_{\vec{c}} f \, ds$. Recall that ds means to evaluate with respect to the arc length.
2. Evaluate $\int_{\vec{c}} \sqrt{xy + 2y + 2} \, ds$ with $\vec{c}$ the line segment from $(0, 1)$ to $(0, -1)$.
3. Evaluate $\int_{\vec{c}} (x - y + z - 2) \, ds$ where $\vec{c}$ is the line segment from $(0, 1, 1)$ to $(1, 0, 1)$.

Line Integrals over Vector Fields

1. Compute $\int_{\vec{r}} f \cdot d\vec{r}$ where $f(x,y) = (y, x^2)$ and $\vec{r}(t) = (4 - t, 4t - t^2)$ for $0 \leq t \leq 3$.
2. Compute $\int_{\vec{r}} f \cdot d\vec{r}$ where $f(x,y) = (y, x^2)$ and $\vec{r}(t) = (t, 4t - t^2)$ for $1 \leq t \leq 4$.
3. Compute $\int_{\vec{r}} F \cdot d\vec{r}$ where $F(x,y) = (-\frac{1}{2}x, -\frac{1}{2}y, \frac{1}{4})$ and $\vec{r}(t) = (\cos(t), \sin(t), t)$, $1 \leq t \leq 4$.

Divergence Theorem and Green's Theorem

1. Verify the divergence theorem for the flow $f(x,y) = (0, y)$ over the circle, $x^2 + y^2 = 5$.
2. Let $f(x,y) = (u(x,y), v(x,y)) = (-x^2y, xy^2)$ and $\vec{c} = \{(x,y) : x^2 + y^2 = 9\}$ and D be the region bounded by $\vec{c}$. Verify Green's Theorem by evaluating both $\oint_{\vec{c}} f(\vec{x}) d\vec{x}$ and $\int \int_D v_x(x,y) - u_y(x,y) dA$.

3. Verify Green's Theorem where $f(x,y) = (4xy, y^2)$ and $\vec{c}$ is the curve $y = x^3$ from the $(0,0)$ to $(2,8)$ and the line segment from $(2,8)$ to $(0,0)$.
4. If you want more practice on verifying Green's and Gauss' theorems, then note that each problem that asks you to verify Gauss' theorem could have asked you to verify Green's theorem and vice-versa. You won't need solutions because you are computing both sides of the equation and they must be equal if all your integration is correct.

Divergence Theorem in Three Dimensions

1. Verify the divergence theorem for $f(x,y,z) = (xy, z, x+y)$ over the region in the first octant bounded by $y = 4$, $z = 4 - x$, $z = 0$, $y = 0$, and $x = 0$.
2. Verify the divergence theorem for $f(x,y,z) = (2x, -2y, z^2)$ over the region $[0,3] \times [0,3] \times [0,3]$.

Line Integrals, Flux, Divergence, and Gauss' Theorem Solutions

Vector Fields, Curl, and Divergence

1. no sketch
2. no sketch
3. no sketch
4. no solution
5. $g(x,y) = e^{xy} + x^2 - y^2$
6. $g(x,y) = e^x \sin(y)$
7. yes, $g(x,y) = x^2y$
8. yes, $g(x,y) = y \sin(x)$
9. yes, $g(x,y,z) = e^x \sin(z) + xyz + \frac{1}{2}y^2 + \frac{1}{3}z^3$
10. no solution
11. no solution

Line Integrals over Scalar Fields

1. 0
2. $8/3$
3. $-\sqrt{2}$

Line Integrals over Vector Fields

1. $69/2$

2. $-69/2$

3. $3\pi/4$

Divergence Theorem and Green's Theorem

1. If both sides are equal, you probably have the right answer!

2. $81\pi/2$

3. $-256/15$

4. no solution

Divergence Theorem in Three Dimensions

1. Hopefully you got the same answer both ways!

2. Hopefully you got the same answer both ways!

Chapter 15

Appendices

15.1 Additional Materials on Limits

Our definition of limits in this book is an intuitive one. The statement that “ $f(x)$ gets close to $f(a)$ as x gets close to a ” is not mathematically precise because of the word ‘close.’ What does ‘close’ mean? Is .5 close to zero? Is .001 close to zero? Mathematicians spent hundreds of years defining what ‘close’ meant and the following definition was the result of the work of Newton and Leibnitz, among others. This definition gave rise to precise definitions for continuity, derivatives, and integrals, which had theretofore been defined only intuitively. This definition made precise the entire field of analysis which is one of the most applied fields in all of mathematics. If the notion seems non-trivial or the definition challenging, then you are in the company of great men. If you find success in understanding the next few problems, then you are in the company of a very few and should exploit your understanding of such deep concepts with a major or minor in the subject.

Definition 1. *If L and a are numbers, then we say that **the limit of f at a is L** if for any two horizontal lines $y = H$ and $y = K$ with the line $y = L$ between them, there are two vertical lines, $x = h$ and $x = k$ with $x = a$ between them, so that if t is any number between h and k (other than a), then $f(t)$ is between H and K .*

Here is a sketch of how we might use this definition to *prove* the statement: $\lim_{x \rightarrow 2} x^2 + 3 = 7$. First, we do some preliminary analysis. That is, even though to prove this statement, we must show that the definition holds for *any* choice of H and K , we will start out with a specific choice of H and K .

Problem 1. *Let $f(x) = x^2 + 3$, $H = 6$, and $K = 8$. Find values h and k so that for all numbers t between h and k we have $f(t)$ between H and K . Be sure to prove that if $h < t < k$ then $H < f(t) < K$.*

Does this *prove* the theorem? No. We must show that we can solve the previous problem for *any* choice of H and K .

Problem 2. *Let $f(x) = x^2 + 3$. Let $y = H$ and $y = K$ be two horizontal lines with $y = 7$ between them. Find vertical lines $x = h$ and $x = k$ satisfying the definition of the limit.*

Problem 3. *Prove: $\lim_{x \rightarrow 2} x^2 + 3 = 7$ by showing that if t satisfies $h < t < k$ then $H < f(t) < K$.*

This definition can be used to prove some of the theorems about limits that we used. Here are a few that you might enjoy working on.

Problem 4. *Prove that if $a \in \mathbb{R}$ then $\lim_{x \rightarrow a} x = a$.*

Problem 5. Prove that if $a \in \mathbb{R}$ then $\lim_{x \rightarrow a} x^2 + 3 = a^2 + 3$.

Problem 6. Prove that if f and g are functions and $\lim_{x \rightarrow a} f(x) = f(a)$ and $\lim_{x \rightarrow a} g(x) = g(a)$ then $\lim_{x \rightarrow a} (f(x) + g(x)) = f(a) + g(a)$.

15.2 The Extended Reals

Have you ever wondered exactly what *infinity* really is? Is it a number? Can you multiply by it? Can you divide by it? Does $\frac{\infty}{\infty} = 1$? Well if you have wondered, then this appendix is for you!

In this text, we try to define precisely most of the words and symbols that we use. Yet we did not define the real numbers, and to develop them from scratch is a non-trivial task. Thus, the real numbers are an “undefined” term in this text. One philosophy that separates mathematics from other less scientific fields is our constant attempt to make clear the distinction between that which we are assuming (axioms or undefined terms) and that which we have proved (theorems, lemmas, or corollaries). We will see examples in this course of each. We will not define what a “real number” is, but we will now define the “extended reals” assuming that the reals exist. We will give only an intuitive notion of the definition of a “limit” but we will define “derivatives” precisely based on our intuitive understanding of limits.

Where does this leave us, if we build the subject on undefined terms and intuitive definitions? It leaves us in better shape than most subjects! While we have not defined everything, we know what we have defined and what we have proved based on those definitions. Should you find mathematics worthy of further exploration, you can take courses where the real numbers are developed (based on even more elementary axioms and assumptions) and where limits are studied in rigorous detail. Can we go all the way back to the beginning? I will if you will define the word “beginning”. Enough mathematical philosophy, let’s get back to the extended reals.

Definition 1. Assuming the existence of the real numbers, we define the **extended reals** to be the set of all real numbers along with two new numbers which we will call $+\infty$ and $-\infty$.

We already know many properties of the real numbers. We know how to add, subtract, multiply, and divide. But which of these properties follow for the real numbers? Here are a few:

Axiom 1. $\infty + a = \infty$ for all real numbers, a .

Axiom 2. $\infty + \infty = \infty$.

Axiom 3. $\infty \cdot \pm a = \pm \infty$ for all positive real numbers, a .

Axiom 4. $\infty \cdot \infty = \infty$.

Axiom 5. $\infty / \pm a = \pm \infty$ for all positive real numbers, a .

Axiom 6. $a / \infty = 0$ for all real numbers, a .

Our interest with the numbers $\pm \infty$ stems from limits. For example, what is

$$\lim_{x \rightarrow \infty} 3x?$$

Some would say that the limit does not exist since there is no *real* number that $f(x) = 3x$ approaches as x approaches ∞ . Others would say that the function f increases without bound as x increases without bound. The latter is our personal favorite, but in the interest of notational efficiency, and having defined the extended reals, we will write,

$$\lim_{x \rightarrow \infty} 3x = \infty.$$

Note that this follows from Axiom 3 above.

Knowing what is **not** true is at least as important as knowing what **is** true. Here are some cases called **indeterminate** forms where the rules you might expect to be true do not necessarily hold. Consider $f(x) = x - 5$ and $g(x) = x$. Clearly,

$$\lim_{x \rightarrow \infty} f(x) = \infty \quad \text{and} \quad \lim_{x \rightarrow \infty} g(x) = \infty.$$

But what about $\lim_{x \rightarrow \infty} (f(x) - g(x))$?

$$\lim_{x \rightarrow \infty} (f(x) - g(x)) = \lim_{x \rightarrow \infty} (x - 5 - x) = \lim_{x \rightarrow \infty} -5 = -5.$$

Yet, if we consider $h(x) = x - 5$ and $k(x) = x^2$ then

$$\lim_{x \rightarrow \infty} h(x) = \infty \quad \text{and} \quad \lim_{x \rightarrow \infty} k(x) = \infty.$$

But what about $\lim_{x \rightarrow \infty} (h(x) - k(x))$?

$$\lim_{x \rightarrow \infty} (h(x) - k(x)) = \lim_{x \rightarrow \infty} (x - 5 - x^2) = \lim_{x \rightarrow \infty} (-x^2 + x - 5) = -\infty.$$

Thus we say that $\infty - \infty$ is an **indeterminate** form because when we consider two functions both of whose limits tend to ∞ , the limit of their differences is not determined.

Problem 1. Show that $\frac{\infty}{\infty}$ is an indeterminate form by finding functions f, g, h and k so that

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} g(x) = \lim_{x \rightarrow \infty} h(x) = \lim_{x \rightarrow \infty} k(x) = \infty,$$

but

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} \neq \lim_{x \rightarrow \infty} \frac{h(x)}{k(x)}.$$

Problem 2. Show that each of ∞^0 , 0^∞ , and $0 \cdot \infty$ is an indeterminate form.

15.3 Exponential and Logarithmic Functions

The goal of this appendix is to review exponential and logarithmic functions. Even though you should definitely have seen these functions before you made it to Calculus, it is my experience that many students benefit from a review of the basics. Use this as a review and ask if you have any troubles.

Problem 1. Use a limit table to convince yourself that the $\lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n$ exists by substituting in $n = 10, n = 100$, and $n = 1000$.

Definition 1. We define the real number e by $e = \lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n$.

Definition 2. Any function of the form $f(x) = b^x$ where b is any positive real number other than 1 is called an **exponential** function.

Problem 2. Graph $r(x) = 2^x, s(x) = 3^x, u(t) = 2^{-t}$, and $v(t) = 3^{-t}$ all on the same pair of coordinate axes by either plotting points (preferable) or by using your favorite technological weapon. Pay special attention to the domain and range, x - and y -intercepts, and any vertical or horizontal asymptotes.

Definition 3. The function $f(x) = e^x$ is called the **natural exponential function**.

Problem 3. Graph each of the following variants of the exponential function, listing domain, range, intercepts, and asymptotes.

1. $f(x) = e^x$
2. $g(x) = e^{-x}$
3. $h(x) = e^{x-2}$
4. $i(x) = e^x - 3$

Definition 4. We define the inverse of $f(x) = e^x$ by $g(x) = \ln(x)$ and refer to this as the **natural logarithm function**.

Definition 5. We define the inverse of $f(x) = b^x$ by $g(x) = \log_b(x)$ and refer to this as the **logarithm to the base b**.

By definition of inverses, we have these Inverse Properties.

1. $\ln(e^x) = x$ for all $x \in \mathbb{R}$ and
2. $e^{\ln(x)} = x$ for all $x > 0$.

The following theorem relates the natural logarithm function to all other base logarithm functions which means that if you really understand the natural logarithm function then you can always convert the other logarithms to the natural logarithm function.

Theorem 1. Change of Base Theorem. For any $b \in \mathbb{N}$, $\log_b(x) = \frac{\ln(x)}{\ln(b)}$.

Theorem 2. Algebra of Exponential Theorem.

1. $e^x e^y = e^{x+y}$
2. $e^x / e^y = e^{x-y}$
3. $(e^x)^y = e^{xy}$

Each of these laws leads to a corresponding statement about logarithms.

Theorem 3. Algebra of Natural Log Theorem.

1. $\ln(xy) = \ln(x) + \ln(y)$
2. $\ln(x/y) = \ln(x) - \ln(y)$
3. $\ln(x^y) = y\ln(x)$

Problem 4. Prove the Logarithmic Laws using the Exponential Laws and the Inverse Properties.

Problem 5. Evaluate each of the following without using a calculator.

1. If $e^4 = 54.59815\dots$, then $\ln(54.59815\dots) = \underline{\hspace{2cm}}$.
2. If $\ln(24) = 3.1780538\dots$, then $e^{3.1780538} = \underline{\hspace{2cm}}$.
3. $e^{\ln(4)} = \underline{\hspace{2cm}}$.

4. $\ln(e^x) =$ _____.

Problem 6. Evaluate each of the following using a calculator.

1. $\ln(3.41) =$ _____.

2. $e^{4\ln(2)} =$ _____.

Problem 7. Write each of the following as a single logarithm.

1. $3\ln(2x) - 2\ln(x) + \ln(y) - \ln(z)$

2. $2\ln(x+y) + \ln\left(\frac{1}{x-y}\right)$

3. $\ln(x) + 3$ *Hint : $3 = \ln(\text{what})$?*

4. Which of the following is correct? Why?

(a) $\ln(x) - \ln(y) + \ln(z) = \ln\left(\frac{x}{yz}\right)$ or

(b) $\ln(x) - \ln(y) + \ln(z) = \ln\left(\frac{xz}{y}\right)$

Problem 8. Expand each of the following into a sum, difference, or multiple of the logarithms.

1. $\ln(9x^2y)$

2. $\ln(4x^{-1}y^2)$

3. $\ln\left(\frac{xy^2}{z^3}\right)$

4. $\ln\left(\frac{(x+y)^2}{(x-y)^2}\right)$

Problem 9. Solve each of the following for x ; give both exact and approximate answers.

1. $\ln(3x) = 2$

2. $\ln(6) + \ln(2x) = 3$

3. $\ln(x^2 - x - 5) = 0$

4. $\ln(x+3) + \ln(x-2) = 2$

5. $\ln^2 x - \ln x^5 + 4 = 0$ *Notation: $\ln^2 x$ means $\ln(x)^2$ and $\ln x^5$ means $\ln(x^5)$*

6. $\ln^2 x + \ln x^2 - 3 = 0$

Problem 10. Solve for x ; give both exact and approximate answers.

1. $e^x = 19$

2. $4^x = \frac{2}{3}$

3. $4^{x+1} = e$

4. $\left(\frac{1}{2}\right)^x = 3$

5. $x^2e^x - 9e^x = 0$ *Hint: Factor, factor, factor!*

6. $e^x - 8e^{-x} = 2$

7. $5^x = 4^{x+1}$

8. $3^{2x+1} = 2^{x-1}$

Solutions to Exponential and Logarithmic Exercises

Solution 1. *These are the solutions to Problem 5.*

1. 4

2. 24

3. 4

4. x

Solution 2. *These are the solutions to Problem 6.*

1. 1.2267123

2. 16

Solution 3. *These are the solutions to Problem 7.*

1. $\ln\left(\frac{8xy}{z}\right)$

2. $\ln\left(\frac{(x+y)^2}{(x-y)}\right)$

3. $\ln(xe^3)$

4. *Only the second statement is true.*

Solution 4. *These are the solutions to Problem 8.*

1. $\ln(9) + 2\ln(x) + \ln(y)$

2. $\ln(4) + 2\ln(y) - \ln(x)$

3. $\ln(x) + 2\ln(y) - 3\ln(z)$

4. $2\ln(x+y) - 2\ln(x-y)$

Solution 5. *These are the solutions to Problem 9.*

1. $e^2/3$

2. $e^3/12$

3. $-2, 3$

4. $\frac{-1 + \sqrt{25 + 4e^2}}{2}$

5. e, e^4

6. $\frac{1}{e^3}, e$

Solution 6. *These are the solutions to Problem 10.*

1. $\ln 19$

2. $\frac{\ln(\frac{2}{3})}{\ln 4}$

3. $\frac{1}{\ln 4} - 1$

4. $\frac{\ln 3}{\ln \frac{1}{2}}$

5. $-3, 3$

6. $\ln 4$

7. $\frac{\ln 4}{\ln(\frac{5}{4})}$

8. $\frac{\ln \frac{1}{6}}{\ln \frac{9}{2}}$

15.4 Trigonometry

We state here only the very minimal definitions and identities from trigonometry that we expect you to know. By “know” I mean memorize and be prepared to use on a test.

Definition 1. *The **unit circle** is the circle of radius one, centered at the origin.*

Definition 2. *If $P = (x, y)$ is a point on the unit circle and L is the line through the origin and P , and θ is the angle between the x -axis and L , then we define **cos**(θ) to be x and **sin**(θ) to be y .*

Definition 3. *These are the definitions of the remaining four trigonometric functions:*

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}, \quad \cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)}, \quad \csc(\theta) = \frac{1}{\sin(\theta)}, \quad \text{and} \quad \sec(\theta) = \frac{1}{\cos(\theta)}$$

Problem 1. *Graph each of $\sin(-\theta)$, $-\sin(\theta)$, $\cos(-\theta)$ and $\cos(\theta)$ to convince yourself of the following theorem.*

Theorem 1. Even/Odd Identities.

1. $\sin(-\theta) = -\sin(\theta)$

2. $\cos(-\theta) = \cos(\theta)$

The Pythagorean Identity follows immediately from Definition 2.

Theorem 2. The Pythagorean Identity. $\sin^2(\theta) + \cos^2(\theta) = 1$

Axiom 1. *It's easier to memorize one identity than three.*

Problem 2. *Divide both sides of the Pythagorean Identity by $\sin^2(\theta)$ to show that $1 + \tan^2(\theta) = \sec^2(\theta)$. Divide by $\cos^2(\theta)$ to show that $1 + \cot^2(\theta) = \csc^2(\theta)$.*

These next ones are a bit tricky to derive, but easy to remember if you memorize the Double Angle Identities.

Theorem 3. Sum/Difference Identities.

1. $\sin(x \pm y) = \sin(x) \cos(y) \pm \cos(x) \sin(y)$
2. $\cos(x \pm y) = \cos(x) \cos(y) \mp \sin(x) \sin(y)$

Theorem 4. Double Angle Identities.

1. $\sin(2x) = 2 \sin(x) \cos(x)$
2. $\cos(2x) = \cos^2(x) - \sin^2(x)$

Problem 3. Use the first Sum/Difference Identity to prove the first Double Angle Identity.

Problem 4. Use the second Sum/Difference Identity to prove the second Double Angle Identity.

Theorem 5. Half Angle Identities.

1. $\sin^2(x/2) = \frac{1 - \cos(x)}{2}$
2. $\cos^2(x/2) = \frac{1 + \cos(x)}{2}$

Theorem 6. Product Identities.

1. $\sin(mx) \sin(nx) = \frac{1}{2} [\cos(m - n)x - \cos(m + n)x]$
2. $\sin(mx) \cos(nx) = \frac{1}{2} [\sin(m - n)x + \sin(m + n)x]$
3. $\cos(mx) \cos(nx) = \frac{1}{2} [\cos(m - n)x + \cos(m + n)x]$

15.5 Summation Formulas

These formulas are useful for computing Riemann Sums.

1. $\sum_{i=1}^n c = nc$ where c is any real number
2. $\sum_{i=1}^n i = \frac{n(n+1)}{2}$
3. $\sum_{i=1}^n i^3 = \left[\frac{n(n+1)}{2} \right]^2$
4. $\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$
5. $\sum_{i=1}^n i^4 = \frac{n(n+1)(2n+1)(3n^2+3n+1)}{30}$