



Novel Mechanism for Asymmetric Leptoquark Pair Production at Hadron Colliders

Casey Hampson

(Department of Physics, Kennesaw State University)



Dr. André Sopczak (CERN; ATLAS Collaboration)



Dr. Marco Guzzi (Department of Physics, Kennesaw State University)


**91st Annual Meeting of the Southeastern Section
of the American Physics Society**

Oct 24–26, 2024 University of North Carolina
at Charlotte, Charlotte, NC, USA





Introduction

- The Standard Model (SM) accurately describes 99% of the phenomena we observe in the universe.
- As we explore high-energy scales in proton-proton collisions at the CERN Large Hadron Collider (LHC), discrepancies between data and theory may hide signatures of new physics, and we must consider theories that extend the SM.
- One such discrepancy is the ratio of decays of the B-meson into e^-e^+ and $\mu^-\mu^+$ ([LHCb 2406.03387](#), [LHCb 2312.09115](#), [Pacey 2405.11572](#))



The Leptoquark

- Some BSM theories introduce the **Leptoquark (LQ)**, a (spin-0,1) boson that couples directly to both quarks and leptons.
- The LQ can be used to explain the aforementioned discrepancy, as well as generate neutrino masses via quantum corrections. [Babu, Julio, NPB841 \(2010\)](#).
- Our goal for this project was to study a novel method for pair producing LQ's at the LHC



Asymmetric Pair Production

To achieve this novel asymmetric pair production mechanism as proposed in ([Doršner, et. al. JHEP \(2023\)](#), [Doršner, et. al. JHEP \(2021\)](#)) we require:


- Produced via a t-channel lepton exchange
- Two LQ's in the final states are not charge conjugates
- They couple to a lepton of the same chirality and flavor

Two such LQ's that can satisfy this criteria are denoted as the S_1 and R_2

$SU(3)_c \times SU(2)_L \times U(1)_Y$	Symbol	Q-L Chirality
$(\mathbf{3}, \mathbf{2}, 7/6)$	R_2	RL, LR
$(\mathbf{3}, \mathbf{1}, 1/3)$	S_1	LL, RR

Asymmetric Pair Production

- We worked in the 2lSS+1 τ group which specializes in more massive decay products
- Along with the symmetry constraints from the Lagrangian we specify Yukawa parameters that force decays into top quarks and τ 's





Analysis

- We used advanced event generators and parton showering programs like MadGraph5 and Pythia8 for the simulation
- The decays of these heavy particles give a final state that is almost identical to that of other SM processes
- Our goal is to differentiate our final-state signature (the signal) from these others (background)
- We use Machine Learning (ML) on the vast number of output variables produced by the simulations (n-tuples)

Analysis



- I heavily updated the analysis code (previously tailored to top-quark pair in association with Higgs production ($t\bar{t}H$)) to process the LQ's and the improved n-tuple models
- Kept old ML model setups, including the number of layers, epochs, etc., only changed number of input parameters



Results

- We show results of the analysis code on the previous ttH data to compare the performance
- Full LQ results are still a work in progress
- After the model was trained, we used a program called TRExFitter to display some histograms of the model's output
- It also generated some preliminary info on the statistical uncertainty


Results – Signal Separation



Results – Statistical Uncertainties

To obtain a measure of the statistical uncertainties associated with the limited dataset size in relation to our model output we consider the median signal strength parameter μ defined as

$$(\text{background}) + \mu \cdot (\text{signal}) = \text{observed}$$



Conclusions and Future Plans

- We have shown that the analysis code prepared for this novel LQ production method performs very well after preliminary testing
- Inputting the LQ samples are next
- Fine tune model parameters
- Investigating systematics and things like fake-classification which are left out in the newest n-tuple model
- This machinery will be used to explore the parameter space of a variety of BSM models

Backup

References

- [1] LHCb collaboration. *Measurement of the branching fraction ratios $R(D^+)$ and $R(D^{*+})$ using muonic τ decays*. 2024. arXiv: 2406.03387 [hep-ex]. URL: <https://arxiv.org/abs/2406.03387>.
- [2] LHCb Collaboration. “Amplitude Analysis of the B^0 Decay”. In: *Physical Review Letters* 132.13 (Mar. 2024). ISSN: 1079-7114. DOI: 10.1103/physrevlett.132.131801. URL: <http://dx.doi.org/10.1103/PhysRevLett.132.131801>.
- [3] Holly Pacey. *Searches for Lepton Flavour Violation at ATLAS and CMS*. 2024. arXiv: 2405.11572 [hep-ex]. URL: <https://arxiv.org/abs/2405.11572>.
- [4] K.S. Babu and J. Julio. “Two-loop neutrino mass generation through leptoquarks”. In: *Nuclear Physics B* 841.1–2 (Dec. 2010), 130–156. ISSN: 0550-3213. DOI: 10.1016/j.nuclphysb.2010.07.022. URL: <http://dx.doi.org/10.1016/j.nuclphysb.2010.07.022>.
- [5] Ilja Doršner, Ajla Lejlić, and Shaikh Saad. “Asymmetric leptoquark pair production at LHC”. In: *Journal of High Energy Physics* 2023.3 (Mar. 2023). ISSN: 1029-8479. DOI: 10.1007/jhep03(2023)025. URL: [http://dx.doi.org/10.1007/JHEP03\(2023\)025](http://dx.doi.org/10.1007/JHEP03(2023)025).
- [6] Ilja Doršner, Svjetlana Fajfer, and Ajla Lejlić. “Novel leptoquark pair production at LHC”. In: *Journal of High Energy Physics* 2021.5 (May 2021). ISSN: 1029-8479. DOI: 10.1007/jhep05(2021)167. URL: [http://dx.doi.org/10.1007/JHEP05\(2021\)167](http://dx.doi.org/10.1007/JHEP05(2021)167).
- [7] Vladyslav Yazykov. “Optimization of the ttH Selection Including Systematics Using Machine Learning with ATLAS Data”. Presented 05 Sep 2023. Prague, Tech. U., 2023. URL: <https://cds.cern.ch/record/2871484>.