

Balls, Urns, and the Supreme Court

- Supreme Court case: *Berghuis v. Smith*

If a group is underrepresented in a jury pool, how do you tell?

- Article by Erin Miller – Friday, January 22, 2010
- Thanks to (former CS109er) Josh Falk for this article

Justice Breyer [Stanford Alum] opened the questioning by invoking the binomial theorem. He hypothesized a scenario involving **“an urn with a thousand balls, and sixty are red, and nine hundred forty are black, and then you select them at random... twelve at a time.”** According to Justice Breyer and the binomial theorem, if the red balls were black jurors then **“you would expect... something like a third to a half of juries would have at least one black person”** on them.

- Justice Scalia’s rejoinder: “We don’t have any urns here.”

Justice Breyer Meets CS109

- Should model this combinatorially ($X \sim \text{HypGeo}$)
 - Ball draws not independent trials (balls not replaced)
- Exact solution:
$$P(\text{draw 12 black balls}) = \frac{\binom{940}{12}}{\binom{1000}{12}} \approx 0.4739$$
$$P(\text{draw} \geq 1 \text{ red ball}) = 1 - P(\text{draw 12 black balls}) \approx 0.5261$$
- Approximation using Binomial distribution
 - Assume $P(\text{red ball})$ constant for every draw = $60/1000$
 - $X = \#$ red balls drawn. $X \sim \text{Bin}(12, 60/1000 = 0.06)$
 - $P(X \geq 1) = 1 - P(X = 0) \approx 1 - 0.4759 = 0.5240$

In Breyer's description, should actually expect just over half of juries to have at least one black person on them

Demo

From Discrete to Continuous

- So far, all random variables we saw were *discrete*
 - Have finite or countably infinite values (e.g., integers)
 - Usually, values are binary or represent a count
- Now it's time for *continuous* random variables
 - Have (uncountably) infinite values (e.g., real numbers)
 - Usually represent measurements (arbitrary precision)
 - Height (centimeters), Weight (lbs.), Time (seconds), etc.
- Difference between how many and how much
- Generally, it means replace $\sum_{x=a}^b f(x)$ with $\int_a^b f(x)dx$

Continuous Random Variables

- X is a **Continuous Random Variable** if there is function $f(x) \geq 0$ for $-\infty \leq x \leq \infty$, such that:

$$P(a \leq X \leq b) = \int_a^b f(x)dx$$

- f is a Probability Density Function (PDF) if:

$$P(-\infty < X < \infty) = \int_{-\infty}^{\infty} f(x)dx = 1$$

Probability Density Functions

- Say f is a **Probability Density Function** (PDF)

$$P(-\infty < X < \infty) = \int_{-\infty}^{\infty} f(x)dx = 1$$

- $f(x)$ is **not** a probability, it is probability/units of X
- Not meaningful without some subinterval over X

$$P(X = a) = \int_a^a f(x)dx = 0$$

- Contrast with Probability Mass Function (PMF) in discrete case: $p(a) = P(X = a)$

where $\sum_{i=1}^{\infty} p(x_i) = 1$ for X taking on values x_1, x_2, x_3, \dots

Cumulative Distribution Functions

- For a continuous random variable X , the **Cumulative Distribution Function** (CDF) is:

$$F(a) = P(X < a) = P(X \leq a) = \int_{-\infty}^a f(x)dx$$

- Density f is derivative of CDF F : $f(a) = \frac{d}{da} F(a)$
- For continuous f and small ε :

$$P(a - \frac{\varepsilon}{2} \leq X \leq a + \frac{\varepsilon}{2}) = \int_{a-\varepsilon/2}^{a+\varepsilon/2} f(x)dx \approx \varepsilon f(a)$$


- So, ratio of probabilities can still be meaningful:
 - $P(X = 1)/P(X = 2) \approx (\varepsilon f(1))/(\varepsilon f(2)) = f(1)/f(2)$

Simple Example

“Support” of PDF

- X is continuous random variable (CRV) with PDF:

$$f(x) = \begin{cases} C(4x - 2x^2) & \text{when } 0 < x < 2 \\ 0 & \text{otherwise} \end{cases}$$



- What is C ?

$$\int_0^2 C(4x - 2x^2) dx = 1 \Rightarrow C \left(2x^2 - \frac{2x^3}{3} \right) \Big|_0^2 = 1$$

$$C \left(\left(8 - \frac{16}{3} \right) - 0 \right) = 1 \Rightarrow C \frac{8}{3} = 1 \Rightarrow C = \frac{3}{8}$$

- What is $P(X > 1)$?

$$\int_1^{\infty} f(x) dx = \int_1^2 \frac{3}{8} (4x - 2x^2) dx = \frac{3}{8} \left(2x^2 - \frac{2x^3}{3} \right) \Big|_1^2 = \frac{3}{8} \left[\left(8 - \frac{16}{3} \right) - \left(2 - \frac{2}{3} \right) \right] = \frac{1}{2}$$

Disk Crashes

- X = days of use before your disk crashes

$$f(x) = \begin{cases} \lambda e^{-x/100} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

- First, determine λ to have actual PDF

- Good integral to know: $\int e^u du = e^u$

$$1 = \int \lambda e^{-x/100} dx = -100\lambda \int \frac{-1}{100} e^{-x/100} dx = -100\lambda e^{-x/100} \Big|_0^\infty = 100\lambda \Rightarrow \lambda = \frac{1}{100}$$

- What is $P(50 < X < 150)$?

$$F(150) - F(50) = \int_{50}^{150} \frac{1}{100} e^{-x/100} dx = -e^{-x/100} \Big|_{50}^{150} = -e^{-3/2} + e^{-1/2} \approx 0.383$$

- What is $P(X < 10)$?

$$F(10) = \int_0^{10} \frac{1}{100} e^{-x/100} dx = -e^{-x/100} \Big|_0^{10} = -e^{-1/10} + 1 \approx 0.095$$

Expectation and Variance

For discrete RV X :

$$E[X] = \sum_x x p(x)$$

$$E[g(X)] = \sum_x g(x) p(x)$$

$$E[X^n] = \sum_x x^n p(x)$$

For continuous RV X :

$$E[X] = \int_{-\infty}^{\infty} x f(x) dx$$

$$E[g(X)] = \int_{-\infty}^{\infty} g(x) f(x) dx$$

$$E[X^n] = \int_{-\infty}^{\infty} x^n f(x) dx$$

For both discrete and continuous RVs:

$$E[aX + b] = aE[X] + b$$


$$\text{Var}(X) = E[(X - \mu)^2] = E[X^2] - (E[X])^2$$

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Linearly Increasing Density

- X is a continuous random variable with PDF:

$$f(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$



- What is $E[X]$?

$$E[X] = \int_{-\infty}^{\infty} x f(x) dx = \int_0^1 2x^2 dx = \frac{2}{3} x^3 \Big|_0^1 = \frac{2}{3}$$

- What is $\text{Var}(X)$?

$$E[X^2] = \int_{-\infty}^{\infty} x^2 f(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} x^4 \Big|_0^1 = \frac{1}{2}$$


$$\text{Var}(X) = E[X^2] - (E[X])^2 = \frac{1}{2} - \left(\frac{2}{3}\right)^2 = \frac{1}{18}$$

Uniform Random Variable

- X is a **Uniform Random Variable**: $X \sim \text{Uni}(\alpha, \beta)$

- Probability Density Function (PDF):

$$f(x) = \begin{cases} \frac{1}{\beta - \alpha} & \alpha \leq x \leq \beta \\ 0 & \text{otherwise} \end{cases}$$



- Sometimes defined over range $\alpha < x < \beta$

- $P(a \leq x \leq b) = \int_a^b f(x) dx = \frac{b - a}{\beta - \alpha} \quad (\text{for } \alpha \leq a \leq b \leq \beta)$
- $E[X] = \int_{-\infty}^{\infty} x f(x) dx = \int_{\alpha}^{\beta} \frac{x}{\beta - \alpha} dx = \frac{x^2}{2(\beta - \alpha)} \Big|_{\alpha}^{\beta} = \frac{\beta^2 - \alpha^2}{2(\beta - \alpha)} = \frac{\alpha + \beta}{2}$
- $\text{Var}(X) = \frac{(\beta - \alpha)^2}{12}$

Fun with the Uniform Distribution

- $X \sim \text{Uni}(0, 20)$

$$f(x) = \begin{cases} \frac{1}{20} & 0 \leq x \leq 20 \\ 0 & \text{otherwise} \end{cases}$$

- $P(X < 6)$?

$$P(x < 6) = \int_0^6 \frac{1}{20} dx = \frac{6}{20}$$

- $P(4 < X < 17)$?

$$P(4 < x < 17) = \int_4^{17} \frac{1}{20} dx = \frac{17}{20} - \frac{4}{20} = \frac{13}{20}$$

Riding the Marguerite Bus

- Say the Marguerite bus stops at the Gates bldg. at 15 minute intervals (2:00, 2:15, 2:30, etc.)
 - Passenger arrives at stop uniformly between 2-2:30pm
 - $X \sim \text{Uni}(0, 30)$

- $P(\text{Passenger waits} < 5 \text{ minutes for bus})?$
 - Must arrive between 2:10-2:15pm or 2:25-2:30pm

$$P(10 < X < 15) + P(25 < x < 30) = \int_{10}^{15} \frac{1}{30} dx + \int_{25}^{30} \frac{1}{30} dx = \frac{5}{30} + \frac{5}{30} = \frac{1}{3}$$

- $P(\text{Passenger waits} > 14 \text{ minutes for bus})?$
 - Must arrive between 2:00-2:01pm or 2:15-2:16pm

$$P(0 < X < 1) + P(15 < x < 16) = \int_0^1 \frac{1}{30} dx + \int_{15}^{16} \frac{1}{30} dx = \frac{1}{30} + \frac{1}{30} = \frac{1}{15}$$