

Likelihood of Data

- Consider n I.I.D. random variables X_1, X_2, \dots, X_n
 - X_i is a sample from density function $f(X_i | \theta)$
 - Note: now explicitly specify parameter θ of distribution
 - We want to determine how “likely” the observed data (x_1, x_2, \dots, x_n) is based on density $f(X_i | \theta)$
 - Define the **Likelihood function**, $L(\theta)$:

$$L(\theta) = \prod_{i=1}^n f(X_i | \theta)$$

- This is just a product since X_i are I.I.D.
 - Intuitively: what is probability of observed data using density function $f(X_i | \theta)$, for some choice of θ

[Demo](#)

Maximum Likelihood Estimator

- The **Maximum Likelihood Estimator** (MLE) of θ , is the value of θ that maximizes $L(\theta)$
 - More formally: $\theta_{MLE} = \arg \max_{\theta} L(\theta)$
 - More convenient to use **log-likelihood function**, $LL(\theta)$:

$$LL(\theta) = \log L(\theta) = \log \prod_{i=1}^n f(X_i | \theta) = \sum_{i=1}^n \log f(X_i | \theta)$$

- Note that \log function is “monotone” for positive values
 - Formally: $x \leq y \Leftrightarrow \log(x) \leq \log(y)$ for all $x, y > 0$
- So, θ that maximizes $LL(\theta)$ also maximizes $L(\theta)$
 - Formally: $\arg \max_{\theta} LL(\theta) = \arg \max_{\theta} L(\theta)$
 - Similarly, for any positive constant c (not dependent on θ):
$$\arg \max_{\theta} (c \cdot LL(\theta)) = \arg \max_{\theta} LL(\theta) = \arg \max_{\theta} L(\theta)$$

Computing the MLE

- General approach for finding MLE of θ
 - Determine formula for $LL(\theta)$
 - Differentiate $LL(\theta)$ w.r.t. (each) θ : $\frac{\partial LL(\theta)}{\partial \theta}$
 - To maximize, set $\frac{\partial LL(\theta)}{\partial \theta} = 0$
 - Solve resulting (simultaneous) equations to get θ_{MLE}
 - Make sure that derived $\hat{\theta}_{MLE}$ is actually a maximum (and not a minimum or saddle point). E.g., check $LL(\theta_{MLE} \pm \varepsilon) < LL(\theta_{MLE})$
 - This step often ignored in expository derivations
 - So, we'll ignore it here too (and won't require it in this class)

Maximizing Likelihood with Bernoulli

- Consider I.I.D. random variables X_1, X_2, \dots, X_n
 - $X_i \sim \text{Ber}(p)$
 - Probability mass function, $f(X_i | p)$, can be written as:

$$f(X_i | p) = p^{x_i} (1-p)^{1-x_i} \quad \text{where } x_i = 0 \text{ or } 1$$

- Likelihood: $L(\theta) = \prod_{i=1}^n p^{X_i} (1-p)^{1-X_i}$

- Log-likelihood:

$$\begin{aligned} LL(\theta) &= \sum_{i=1}^n \log(p^{X_i} (1-p)^{1-X_i}) = \sum_{i=1}^n [X_i (\log p) + (1-X_i) \log(1-p)] \\ &= Y (\log p) + (n-Y) \log(1-p) \quad \text{where } Y = \sum_{i=1}^n X_i \end{aligned}$$

- Differentiate w.r.t. p , and set to 0:

$$\frac{\partial LL(p)}{\partial p} = Y \frac{1}{p} + (n-Y) \frac{-1}{1-p} = 0 \quad \Rightarrow \quad p_{MLE} = \frac{Y}{n} = \frac{1}{n} \sum_{i=1}^n X_i$$

Maximizing Likelihood with Poisson

- Consider I.I.D. random variables X_1, X_2, \dots, X_n
 - $X_i \sim \text{Poi}(\lambda)$
 - PMF: $f(X_i | \lambda) = \frac{e^{-\lambda} \lambda^{x_i}}{x_i!}$ Likelihood: $L(\theta) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$

- Log-likelihood:

$$\begin{aligned} LL(\theta) &= \sum_{i=1}^n \log\left(\frac{e^{-\lambda} \lambda^{X_i}}{X_i!}\right) = \sum_{i=1}^n [-\lambda \log(e) + X_i \log(\lambda) - \log(X_i!)] \\ &= -n\lambda + \log(\lambda) \sum_{i=1}^n X_i - \sum_{i=1}^n \log(X_i!) \end{aligned}$$

- Differentiate w.r.t. λ , and set to 0:

$$\frac{\partial LL(\lambda)}{\partial \lambda} = -n + \frac{1}{\lambda} \sum_{i=1}^n X_i = 0 \quad \Rightarrow \quad \lambda_{MLE} = \frac{1}{n} \sum_{i=1}^n X_i$$

Maximizing Likelihood with Normal

- Consider I.I.D. random variables X_1, X_2, \dots, X_n

- $X_i \sim N(\mu, \sigma^2)$

- PDF: $f(X_i | \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(X_i - \mu)^2 / (2\sigma^2)}$

- Log-likelihood:

$$LL(\theta) = \sum_{i=1}^n \log\left(\frac{1}{\sqrt{2\pi}\sigma} e^{-(X_i - \mu)^2 / (2\sigma^2)}\right) = \sum_{i=1}^n \left[-\log(\sqrt{2\pi}\sigma) - (X_i - \mu)^2 / (2\sigma^2) \right]$$

- First, differentiate w.r.t. μ , and set to 0:

$$\frac{\partial LL(\mu, \sigma^2)}{\partial \mu} = \sum_{i=1}^n 2(X_i - \mu) / (2\sigma^2) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

- Then, differentiate w.r.t. σ , and set to 0:

$$\frac{\partial LL(\mu, \sigma^2)}{\partial \sigma} = \sum_{i=1}^n -\frac{1}{\sigma} + 2(X_i - \mu)^2 / (2\sigma^3) = -\frac{n}{\sigma} + \sum_{i=1}^n (X_i - \mu)^2 / (\sigma^3) = 0$$

Being Normal, Simultaneously

- Now have two equations, two unknowns:

$$\frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \quad -\frac{n}{\sigma} + \sum_{i=1}^n (X_i - \mu)^2 / (\sigma^3) = 0$$

- First, solve for μ_{MLE} :

$$\frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \Rightarrow \sum_{i=1}^n X_i = n\mu \Rightarrow \mu_{MLE} = \frac{1}{n} \sum_{i=1}^n X_i$$

- Then, solve for σ^2_{MLE} :

$$-\frac{n}{\sigma} + \sum_{i=1}^n (X_i - \mu)^2 / (\sigma^3) = 0 \Rightarrow n\sigma^2 = \sum_{i=1}^n (X_i - \mu)^2$$

$$\sigma^2_{MLE} = \frac{1}{n} \sum_{i=1}^n (X_i - \mu_{MLE})^2$$



- Note: μ_{MLE} unbiased, but σ^2_{MLE} biased (same as MOM)

Maximizing Likelihood with Uniform

- Consider I.I.D. random variables X_1, X_2, \dots, X_n
 - $X_i \sim \text{Uni}(\alpha, \beta)$
 - PDF: $f(X_i | \alpha, \beta) = \begin{cases} \frac{1}{\beta - \alpha} & \alpha \leq x_i \leq \beta \\ 0 & \text{otherwise} \end{cases}$
 - Likelihood: $L(\theta) = \begin{cases} \left(\frac{1}{\beta - \alpha}\right)^n & \alpha \leq x_1, x_2, \dots, x_n \leq \beta \\ 0 & \text{otherwise} \end{cases}$
 - Constraint $\alpha \leq x_1, x_2, \dots, x_n \leq \beta$ makes differentiation tricky
 - Intuition: want interval size $(\beta - \alpha)$ to be as small as possible to maximize likelihood function for each data point
 - But need to make sure all observed data contained in interval
 - If all observed data not in interval, then $L(\theta) = 0$
 - Solution: $\alpha_{MLE} = \min(x_1, \dots, x_n)$ $\beta_{MLE} = \max(x_1, \dots, x_n)$

Understanding MLE with Uniform

- Consider I.I.D. random variables X_1, X_2, \dots, X_n
 - $X_i \sim \text{Uni}(0, 1)$
 - Observe data:
 - 0.15, 0.20, 0.30, 0.40, 0.65, 0.70, 0.75



Once Again, Small Samples = Problems

- How do small samples affect MLE?

- In many cases, $\mu_{MLE} = \frac{1}{n} \sum_{i=1}^n X_i$ = sample mean
 - Unbiased. Not too shabby...
- As seen with Normal, $\sigma_{MLE}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu_{MLE})^2$
 - Biased. Underestimates for small n (e.g., 0 for $n = 1$)
- As seen with Uniform, $\alpha_{MLE} \geq \alpha$ and $\beta_{MLE} \leq \beta$
 - Biased. Problematic for small n (e.g., $\alpha = \beta$ when $n = 1$)
- Small sample phenomena intuitively make sense:
 - Maximum likelihood \Rightarrow best explain data we've seen
 - Does not attempt to generalize to unseen data

Properties of MLE

- Maximum Likelihood Estimators are generally:
 - Consistent: $\lim_{n \rightarrow \infty} P(|\hat{\theta} - \theta| < \varepsilon) = 1$ for $\varepsilon > 0$
 - Potentially biased (though asymptotically less so)
 - Asymptotically optimal
 - Has smallest variance of “good” estimators for large samples
 - Often used in practice where sample size is large relative to parameter space
 - But be careful, there are some very large parameter spaces
 - Joint distributions of several variables can cause problems
 - Parameter space grows exponentially
 - Parameter space for 10 *dependent* binary variables $\approx 2^{10}$


Maximizing Likelihood with Multinomial

- Consider I.I.D. random variables Y_1, Y_2, \dots, Y_n
 - $Y_k \sim \text{Multinomial}(p_1, p_2, \dots, p_m)$, where $\sum_{i=1}^m p_i = 1$
 - X_i = number of trials with outcome i where $\sum_{i=1}^m X_i = n$
 - PDF: $f(X_1, \dots, X_m \mid p_1, \dots, p_m) = \frac{n!}{x_1! x_2! \dots x_m!} p_1^{x_1} p_2^{x_2} \dots p_m^{x_m}$
 - Log-likelihood: $LL(\theta) = \log(n!) - \sum_{i=1}^m \log(X_i!) + \sum_{i=1}^m X_i \log(p_i)$
 - Account for constraint $\sum_{i=1}^m p_i = 1$ when differentiating $LL(\theta)$
 - Use Lagrange multipliers (drop non- p_i terms):

$$A(\theta) = \sum_{i=1}^m X_i \log(p_i) + \lambda \left(\sum_{i=1}^m p_i - 1 \right)$$

Rock on!

Joseph-Louis Lagrange
(1736-1813)



Home on Lagrange

- Want to maximize:

$$A(\theta) = \sum_{i=1}^m X_i \log(p_i) + \lambda \left(\sum_{i=1}^m p_i - 1 \right)$$

- Differentiate w.r.t. each p_i , in turn:

$$\frac{\partial A(\theta)}{\partial p_i} = X_i \frac{1}{p_i} + \lambda = 0 \quad \Rightarrow \quad p_i = \frac{-X_i}{\lambda}$$

- Solve for λ , noting $\sum_{i=1}^m X_i = n$ and $\sum_{i=1}^m p_i = 1$:

$$\sum_{i=1}^m p_i = \sum_{i=1}^m \frac{-X_i}{\lambda} \quad \Rightarrow \quad 1 = \frac{-n}{\lambda} \quad \Rightarrow \quad \lambda = -n$$

- Substitute λ into p_i , yielding: $p_i = \frac{X_i}{n}$
- Intuitive result: probability p_i = proportion of outcome i

When MLE's Attack!

- Consider 6-sided die
 - $X \sim \text{Multinomial}(p_1, p_2, p_3, p_4, p_5, p_6)$
 - Roll $n = 12$ times
 - Result: 3 ones, 2 twos, 0 threes, 3 fours, 1 fives, 3 sixes
 - Consider MLE for p_i :
 - $p_1 = 3/12, p_2 = 2/12, p_3 = 0/12, p_4 = 3/12, p_5 = 1/12, p_6 = 3/12$
 - Based on estimate, infer that you will **never** roll a three
 - Do you really believe that?
 - Frequentist: Need to roll more! Probability = frequency in limit
 - Bayesian: Have prior beliefs of probability, even before any rolls!

Need a Volunteer

So good to see
you again!



Two Envelopes

- I have two envelopes, will allow you to have one
 - One contains \$X, the other contains \$2X
 - Select an envelope
 - Open it!
 - Now, would you like to switch for other envelope?
 - To help you decide, compute $E[\$ \text{ in other envelope}]$
 - Let $Y = \$ \text{ in envelope you selected}$
$$E[\$ \text{ in other envelope}] = \frac{1}{2} \cdot \frac{Y}{2} + \frac{1}{2} \cdot 2Y = \frac{5}{4} Y$$
 - Before opening envelope, think either equally good
 - So, what happened by opening envelope?
 - And does it really make sense to switch?