

Clinical Medicine Dosage & Usage Guide

MediCart Clinical Reference Library | Verified Guidelines

1. Analgesics & Antipyretics

Paracetamol / Acetaminophen (Dolo 650, Crocin, Calpol): Standard adult dose is 500mg to 650mg every 4 to 6 hours as needed. Do not exceed 4000mg (4g) within a 24-hour window to prevent acute liver toxicity. Side effects are rare but include skin rash or liver enzyme elevations when overdosed. Avoid combining with alcohol due to heightened hepatotoxicity hazards.

Ibuprofen (Brufen 400mg): Nonsteroidal anti-inflammatory drug (NSAID). Standard adult dosage is 400mg every 6 to 8 hours, preferably taken after meals to avoid gastrointestinal irritation. Maximum daily dose is 1200mg. Commonly causes stomach pain, heartburn, or nausea. Contraindicated in patients with active peptic ulcers, chronic kidney disease, or severe heart failure.

2. Antibiotics

Azithromycin (Azithral 500mg): Broad-spectrum macrolide antibiotic. Typical course is 500mg once daily for 3 to 5 days. Must be taken 1 hour before or 2 hours after meals for optimal absorption. Common side effects: mild diarrhea, nausea, stomach cramps, or temporary hearing changes at high doses. Ensure full compliance to prevent antibiotic resistance.

3. Gastrointestinal Drugs

Pantoprazole (Pan-40): Proton Pump Inhibitor (PPI) that reduces gastric acid production. Recommended dosage is 40mg once daily, taken 30 to 45 minutes before breakfast. Useful for GERD, acid reflux, and stomach ulcers. Long term usage may lead to vitamin B12 or magnesium deficiency.

4. Antidiabetics

Metformin 500mg (Glycomet): Standard initial dose is 500mg once or twice daily with meals to reduce gastrointestinal discomfort. Helps control blood glucose in Type 2 Diabetes. Major risk includes lactic acidosis (rare but severe), presenting as muscle pain, abdominal distress, and drowsiness.