

1 Results

Definition 1. (Interference) Given two sets U, W , we say U k -interferes with W if

$$|U \cap W| \geq \frac{|W|}{k} \quad (1)$$

for some $k \in (0, |W|]$

Corollary 2. If $|U| = |W|$, then U k -interferes with W if and only if W k -interferes with U .

We restrict the upper range of k to $|W|$ for convenience, as beyond that all values of $\frac{|W|}{k}$ will be less than 1.

Definition 3. ((r,T,k)-Subgraph capacity) Given a vertex set $V = \{v_1, \dots, v_n\}$, the (r, T, k) -subgraph capacity of V is the maximum number of subgraphs of size r subject to the constraint that for any randomly generated subgraph U of size r ,

$$\mathbb{E}[X] \leq T \quad (2)$$

where X is the number of interferences caused by adding U to the existing collection of subgraphs.

Lemma 4. Given a vertex set V with n vertices and two subsets U, W of size r , the probability that U k -interferes with W is

$$\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}}$$

Proof. if $V = \{v_1, \dots, v_n\}$, we can represent a subset U as a vector of length n , u defined by

$$u_i = \begin{cases} 1 & \text{if } v_i \in U \\ 0 & \text{if } v_i \notin U \end{cases}$$

With this representation, U, W intersect at the indices where both vectors u, w have a 1. Let Y be a random variable denoting the number of indices where both u, w have a 1. Then

$$P(Y = y) = \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}} \quad (3)$$

Then the probability that U k -interferes with W is

$$\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}} \quad (4)$$

□

Theorem 5. Given a vertex set V with n vertices, the (r, T, k) -subgraph capacity of V is

$$\left\lfloor \frac{2T}{\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}}} + 1 \right\rfloor$$

Proof. Let $M - 1$ be the number of subgraphs of size r currently generated. Add an arbitrary subgraph U of size r . From lemma 4., we know that the probability of U k -interfering with another subgraph is $\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}}$. Since there are $M - 1$ other subgraphs, the expected number of k -interferences by adding U is

$$\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}} \frac{M - 1}{2} \quad (5)$$

Note that we are dividing by 2 since the probability from lemma 4. is the probability that adding U will cause 2 interferences due to the relationship described in corollary 2. From equation (2), we have

$$\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}} \frac{(M - 1)}{2} \leq T \implies M \leq \frac{2T}{\sum_{y=\lceil \frac{r}{k} \rceil}^r \frac{\binom{r}{y} \binom{n-r}{r-y}}{\binom{n}{r}}} + 1 \quad (6)$$

The (r, T, k) -subgraph capacity of V is the largest integer M that satisfies equation (6). □