


3.1.3 $n?$ if $T_n < -31$
 $\therefore -4n + 14 < -31$
 $\therefore -4n < -45$
 $\div (-4) \therefore n > 11\frac{1}{4}$
 \therefore The 12th term \blacktriangleleft



3.2 The even numbers: 6; 12; 18 ...
 \therefore The 13th even number = $13 \times 6 = 78 \blacktriangleleft$

OR: The 13th even number
 = the 26th term of the pattern
 = 26×3
 = 78

4.1 $P = 4\,500$; $i = \frac{4,25}{100} = 0,0425$; $n = \frac{30}{12} = 2\frac{1}{2}$; $A?$
 $A = P(1+i)^n = 4\,500(1+0,0425)^{2,5} = R4\,993,47 \blacktriangleleft$

4.2.1 The loan amount = $R5\,999 - R600 = R5\,399$
 The accumulated amount, $A = P(1+in)$
 where $P = 5\,399$; $i = 8\% = 0,08$; $n = 1\frac{1}{2}$ years; $A?$
 $\therefore A = 5\,399 \left[1 + (0,08) \left(\frac{3}{2} \right) \right]$
 = R6 046,88

\therefore The monthly amount to be paid = $\frac{6\,046,88}{18}$
 = R335,94 \blacktriangleleft



4.2.2 The amount of interest
 = The total amount paid over the 18 months
 – the loan amount
 = R6 046,88 – R5 399
 = R647,88

4.3 28,35 g is worth \$978,34 = $R978,34 \times 8,79$
 = R8 599,61

\therefore 1 g is worth $\frac{R8\,599,61}{28,35}$
 \therefore 1 kg is worth $R \frac{8\,599,61}{28,35} \times 1\,000 \dots 1\,kg = 1\,000\,g$
 $\approx R303\,337,16 \blacktriangleleft$

5.1.1 $A \cap B \blacktriangleleft$ [OR: A and B \blacktriangleleft]

5.1.2 $A' \blacktriangleleft$ [OR: not A \blacktriangleleft]

5.2 Set B \blacktriangleleft

5.3.1 Of the 40 learners, 7 are left-handed
 $\therefore 40 - 7 = 33$ are right-handed




Of the 18 learners who play soccer,
 4 are left-handed

\therefore 14 learners who play soccer are right-handed
 \therefore The number of learners who are right-handed
 and DON'T play soccer
 = $33 - 14 = 19 \blacktriangleleft$



5.3.2




5.3.3 (a) $n(L \text{ or } S) = 3 + 4 + 14 = 21$

$\therefore P(L \text{ or } S) = \frac{21}{40} \blacktriangleleft$

(b) $n(R \text{ and } S) = 14 \dots$ where R is the set
 of all right-handed
 people
 $\therefore P(R \text{ and } S) = \frac{14}{40}$
 = $\frac{7}{20} \blacktriangleleft$

6.1



$f: y = \frac{3}{x} + 1$

y-intercept ($x = 0$): none

x-intercept ($y = 0$): $\frac{3}{x} + 1 = 0$

$\therefore \frac{3}{x} = -1$

$\therefore x = -3$

$g: y = -2x - 4$

y-intercept ($x = 0$): $y = -4$

x-intercept ($y = 0$): $-2x - 4 = 0$

$\therefore -2x = 4$

$\therefore x = -2$

6.2 Asymptotes: $y = 1$ <
& $x = 0$ (the y-axis) <

6.3 Domain of f : $x \neq 0$; $x \in \mathbb{R}$ <
... $(-\infty; 0) \cup (0; \infty)$

$$\begin{aligned} 6.4 \quad f(x) = g(x) &\Rightarrow \frac{3}{x} + 1 = -2x - 4 \\ &\times x \quad \therefore 3 + x = -2x^2 - 4x \\ &\therefore 2x^2 + 5x + 3 = 0 \\ &\therefore (2x + 3)(x + 1) = 0 \\ &\therefore 2x + 3 = 0 \quad \text{or} \quad x + 1 = 0 \\ &\therefore 2x = -3 \quad \therefore x = -1 \quad \text{<} \\ &\therefore x = -\frac{3}{2} \quad \text{<} \end{aligned}$$

Note: These are the x -coordinates of the points of intersection of f and g :



$(-1\frac{1}{2}; -1)$ & $(-1; -2)$

$$\begin{aligned} 6.5 \quad -1 &\leq g(x) < 3 \\ \therefore -1 &\leq -2x - 4 < 3 \quad \dots g(x) = -2x - 4 \end{aligned}$$

$$\begin{aligned} \text{add 4: } \therefore 3 &\leq -2x < 7 \quad \text{When one divides by a negative number, the direction of the 'inequality' changes.} \\ \div (-2): \therefore -\frac{3}{2} &\geq x > -\frac{7}{2} \quad \dots \\ \therefore -\frac{7}{2} &< x \leq -\frac{3}{2} \quad \dots \text{the inequality has been rewritten with the smaller value on the left} \end{aligned}$$


$$\text{i.e. } -3\frac{1}{2} < x \leq -1\frac{1}{2} \quad \text{<} \quad \left[\text{OR: } \left(-3\frac{1}{2}; -1\frac{1}{2}\right] \text{<} \right]$$



(means excluding ;] means including

$$\begin{aligned} 6.6 \quad k(x) &= 2g(x) = 2(-2x - 4) = -4x - 8 \\ \therefore \text{The equation of } k: y &= -4x - 8 \\ \therefore \text{The y-intercept of } k: (0; -8) &\text{<} \quad \dots \text{substitute } x = 0 \end{aligned}$$

6.7 x -intercept of g : $(-2; 0)$
& x -intercept of h : $(2; 0)$ <
 y -intercept of g : $(0; -4)$
& y -intercept of h : $(0; -4)$ <



Notice: The reflected points have the same y -coordinate, but the x -coordinates are opposite in sign.



7.1 $C(-2; 0)$ < ... symmetrical about the y -axis

7.2 The equation of f : $y = a(x+2)(x-2)$... roots -2 & 2
 $\therefore y = a(x^2 - 4)$

$$\begin{aligned} \text{Subst. } B\left(-3; \frac{5}{2}\right): \therefore \frac{5}{2} &= a[(-3)^2 - 4] \\ \therefore \frac{5}{2} &= a(5) \\ \div 5: \therefore a &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \therefore \text{The equation of } f: y &= \frac{1}{2}(x^2 - 4) \\ \therefore y &= \frac{1}{2}x^2 - 2 \quad \text{<} \end{aligned}$$

7.3 The y -intercept of f is $(0; -2)$
 \therefore The range of f : $y \geq -2$ < [OR: $[-2; \infty)$ <]




Graphs are easier than you thought!

The Answer Series offers excellent material in several subjects for Gr 10 - 12.

See our website www.theanswer.co.za

7.4 The graph of h is obtained by flipping f ... $-f(x)$
then, shifting down 2 units ... -2




\therefore The range of h : $y \leq 0$ <

OR: $(-\infty; 0]$ <

$$\text{OR: } h(x) = -\left(\frac{1}{2}x^2 - 2\right) - 2$$

$$\therefore h(x) = -\frac{1}{2}x^2 + 2 - 2$$

$$\therefore h(x) = -\frac{1}{2}x^2$$



\therefore The range of h : $y \leq 0$ <

7.5 $q = -4$... range $y > -4 \Rightarrow y = -4$ is an asymptote

\therefore Equation of g :
 $y = b^x - 4$; $b > 0$

Substitute $A(2; 0)$:

$$0 = b^2 - 4$$

$$\therefore b^2 = 4$$

$$\therefore b = 2 \quad \dots b \neq -2 \quad \therefore b > 0$$

\therefore Equation of g :

$$y = 2^x - 4 \quad \text{<}$$

\therefore means therefore
 \therefore means because



1.1 The mean,

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n} \quad \dots \quad \frac{\text{total number of scores}}{\text{total number of days}}$$

$$= \frac{929}{19}$$


$$\approx 48,89 \quad \leftarrow$$

1.2 31; 31; 34; 36; 37; 39; 40; 43; 46; 46; 48;

(Q₁) **(Q₂)**
 52; 56; 60; 62; 63; 65; 66; 74

The median **(Q₂)** = 46 ←1.3 The lower quartile **(Q₁)** = 37 ←The upper quartile **(Q₃)** = 62 ←

1.4 Min value = 31 & Max value = 74



Our solutions are set out in such a way as to promote thorough understanding and logic!

We trust that this package will help you grow in confidence as you prepare for your exams. The Answer Series study guides have been the key to exam success for many learners. Visit our website to find appropriate resources for **your** success!

www.theanswer.co.za
2.1 $2\,500 \leq x < 4\,500$

The **sum** of ...
 the **products** of the **frequency**
 and the **mid-value** for each interval

2.2 Estimated mean, \bar{X}

$$= \frac{103 \times 3\,500 + 19 \times 5\,500 + 70 \times 7\,500 + 77 \times 9\,500 \dots}{103 + 19 + 70 + 77 + 85 + 99}$$

The sum of the frequencies

$$\dots + 85 \times 11\,500 + 99 \times 13\,500$$

$$= \frac{4\,035\,500}{453}$$

$$\approx 8\,908,39 \text{ kg} \quad \leftarrow$$

$$\bar{X} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i}$$

2.3 The estimated mean ←

This value is at the centre of the set, whereas the modal class is an extreme situation in relation to the other intervals. ←

$$\begin{aligned} 3.1.1 \quad DE^2 &= (3+3)^2 + (-5-3)^2 \\ &= 36 + 64 \\ &= 100 \\ \therefore DE &= 10 \text{ units} \quad \leftarrow \end{aligned}$$



3.1.2 Gradient of DE,

$$m_{DE} = \frac{-5-3}{3+3} = \frac{-8}{6} = -\frac{4}{3} \quad \leftarrow$$

$$3.1.3 \quad m_{EF} = \frac{k+5}{-1-3} = \frac{k+5}{-4}$$


$$D\hat{E}F = 90^\circ \Rightarrow m_{EF} = +\frac{3}{4} \quad \dots \quad EF \perp DE$$

$$\therefore \frac{k+5}{-4} = \frac{3}{4}$$

$$\begin{aligned} \times (-4) \quad \therefore k+5 &= -3 \\ \therefore k &= -8 \quad \leftarrow \end{aligned}$$

$$3.1.4 \quad M\left(\frac{-3+(-1)}{2}; \frac{3+(-8)}{2}\right)$$

$$\therefore M\left(-2; -\frac{5}{2}\right) \quad \leftarrow$$



3.1.5

DEFG will be a \parallel^m if M is the midpoint of EG too.

& Since $D\hat{E}F = 90^\circ$,

DEFG will be a rectangle.

... if one \angle of a \parallel^m is a right \angle ,
 then the \parallel^m is a rectangle.



$$\begin{aligned} \frac{x_G+3}{2} &= -2 \quad \text{and} \quad \frac{y_G+(-5)}{2} = -\frac{5}{2} \\ \times 2) \quad \therefore x_G+3 &= -4 \quad \therefore y_G-5 = -5 \\ \therefore x_G &= -7 \quad \therefore y_G = 0 \\ \therefore G(-7; 0) &\quad \leftarrow \end{aligned}$$

OR: The translation F to G equals that of E to D

$$\begin{aligned} \therefore G(-1-6; -8+8) \\ \therefore G(-7; 0) \quad \leftarrow \end{aligned}$$

OR: The translation D to G equals that of E to F

$$\begin{aligned} \therefore G(-3-4; 3-3) \\ \therefore G(-7; 0) \quad \leftarrow \end{aligned}$$



3.2 $CD^2 = (x-1)^2 + (5+2)^2 = (\sqrt{53})^2$
 $\therefore (x-1)^2 + 49 = 53$
 $\therefore (x-1)^2 = 4$
 $\therefore x-1 = \pm 2$
 $\therefore x = 3 \text{ or } -1$


Note: x must be negative.

But $x < 0$ in the second quadrant

$\therefore x = -1 \leftarrow \dots$ only the neg. value of x is valid

4.1.1 $\sin C = \frac{AB}{AC} \leftarrow$

4.1.2 $\cot A = \frac{AB}{BC}$

Note: $\tan A = \frac{BC}{AB}$; $\cot A = \frac{1}{\tan A}$ 

4.2 The expression


$$= \frac{\frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{3}}}{\sqrt{2}}$$

$$= \frac{1}{2} \times \frac{1}{\sqrt{2}}$$

$$= \frac{1}{2\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \dots \text{The denominator must be rationalised}$$

$$= \frac{\sqrt{2}}{4} \leftarrow \dots \sqrt{2} \times \sqrt{2} = 2$$

4.3.1 $OP = 13$ units $\dots 5 : 12 : 13 \Delta$; Pythagoras



$$\therefore \cos \theta = \frac{-5}{13} = -\frac{5}{13} \leftarrow \dots \cos \theta = \frac{x}{r}$$

4.3.2 $\sin \theta = \frac{12}{13} \rightarrow \operatorname{cosec} \theta = \frac{13}{12}$
 $\therefore \operatorname{cosec}^2 \theta + 1 = \left(\frac{13}{12}\right)^2 + 1 = \frac{169}{144} + 1$
 $= \frac{169 + 144}{144} = \frac{313}{144} \leftarrow \left(= 2\frac{25}{144} \leftarrow\right)$

5.1.1 $5 \cos x = 3$
 $\div 5) \therefore \cos x = \frac{3}{5} (= 0,6)$

$$\therefore x \approx 53,1^\circ \leftarrow \dots \cos^{-1}\left(\frac{3}{5}\right) =$$

5.1.2 $\tan 2x = 1,19$
 $\therefore 2x = 49,958\dots^\circ \dots \tan^{-1} 1,19 =$
 $\div 2) \therefore x \approx 25,0^\circ \leftarrow$

5.1.3 $4 \sec x - 3 = 5$
 $+ 3) \therefore 4 \sec x = 8$
 $\div 4) \therefore \sec x = 2$
 $\therefore \cos x = \frac{1}{2}$
 $x = 60^\circ \leftarrow \dots \cos^{-1}\left(\frac{1}{2}\right) =$

5.2.1 $\hat{JKD} = 8^\circ \leftarrow \dots$ alternate \angle 's; \parallel lines

5.2.2 In $\triangle JDK$: $\frac{DK}{5} = \cot 8^\circ \dots = \frac{1}{\tan 8^\circ}$

$$\times 5) \therefore DK = \frac{5}{\tan 8^\circ}$$

$$= 35,5768\dots \text{ km}$$

$$= 35\,576,8 \text{ metres}$$

$$\approx 35\,577 \text{ metres} \leftarrow$$


\dots correct to the nearest metre



5.2.3 $DS = DK - SK$
 $= 35,58 \text{ km} - 8 \text{ km}$
 $= 27,58 \text{ km} \leftarrow$

5.2.4 $\tan \hat{JSD} = \frac{5}{27,58}$
 $\therefore \hat{JSD} \approx 10,3^\circ \leftarrow \dots \tan^{-1}\left(\frac{5}{27,58}\right) =$
correct to 1 dec. place

6.1.1




6.1.2 $y = -2 \tan x \leftarrow$


6.2.1 $a = 4 \leftarrow$ $g(x) = a \sin x \rightarrow g(90^\circ) = a \sin 90^\circ$
 $\rightarrow 4 = a$

6.2.2 The range of h :
 $-2 \leq y \leq 6 \leftarrow \dots$ the values of y

7.




$$\begin{aligned} 7.1.1 \quad AH^2 &= 0,8^2 + 1,5^2 \\ &= 2,89 \\ \therefore AH &\approx 1,7 \text{ m} \end{aligned}$$



OR: Pythag. triple: 8 : 15 : 17
 $\Rightarrow 0,8 : 1,5 : 1,7$

$$\begin{aligned} 7.1.2 \quad \text{Surface area of roof} &= 4 \times \text{area of } \triangle ABC \\ &= 4 \times \frac{1}{2} (3)(1,7) \\ &= 10,2 \text{ m}^2 \end{aligned}$$



$$\begin{aligned} 7.1.3 \quad \text{Surface area of the walls} &= 4 \times \text{area of GFEC} \\ &= 4 \times (3)(2,1) \\ &= 25,2 \text{ m}^2 \end{aligned}$$

$$\begin{aligned} \therefore \text{The total surface area of the tent} &= 10,2 + 25,2 \\ &= 35,4 \text{ m}^2 \end{aligned}$$



$$7.2.1 \quad \text{Volume} = \frac{4}{3} \pi r^3 = \frac{4}{3} \pi (8)^3 \approx 2\,144,66 \text{ mm}^3$$

$$\begin{aligned} 7.2.2 \quad &2^3 : 1 \\ &= 8 : 1 \end{aligned}$$

$$\frac{\text{New volume}}{\text{Original volume}} = \frac{\frac{4}{3} \pi (2r)^3}{\frac{4}{3} \pi r^3} = \frac{2^3 r^3}{r^3} = \frac{8}{1}$$

$$\begin{aligned} 7.2.3 \quad \text{Volume of silver} &= \frac{4}{3} \pi (8 + 1)^3 - \frac{4}{3} \pi (8)^3 \dots \text{The volume of silver covering the ball} \\ &= 908,967 \dots \\ &\approx 908,97 \text{ mm}^3 \end{aligned}$$



Geometry is easier than you thought!
 The Answer Series offers excellent material (especially Geometry) for Gr 10 - 12.
 See our website www.theanswer.co.za

$$8.1 \quad OQ = 2 \text{ cm} \dots \text{the longer diagonal of a kite bisects the shorter diagonal}$$


$$8.2 \quad \angle POQ = 90^\circ \dots \text{the diagonals of a kite intersect at right angles}$$

$$\begin{aligned} 8.3 \quad \angle QPO &= 20^\circ \dots \text{the longer diagonal of a kite bisects the (opposite) angles of a kite} \\ \therefore \angle QPS &= 40^\circ \end{aligned}$$

9.

Hint:

Use hiliters to mark the various \parallel^m s and Δ^s



The hilitied Δ^s (and their sides) refer to Question 9.3.

$$9.1 \quad \text{In } \triangle DBA: \quad O \text{ is the midpt of } BD \dots \text{diagonals of } \parallel^m BCDE \text{ bisect each other}$$

$$\& F \text{ is the midpt of } AD \dots \text{diagonals of } \parallel^m AODE \text{ bisect each other}$$

$$\therefore OF \parallel AB \dots \text{the line joining the midpoints of two sides of a } \Delta \text{ is } \parallel \text{ to the 3}^{rd} \text{ side}$$

$$9.2 \quad \begin{aligned} AE &\parallel OD \dots \text{opp. sides of } \parallel^m AODE \\ \therefore AE &\parallel BO \end{aligned}$$

$$\text{and } OF \parallel AB \dots \text{proven above}$$

$$\therefore OE \parallel AB$$

$$\therefore ABOE \text{ is a } \parallel^m \dots \text{both pairs of opposite sides are parallel}$$

$$\text{OR: In } \parallel^m AODE: AE = \text{ and } \parallel OD \dots \text{opp. sides of } \parallel^m$$

$$\text{But } OD = \text{ and } \parallel BO \dots O \text{ proved midpt of } BD \text{ in } 9.1$$

$$\therefore AE = \text{ and } \parallel BO$$

$$\therefore ABOE \text{ is a } \parallel^m \dots 1 \text{ pr of opp. sides} = \text{ and } \parallel$$

$$9.3 \quad \text{In } \Delta^s ABO \text{ and } EOD$$

$$1) AB = EO \dots \text{opposite sides of } \parallel^m ABOE$$

$$2) BO = OD \dots \text{proved in } 9.1$$

$$3) AO = ED \dots \text{opposite sides of } \parallel^m AODE$$

$$\therefore \triangle ABO \equiv \triangle EOD \dots \text{SSS}$$

