

第七章 重積分

7.1 二重積分

概念與基本性質

二重積分 \approx (帶符號) 體積: z 軸上方為正, 下方為負.

定義. 給定 $\Omega = [a, b] \times [c, d]$, $b \geq a$, $d \geq c$, $f: \Omega \rightarrow \mathbb{R}$.

- Ω 分割 $\mathbb{P}: a = x_0 < x_1 < x_2 < \cdots < x_n = b$, $c = y_0 < y_1 < y_2 < \cdots < y_m = d$
- $\Delta x_k = x_k - x_{k-1}$, $\Delta y_l = y_l - y_{l-1}$, $k = 1, 2, \dots, n$; $l = 1, 2, \dots, m$
- $\|\mathbb{P}\| = \max \{\Delta x_k \Delta y_l \mid 1 \leq k \leq n, 1 \leq l \leq m\}$
- 樣本點 $(\xi_k, \zeta_l): x_{k-1} \leq \xi_k \leq x_k$, $y_{l-1} \leq \zeta_l \leq y_l$, $k = 1, 2, \dots, n$; $l = 1, 2, \dots, m$.
- $u_{k,l} = \max \{f(x, y) \mid x_{k-1} \leq x \leq x_k, y_{l-1} \leq y \leq y_l\}$, $\ell_{k,l} = \min \{f(x, y) \mid x_{k-1} \leq x \leq x_k, y_{l-1} \leq y \leq y_l\}$, $k = 1, 2, \dots, n$; $l = 1, 2, \dots, m$.
- $R(f, \mathbb{P}) = \sum_{l=1}^m \sum_{k=1}^n f(\xi_k, \zeta_l) \Delta x_k \Delta y_l$, $U(f, \mathbb{P}) = \sum_{l=1}^m \sum_{k=1}^n u_{k,l} \Delta x_k \Delta y_l$, $L(f, \mathbb{P}) = \sum_{l=1}^m \sum_{k=1}^n \ell_{k,l} \Delta x_k \Delta y_l$;
顯然 $L(f, \mathbb{P}) \leq R(f, \mathbb{P}) \leq U(f, \mathbb{P})$.
- 求 $\lim_{\|\mathbb{P}\| \rightarrow 0} R(f, \mathbb{P})$. 若對不同分割與樣本點選取此極限均存在且相等, 稱 f 在 Ω 可積 (分); $f(x, y)$ 在 Ω 的定積分為 $\int_{\Omega} f(x, y) dA \equiv \lim_{\|\mathbb{P}\| \rightarrow 0} R(f, \mathbb{P})$.

性質.

- 若 f 在 Ω 有界, 且除了 Ω 中有限個平滑曲線外 f 在 Ω 連續, 則 f 在 Ω 可積分.
- 若 $\Omega = \Omega_1 \cup \Omega_2$ 且 Ω_1, Ω_2 均為矩形, f 在 Ω_1, Ω_2 均可積, 則 $\int_{\Omega} f dA = \int_{\Omega_1} f dA + \int_{\Omega_2} f dA$.
- 若 f_1, f_2 均在 Ω 可積且 $f_1(x, y) \leq f_2(x, y) \forall (x, y) \in \Omega$, 則 $\int_{\Omega} f_1 dA \leq \int_{\Omega} f_2 dA$.
- 若 $\alpha, \beta \in \mathbb{R}$, f_1, f_2 均在 Ω 可積, 則 $\int_{\Omega} (\alpha f_1 + \beta f_2) dA = \alpha \int_{\Omega} f_1 dA + \beta \int_{\Omega} f_2 dA$.

逐次積分

矩形積分區域

定理 (Fubini). 若 $\Omega = [a, b] \times [c, d]$, $f: \Omega \rightarrow \mathbb{R}$ 為可積, 則

$$\int_{\Omega} f(x, y) dA = \int_c^d \left\{ \int_a^b f(x, y) dx \right\} dy = \int_a^b \left\{ \int_c^d f(x, y) dy \right\} dx$$

例. 令 $\Omega = \{(x, y) \mid -1 \leq x \leq 1, -2 \leq y \leq 2\}$, 則 $\int_{\Omega} \sqrt{1-x^2} dA = \int_{-2}^2 \left\{ \int_{-1}^1 \sqrt{1-x^2} dx \right\} dy = \frac{\pi}{2} \times 4 = 2\pi$.

例. 若 $\Omega = [0, 2] \times [1, 3]$, 求 $\int_{\Omega} x^2 y dA$.

解.

- 由 Fubini 定理 $\int_{\Omega} x^2 y dA = \int_0^2 \int_1^3 x^2 y dy dx = \int_1^3 \int_0^2 x^2 y dx dy$.

- $\int_0^2 \int_1^3 x^2 y \, dy \, dx = \int_0^2 x^2 \left(\frac{y^2}{2} \Big|_1^3 \right) dx = \int_0^2 x^2 \left(\frac{9}{2} - \frac{1}{2} \right) dx = \frac{4}{3} x^3 \Big|_0^2 = \frac{32}{3}.$
- $\int_1^3 \int_0^2 x^2 y \, dx \, dy = \int_1^3 y \left(\frac{x^3}{3} \Big|_0^2 \right) dy = \int_1^3 y \frac{8}{3} dy = \frac{4}{3} y^2 \Big|_1^3 = \frac{36}{3} - \frac{4}{3} = \frac{32}{3}.$

例. 若 $\Omega = [0, 1] \times [0, 3]$, 求 $\int_{\Omega} e^{2x+y} \, dA$.

解.

- 由 Fubini 定理 $\int_{\Omega} e^{2x+y} \, dA = \int_0^3 \int_0^1 e^{2x+y} \, dx \, dy = \int_0^3 \int_0^1 e^{2x+y} \, dy \, dx.$
- $\int_0^3 \int_0^1 e^{2x+y} \, dx \, dy = \int_0^3 e^y \left(\int_0^1 e^{2x} \, dx \right) dy = \int_0^3 e^y \left(\frac{e^{2x}}{2} \Big|_0^1 \right) dy = \frac{(e^2 - 1)(e^3 - 1)}{2}.$
- $\int_0^1 \int_0^3 e^{2x+y} \, dy \, dx = \int_0^1 e^{2x} \left(\int_0^3 e^y \, dy \right) dx = \int_0^1 e^{2x} \left(\frac{e^y}{1} \Big|_0^3 \right) dx = \frac{(e^2 - 1)(e^3 - 1)}{2}.$

例. 若 $\Omega = [0, \pi] \times [0, 2\pi]$, 求 $\int_{\Omega} \sin(x+y) \, dA$.

解.

- 由 Fubini 定理 $\int_{\Omega} \sin(x+y) \, dA = \int_0^{\pi} \int_0^{2\pi} \sin(x+y) \, dy \, dx = \int_0^{\pi} \int_0^{2\pi} \sin(x+y) \, dx \, dy.$
- $\int_0^{\pi} \int_0^{2\pi} \sin(x+y) \, dy \, dx = \int_0^{\pi} \left(-\cos(x+y) \Big|_0^{2\pi} \right) dx = -\int_0^{\pi} (\cos(x+2\pi) - \cos x) \, dx = 0.$
- $\int_0^{2\pi} \int_0^{\pi} \sin(x+y) \, dx \, dy = \int_0^{2\pi} \left(-\cos(x+y) \Big|_0^{\pi} \right) dy = -\int_0^{2\pi} (\cos(\pi+y) - \cos y) \, dy = -\int_0^{2\pi} (\cos \pi \cos y - \sin \pi \sin y - \cos y) \, dy = 2 \int_0^{2\pi} \cos y \, dy = 0.$

例. 若 $\Omega = [1, 2] \times [0, 1]$, 求 $\int_{\Omega} ye^{xy} \, dA$.

解.

- 由 Fubini 定理 $\int_{\Omega} ye^{xy} \, dA = \int_0^1 \int_1^2 ye^{xy} \, dx \, dy = \int_1^2 \int_0^1 ye^{xy} \, dy \, dx.$
- $\int_0^1 \int_1^2 ye^{xy} \, dx \, dy = \int_0^1 y \left(\frac{e^{xy}}{y} \Big|_1^2 \right) dy = \int_0^1 (e^{2y} - e^y) \, dy = \frac{e^2}{2} - e + \frac{1}{2}.$
- $\int_1^2 \int_0^1 ye^{xy} \, dy \, dx = \int_1^2 \left(y \frac{e^{xy}}{x} - \frac{e^{xy}}{x^2} \Big|_0^1 \right) dx = \int_1^2 \left(\underbrace{\frac{e^x}{x} - \frac{e^x}{x^2}}_{\left(\frac{e^x}{x}\right)' = \frac{e^x}{x} - \frac{e^x}{x^2}} + \frac{1}{x^2} \right) dx = \left(\frac{e^x}{x} - \frac{1}{x} \right) \Big|_1^2 = \frac{e^2}{2} - e + \frac{1}{2}.$

習題. 求下列重積分.

- $\int_2^3 \int_1^5 (x+2y) \, dx \, dy$
- $\int_0^{\ln 4} \int_0^{\ln 3} e^{x+y} \, dx \, dy$
- $\int_0^1 \int_0^1 \frac{xy}{\sqrt{x^2+y^2+1}} \, dy \, dx$
- $\int_0^{\frac{\pi}{2}} \int_0^{\cos y} e^x \sin y \, dx \, dy$
- $\int_0^{\frac{\pi}{2}} \int_0^{\cos \theta} \rho^2 \sin \theta \, d\rho \, d\theta$
- $\int_0^1 \int_{y^4}^{y^2} \sqrt{\frac{y}{x}} \, dx \, dy$
- $\int_0^{\frac{\pi}{2}} \int_0^a \frac{r}{\sqrt{a^2 - r^2 \cos^2 \theta}} \, dr \, d\theta$


解.

- $\int_2^3 \int_1^5 (x+2y) dx dy = \int_2^3 \left(\frac{x^2}{2} + 2yx \right) \Big|_1^5 dy = \int_2^3 (12+8y) dy = (12y+4y^2) \Big|_2^3 = 32$
- $\int_0^{\ln 4} \int_0^{\ln 3} e^{x+y} dx dy = \left(\int_0^{\ln 4} e^y dy \right) \cdot \left(\int_0^{\ln 3} e^x dx \right) = (4-1)(3-1) = 6$
- $\int_0^1 \int_0^1 \frac{xy}{\sqrt{x^2+y^2+1}} dy dx = \int_0^1 x \left(\int_0^1 \frac{y}{\sqrt{x^2+y^2+1}} dy \right) dx = \int_0^1 x \left(\frac{1}{2} \int_{x^2+1}^{x^2+2} \frac{1}{\sqrt{u}} du \right) dx$
 $\stackrel{\text{令 } u=y^2+x^2+1, du=2ydy}{=} \int_0^1 x(\sqrt{x^2+2} - \sqrt{x^2+1}) dx = \int_0^1 \underbrace{x\sqrt{x^2+2}}_{\text{令 } u=x^2+2} dx - \int_0^1 \underbrace{x\sqrt{x^2+1}}_{\text{令 } u=x^2+1} dx = \sqrt{3} - \frac{4}{3}\sqrt{2} + \frac{1}{3}$
- $\int_0^{\frac{\pi}{2}} \int_0^{\cos y} e^x \sin y dx dy = \int_0^{\frac{\pi}{2}} \sin y \left(\int_0^{\cos y} e^x dx \right) dy = \int_0^{\frac{\pi}{2}} \sin y (e^{\cos y} - 1) dy = (-e^{\cos y} + \cos y) \Big|_0^{\frac{\pi}{2}} = e - 2$
- $\int_0^{\frac{\pi}{2}} \int_0^{\cos \theta} \rho^2 \sin \theta d\rho d\theta = \int_0^{\frac{\pi}{2}} \sin \theta \left(\int_0^{\cos \theta} \rho^2 d\rho \right) d\theta = \int_0^{\frac{\pi}{2}} \sin \theta \frac{\cos^3 \theta}{3} d\theta = -\frac{\cos^4 \theta}{12} \Big|_0^{\frac{\pi}{2}} = \frac{1}{12}$
- $\int_0^1 \int_{x^2}^{x^3} (x^2+y^2) dy dx = \int_0^1 \left(x^2 y + \frac{y^3}{3} \right) \Big|_{y=x^2}^{y=x^3} dx = \int_0^1 \left(x^2 \cdot x^3 + \frac{x^9}{3} - x^2 \cdot x^2 - \frac{x^6}{3} \right) dx = -\frac{1}{21}$
- $\int_0^y x\sqrt{y^2-x^2} dx$ 中, 令 $u = y^2 - x^2$, 則 $du = -2x dx \Rightarrow x dx = -\frac{1}{2} du$. 積分範圍 x 由 0 至 y , 則變數變換 $u = y^2 - x^2$ 由 $y^2 - 0^2 = y^2$ 至 $y^2 - y^2 = 0$. 則 $\int_0^y x\sqrt{y^2-x^2} dx = \int_{y^2}^0 \sqrt{u} \frac{-1}{2} du = \frac{1}{2} \int_0^{y^2} \sqrt{u} du = \frac{1}{2} \cdot \frac{2}{3} u^{\frac{3}{2}} \Big|_0^{y^2} = \frac{y^3}{3}$. 故 $\int_1^2 \int_0^y x\sqrt{y^2-x^2} dx dy = \int_1^2 \frac{y^3}{3} dy = \frac{y^4}{12} \Big|_1^2 = \frac{5}{4}$.
- $\int_0^1 \int_{y^4}^{y^2} \sqrt{\frac{y}{x}} dx dy = \int_0^1 \sqrt{y} \left(\int_{y^4}^{y^2} \frac{1}{\sqrt{x}} dx \right) dy = \int_0^1 \sqrt{y} (2\sqrt{x}) \Big|_{y^4}^{y^2} dy = 2 \int_0^1 (y^{\frac{3}{2}} - y^{\frac{5}{2}}) dy = \frac{8}{35}$
- $\int_0^a \frac{r}{\sqrt{a^2-r^2 \cos^2 \theta}} dr$ 中, 令 $u = a^2 - r^2 \cos^2 \theta$, 則 $du = -2 \cos^2 \theta r dr \Rightarrow r dr = \frac{-1}{2 \cos^2 \theta} du$. 積分範圍 r 由 0 至 a , 則變數變換 $u = a^2 - r^2 \cos^2 \theta$ 由 $a^2 - 0^2 \cos^2 \theta = a^2$ 至 $a^2 - a^2 \cos^2 \theta = a^2 \sin^2 \theta$. 則 $\int_0^a \frac{r}{\sqrt{a^2-r^2 \cos^2 \theta}} dr = \int_{a^2}^{a^2 \sin^2 \theta} \frac{1}{\sqrt{u}} \frac{-1}{2 \cos^2 \theta} du = \int_{a^2 \sin^2 \theta}^{a^2} \frac{1}{2 \cos^2 \theta \sqrt{u}} du = \frac{\sqrt{u}}{\cos^2 \theta} \Big|_{u=a^2 \sin^2 \theta}^{u=a^2} = a \frac{1 - \sin \theta}{\cos^2 \theta}$.
故 $\int_0^{\frac{\pi}{2}} \int_0^a \frac{r}{\sqrt{a^2-r^2 \cos^2 \theta}} dr d\theta = a \int_0^{\frac{\pi}{2}} \frac{1 - \sin \theta}{\cos^2 \theta} d\theta = a \frac{\sin \theta - 1}{\cos \theta} \Big|_0^{\frac{\pi}{2}} = a \left(\lim_{\theta \rightarrow \frac{\pi}{2}} \frac{\sin \theta - 1}{\cos \theta} + 1 \right) = a$.


一般積分區域

結論 (基本區域積分).


- 選擇初始積分方向 v : 延 x 方向為 dx , 延 y 方向為 dy .
- 想像積分方向 v 為一射線, 進入區域之邊界函數為積分下界, 離開區域之邊界函數為積分上界; 邊界函數本身不含變數 v .
- 次第區域為原始區域延 v 之投影 (令 $v = (\text{常數})$ 之方程式, 或邊界的交集); 若此投影區域維度 > 1 , 選擇積分方向.



$$\int_{\Omega} f(x, y) \, dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) \, dy \, dx$$




$$\int_{\Omega} f(x, y) \, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) \, dx \, dy$$




例. 求 $\int_{\Omega} x \cos y \, dA$, Ω 為 $y=0$, $x=1$, 與 $y=x^2$ 圍成之區域.

$$\begin{aligned} \text{解. } \int_{\Omega} x \cos y \, dA &= \int_0^1 \int_0^{x^2} x \cos y \, dy \, dx = \int_0^1 x \left(\sin y \Big|_0^{x^2} \right) dx \\ &= \int_0^1 x \sin x^2 \, dx = \frac{1 - \cos 1}{2} \end{aligned}$$



例. 求 $\int_{\Omega} e^{-y^2} \, dA$, $\Omega = \{(x, y) \mid 0 \leq y \leq 1, 0 \leq x \leq y\}$.


$$\text{解. } \int_{\Omega} e^{-y^2} \, dA = \int_0^1 \int_0^y e^{-y^2} \, dx \, dy = \int_0^1 y e^{-y^2} \, dy = \frac{1 - e^{-1}}{2}$$



例. 求 $\int_0^1 \int_0^1 e^{\max\{x^2, y^2\}} \, dx \, dy$.

$$\begin{aligned} \text{解. } \int_0^1 \int_0^1 e^{\max\{x^2, y^2\}} \, dx \, dy &= \int_{\Omega_1} e^{y^2} \, dA + \int_{\Omega_2} e^{x^2} \, dA \\ &= \int_0^1 \int_0^y e^{y^2} \, dx \, dy + \int_0^1 \int_0^x e^{x^2} \, dy \, dx = \int_0^1 y e^{y^2} \, dy + \int_0^1 x e^{x^2} \, dx = e - 1. \end{aligned}$$

積分順序交換



例. 求 $\int_0^1 \int_0^{\sqrt{1-x^2}} \sqrt{1-y^2} \, dy \, dx$.

解. $\int_0^1 \int_0^{\sqrt{1-x^2}} \sqrt{1-y^2} \, dy \, dx = \int_0^1 \int_0^{\sqrt{1-y^2}} \sqrt{1-y^2} \, dx \, dy = \int_0^1 (1-y^2) \, dy = \frac{2}{3}$.
 $\int_0^1 \int_0^{\sqrt{1-x^2}} \sqrt{1-y^2} \, dy \, dx = \frac{1}{2} \int_0^1 (\sin^{-1} \sqrt{1-x^2} + x \sqrt{1-x^2}) \, dx = \frac{1}{2} \left(1 + \frac{1}{3}\right) = \frac{2}{3}$.

例. 將下列積分以不同積分順序寫出.

1. $\int_0^2 \int_{x^2}^{2x} f(x, y) \, dy \, dx$

3. $\int_{-1}^2 \int_{x^2}^{x+2} f(x, y) \, dy \, dx$

2. $\int_0^1 \int_{\sin^{-1} y}^{\frac{\pi}{2}} f(x, y) \, dx \, dy$

4. $\int_0^1 \int_0^{2y} f(x, y) \, dx \, dy + \int_1^3 \int_0^{3-y} f(x, y) \, dx \, dy$

解.

1. $\int_0^2 \int_{x^2}^{2x} f(x, y) \, dy \, dx = \int_0^4 \int_{\sqrt{y}}^{\frac{y}{2}} f(x, y) \, dx \, dy$

2. $\int_0^1 \int_{\sin^{-1} y}^{\frac{\pi}{2}} f(x, y) \, dx \, dy = \int_0^{\frac{\pi}{2}} \int_0^{\sin x} f(x, y) \, dx \, dy$

3. $\int_{-1}^2 \int_{x^2}^{x+2} f(x, y) \, dy \, dx = \int_0^1 \int_{-\sqrt{y}}^{\sqrt{y}} f(x, y) \, dx \, dy + \int_1^4 \int_{y-2}^{\sqrt{y}} f(x, y) \, dx \, dy$

4. $\int_0^1 \int_0^{2y} f(x, y) \, dx \, dy + \int_1^3 \int_0^{3-y} f(x, y) \, dx \, dy = \int_0^2 \int_{\frac{x}{2}}^{3-x} f(x, y) \, dy \, dx$

例. 證明 $\int_0^x \int_0^t F(u) \, du \, dt = \int_0^x (x-u)F(u) \, du$.

解. $\int_0^x \int_0^t F(u) \, du \, dt = \int_0^x \int_u^x F(u) \, dt \, du = \int_0^x (x-u)F(u) \, du$.

例. 求 $\int_0^1 \int_x^1 e^{-y^2} \, dy \, dx$.

解. $\int_0^1 \int_x^1 e^{-y^2} \, dy \, dx = \int_0^1 \int_0^y e^{-y^2} \, dx \, dy = \int_0^1 y e^{-y^2} \, dy = \frac{1-e^{-1}}{2}$

例. 求 $\int_0^1 \int_y^1 \frac{\sin x}{x} \, dx \, dy$.

解. $\int_0^1 \int_y^1 \frac{\sin x}{x} \, dx \, dy = \int_0^1 \int_0^x \frac{\sin x}{x} \, dy \, dx = \int_0^1 \sin x \, dx = 1 - \cos 1$.

例. 求 $\int_0^1 \int_{\sin^{-1} y}^{\frac{\pi}{2}} \cos x \sqrt{1+\cos^2 x} \, dx \, dy$.

解.

- $\int_0^1 \int_{\sin^{-1}y}^{\frac{\pi}{2}} \cos x \sqrt{1 + \cos^2 x} \, dx \, dy = \int_0^{\frac{\pi}{2}} \int_0^{\sin x} \cos x \sqrt{1 + \cos^2 x} \, dy \, dx = \int_0^{\frac{\pi}{2}} \sin x \cos x \sqrt{1 + \cos^2 x} \, dx$
 $= \int_0^1 u \sqrt{1 + u^2} \, du = \int_1^2 \sqrt{v} \frac{1}{2} \, dv = \frac{2\sqrt{2} - 1}{3}$, 其中變數變換 $u = \cos x$ 與 $v = 1 + u^2$.
- 令 $u = \sin x$, $\int_{\sin^{-1}y}^{\frac{\pi}{2}} \cos x \sqrt{1 + \cos^2 x} \, dx = \int_y^1 \sqrt{2 - u^2} \, du$. 令 $u = \sqrt{2} \sin \theta$, 則 $\int_y^1 \sqrt{2 - u^2} \, du = \int_{\sin^{-1} \frac{y}{\sqrt{2}}}^{\frac{\pi}{4}} 2 \cos^2 \theta \, d\theta = \int_{\sin^{-1} \frac{y}{\sqrt{2}}}^{\frac{\pi}{4}} (1 + \cos 2\theta) \, d\theta = (\theta + \sin \theta \cos \theta) \Big|_{\sin^{-1} \frac{y}{\sqrt{2}}}^{\frac{\pi}{4}} = \frac{\pi}{4} + \frac{1}{2} - \sin^{-1} \frac{y}{\sqrt{2}} - \frac{y}{\sqrt{2}} \cdot \frac{\sqrt{2 - y^2}}{\sqrt{2}} = \frac{\pi}{4} + \frac{1}{2} - \sin^{-1} \frac{y}{\sqrt{2}} - \frac{y\sqrt{2 - y^2}}{2}$. 故 $\int_0^1 \int_{\sin^{-1}y}^{\frac{\pi}{2}} \cos x \sqrt{1 + \cos^2 x} \, dx \, dy = \int_0^1 \left(\frac{\pi}{4} + \frac{1}{2} - \sin^{-1} \frac{y}{\sqrt{2}} - \frac{y\sqrt{2 - y^2}}{2} \right) \, dy = \frac{\pi}{4} + \frac{1}{2} - \left(\frac{\pi}{4} + 1 - \sqrt{2} \right) - \left(\frac{\sqrt{2}}{3} - \frac{1}{6} \right) = \frac{2\sqrt{2} - 1}{3}$.

例. 求 $\int_0^\infty \frac{e^{-ax} - e^{-bx}}{x} \, dx$, $b > a > 0$.

解. 由 $\int_a^b e^{-xy} \, dy = \frac{1}{-x} e^{-xy} \Big|_{y=a}^{y=b} = \frac{1}{-x} (e^{-xb} - e^{-xa}) = \frac{e^{-ax} - e^{-bx}}{x}$, $\int_0^\infty \frac{e^{-ax} - e^{-bx}}{x} \, dx = \int_0^\infty \left(\int_a^b e^{-xy} \, dy \right) \, dx = \int_a^b \left(\int_0^\infty e^{-xy} \, dx \right) \, dy = \int_a^b \frac{1}{y} \, dy = \ln \frac{b}{a}$.

例. 求 $\int_0^\infty \frac{\tan^{-1} \pi x - \tan^{-1} x}{x} \, dx$.

解. 由 $\int_x^{\pi x} \frac{1}{1 + y^2} \, dy = \tan^{-1} \pi x - \tan^{-1} x$, $\int_0^\infty \frac{\tan^{-1} \pi x - \tan^{-1} x}{x} \, dx = \int_0^\infty \frac{1}{x} \left(\int_x^{\pi x} \frac{1}{1 + y^2} \, dy \right) \, dx = \int_0^\infty \int_x^{\pi x} \frac{1}{x(1 + y^2)} \, dy \, dx = \int_0^\infty \int_{\frac{y}{\pi}}^y \frac{1}{x(1 + y^2)} \, dx \, dy = \int_0^\infty \frac{1}{1 + y^2} \left(\int_{\frac{y}{\pi}}^y \frac{1}{x} \, dx \right) \, dy = \int_0^\infty \frac{1}{1 + y^2} \left(\ln y - \ln \frac{y}{\pi} \right) \, dy = \int_0^\infty \frac{\ln \pi}{1 + y^2} \, dy = \frac{\pi}{2} \ln \pi$.

例. 求 $\int_0^\infty \frac{\sin x}{x} \, dx$.

解. 由 $\int_0^\infty e^{-xy} \, dy = \frac{1}{x}$, $x > 0$; 又 $\int e^{ax} \sin bx \, dx = \frac{e^{ax}(a \cdot \sin bx - b \cdot \cos bx)}{a^2 + b^2}$.
 故 $\int_0^\infty \frac{\sin x}{x} \, dx = \int_0^\infty \sin x \left(\int_0^\infty e^{-xy} \, dy \right) \, dx = \int_0^\infty \left(\int_0^\infty e^{-xy} \sin x \, dx \right) \, dy = \int_0^\infty \left(\int_0^\infty e^{-xy} \sin x \, dx \right) \, dy = \int_0^\infty \frac{e^{-xy}((-y) \cdot \sin x - 1 \cdot \cos x) \Big|_{x=0}^{x=\infty}}{(-y)^2 + 1^2} \, dy = \int_0^\infty \frac{0 - (-1)}{1 + y^2} \, dy = \int_0^\infty \frac{1}{1 + y^2} \, dy = \tan^{-1} y \Big|_0^\infty = \frac{\pi}{2}$.

變數變換

單變數積分 $\int_{[a,b]} f(x) \, dx$ 中, 令 $x = x(u)$, 則 $dx = \frac{dx}{du} \, du$; $\int_{[a,b]} f(x) \, dx = \underbrace{\int_{x^{-1}[a,b]}}_{\text{由小到大}} f(x(u)) \left| \frac{dx}{du} \right| \, du$.

例.

- 求 $\int_0^1 \frac{dx}{\sqrt{1 - x^2}}$: 令 $x = \sin u$, 則 $dx = \cos u \cdot du$; x 由 0 至 1, 則 u 由 $\sin^{-1} 0 = 0$ 至 $\sin^{-1} 1 = \frac{\pi}{2}$. 故 $\int_0^1 \frac{dx}{\sqrt{1 - x^2}} = \int_0^{\frac{\pi}{2}} \frac{\cos u \cdot du}{\sqrt{1 - \sin^2 u}} = \int_0^{\frac{\pi}{2}} \frac{\cos u \cdot du}{\cos u} = \int_0^{\frac{\pi}{2}} 1 \cdot du = \frac{\pi}{2}$.

- 求 $\int_{\frac{1}{2}}^1 (1-2x)^2 dx$: 令 $u = 1-2x$, 亦即 $x = \frac{1}{2} - \frac{u}{2}$, 則 $dx = \frac{-1}{2} du$; x 由 $\frac{1}{2}$ 至 1, 則 u 由 $1-2 \cdot \frac{1}{2} = 0$ 至 $1-2 \cdot 1 = -1$. 故 $\int_{\frac{1}{2}}^1 (1-2x)^2 dx = \int_0^{-1} u^2 \left(\frac{-1}{2}\right) du = \int_{-1}^0 u^2 \frac{1}{2} du = \frac{1}{6}$.

定理 (重積分變數變換). 給定 $\Omega_{\mathbf{x}}, \Omega_{\mathbf{u}} \subseteq \mathbb{R}^n$, $\mathbf{x} = \mathbf{x}(\mathbf{u}) : \Omega_{\mathbf{u}} \rightarrow \Omega_{\mathbf{x}}$ 且

- \mathbf{x} 為嵌射 (亦即 $\mathbf{x}(\Omega_{\mathbf{u}}) = \Omega_{\mathbf{x}}$, $\mathbf{x}^{-1}(\Omega_{\mathbf{x}}) = \Omega_{\mathbf{u}}$)
- \mathbf{x} 之各分量函數 $x_j, j = 1, 2, \dots, n$ 為連續可偏微分

$$\bullet \forall \mathbf{u} \in \Omega_{\mathbf{u}}, \mathbf{x} \text{ 之 Jacobian } J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x_1}{\partial u_1} & \frac{\partial x_1}{\partial u_2} & \cdots & \frac{\partial x_1}{\partial u_n} \\ \frac{\partial x_2}{\partial u_1} & \frac{\partial x_2}{\partial u_2} & \cdots & \frac{\partial x_2}{\partial u_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial u_1} & \frac{\partial x_n}{\partial u_2} & \cdots & \frac{\partial x_n}{\partial u_n} \end{vmatrix}(\mathbf{u}) \neq 0$$

則對於可積函數 $f : \Omega_{\mathbf{x}} \rightarrow \mathbb{R}$, $\int_{\Omega_{\mathbf{x}}} f(\mathbf{x}) d\mathbf{x} = \int_{\mathbf{x}^{-1}(\Omega_{\mathbf{x}})} f(\mathbf{x}(\mathbf{u})) |J_{\mathbf{x}}(\mathbf{u})| d\mathbf{u} = \int_{\mathbf{x}^{-1}(\Omega_{\mathbf{x}})} f(\mathbf{x}(\mathbf{u})) \left| \frac{\partial \mathbf{x}}{\partial \mathbf{u}} \right| d\mathbf{u}$.

例. 利用變數變換 $x = u^2 - v^2, y = 2uv$ 求 $\int_{\Omega} y dA$, Ω 為 $y \geq 0, y^2 = 4 - 4x, y^2 = 4 + 4x$ 所圍成之區域.

解. 由 $\begin{cases} x = u^2 - v^2 \\ y = 2uv \end{cases}$, Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 2u & -2v \\ 2v & 2u \end{vmatrix} = 4(u^2 + v^2)$. 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $uv = 0, u^2v^2 + u^2 - v^2 - 1 = (v^2 + 1)(u^2 - 1) = 0, u^2v^2 - u^2 + v^2 - 1 = (u^2 + 1)(v^2 - 1) = 0$ 圍成, 亦即 $\Omega = \{0 \leq u \leq 1, 0 \leq v \leq 1\}$, 故 $\int_{\Omega} y dA = \int_0^1 \int_0^1 2uv \left| 4(u^2 + v^2) \right| du dv = 2$.

例. 求 $\int_{\Omega} (x+y)^2 dA$, Ω 為 $x+y=0, x+y=1, 2x-y=0, 2x-y=3$ 所圍成之平行四邊形.

解. 令 $\begin{cases} u = x+y \\ v = 2x-y \end{cases}$, 則 $\begin{cases} x = \frac{1}{3}u + \frac{1}{3}v \\ y = \frac{2}{3}u - \frac{1}{3}v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{2}{3} & -\frac{1}{3} \end{vmatrix} = -\frac{1}{3}$. 變數變換 $(x, y) \rightarrow (u, v)$ 後 $\Omega = \{0 \leq u \leq 1, 0 \leq v \leq 3\}$, 故 $\int_{\Omega} (x+y)^2 dA = \int_0^3 \int_0^1 u^2 \left| -\frac{1}{3} \right| du dv = \frac{1}{3}$.

例. 求 $\int_{\Omega} \sqrt{x+y} dA$, Ω 為頂點 $(0, 0), (a, 0), (0, a), a > 0$ 之三角形.

解. 令 $\begin{cases} u = x+y \\ v = x-y \end{cases}$, 則 $\begin{cases} x = \frac{1}{2}u + \frac{1}{2}v \\ y = \frac{1}{2}u - \frac{1}{2}v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$. 變數變換前 Ω 由 $x+y=0, x+y=a, x=0, y=0$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $u=0, u=a, u+v=0, u-v=0$ 圍成, 故 $\int_{\Omega} \sqrt{x+y} dA = \int_0^a \int_{-u}^u \sqrt{u} \left| -\frac{1}{2} \right| dv du = \int_0^a u\sqrt{u} du = \frac{2}{5} a^{\frac{5}{2}}$.

例. 求 $\int_{\Omega} (x+y) e^{x-y} dA$, Ω 為頂點 $(4, 0), (6, 2), (4, 4), (2, 2)$ 之四邊形.

解. 令 $\begin{cases} u = x+y \\ v = x-y \end{cases}$, 則 $\begin{cases} x = \frac{1}{2}u + \frac{1}{2}v \\ y = \frac{1}{2}u - \frac{1}{2}v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$. 變數變換前 Ω 由 $x-y=0, x-y=4, x+y=4, x+y=8$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $v=0, v=4, u=4, u=8$ 圍成, 故 $\int_{\Omega} (x+y) e^{x-y} dA = \int_0^4 \int_4^8 u e^v \left| -\frac{1}{2} \right| du dv = \frac{1}{2} \int_0^4 e^v dv \int_4^8 u du = 12(e^4 - 1)$.

例. 求 $\int_{\Omega} e^{\frac{x+y}{x-y}} dA$, Ω 為頂點 $(1, 0)$, $(2, 0)$, $(0, -2)$, $(0, -1)$ 之梯形.

解. 令 $\begin{cases} u = x + y \\ v = x - y \end{cases}$, 則 $\begin{cases} x = \frac{1}{2}u + \frac{1}{2}v \\ y = \frac{1}{2}u - \frac{1}{2}v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$. 變數變換前 Ω 由 $x - y = 1$, $x - y = 2$, $x = 0$, $y = 0$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $v = 1$, $v = 2$, $u + v = 0$, $u - v = 0$ 圍成, 故 $\int_{\Omega} e^{\frac{x+y}{x-y}} dA = \int_1^2 \int_{-v}^v e^{\frac{u}{v}} \left| -\frac{1}{2} \right| du dv = \frac{1}{2} \int_1^2 \left(v e^{\frac{u}{v}} \Big|_{u=-v}^{u=v} \right) dv = \frac{3(e - e^{-1})}{4}$.

例. 求 $y = x^2$, $y = 2x^2$, $x = y^2$, $x = 3y^2$ 所圍成之區域面積.

解. 令 $\begin{cases} u = \frac{y}{x^2} \\ v = \frac{x}{y^2} \end{cases}$, 則 $\begin{cases} x = (u^2 v)^{-\frac{1}{3}} \\ y = (uv^2)^{-\frac{1}{3}} \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} -\frac{1}{3}(u^2 v)^{-\frac{4}{3}} \cdot 2uv & -\frac{1}{3}(u^2 v)^{-\frac{4}{3}} \cdot u^2 \\ -\frac{1}{3}(uv^2)^{-\frac{4}{3}} \cdot v^2 & -\frac{1}{3}(uv^2)^{-\frac{4}{3}} \cdot 2uv \end{vmatrix}$
 $= \frac{1}{3u^2 v^2}$. 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $u = 1$, $u = 2$, $v = 1$, $v = 3$ 圍成, 亦即 $\Omega = \{1 \leq u \leq 2, 1 \leq v \leq 3\}$, 面積為 $\int_{\Omega} dA = \int_1^3 \int_1^2 \left| \frac{1}{3u^2 v^2} \right| du dv = \frac{1}{9}$.

例. 求 $\int_{\Omega} \frac{xy}{1+x^2y^2} dA$, Ω 為 $xy = 1$, $xy = 5$, $x = 1$, $x = 6$ 所圍成之區域.

解. 令 $\begin{cases} u = xy \\ v = x \end{cases}$, 則 $\begin{cases} x = v \\ y = \frac{u}{v} \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ \frac{1}{v} & -\frac{u}{v^2} \end{vmatrix} = -\frac{1}{v}$. 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $u = 1$, $u = 5$, $v = 1$, $v = 6$ 圍成, 亦即 $\Omega = \{1 \leq u \leq 5, 1 \leq v \leq 6\}$, 故 $\int_{\Omega} \frac{xy}{1+x^2y^2} dA = \int_1^6 \int_1^5 \frac{u}{1+u^2} \left| -\frac{1}{v} \right| du dv = \int_1^6 \frac{1}{v} dv \int_1^5 \frac{u}{1+u^2} du = \frac{\ln 6 \cdot \ln 13}{2}$.

例. 求 $\int_{\Omega} \frac{y}{x} dA$, Ω 為 $y = x$, $x^2 + 4y^2 = 4$, $y \geq 0$ 所圍成之區域.

解. 令 $\begin{cases} x = 2r \cos \theta \\ y = r \sin \theta \end{cases}$, Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} 2 \cos \theta & -2r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = 2r$. 變數變換 $(x, y) \rightarrow (r, \theta)$ 後 Ω 由 $r = 0$, $r = 1$, $\theta = 0$, $\theta = \tan^{-1} 2$ 圍成, 亦即 $\Omega = \{0 \leq r \leq 1, 0 \leq \theta \leq \tan^{-1} 2\}$, 故 $\int_{\Omega} \frac{y}{x} dA = \int_0^{\tan^{-1} 2} \int_0^1 \frac{1}{2} \tan \theta |2r| dr d\theta = \int_0^{\tan^{-1} 2} \tan \theta d\theta \int_0^1 r dr = -\frac{1}{2} \ln |\cos \theta| \Big|_0^{\tan^{-1} 2} = \frac{\ln 5}{4}$.

例. 求 $\int_{\Omega} \sin(9x^2 + 4y^2) dA$, Ω 為 $9x^2 + 4y^2 = 1$, $y \geq 0$, $x \geq 0$ 所圍成之區域.

解. 令 $\begin{cases} x = \frac{1}{3} r \cos \theta \\ y = \frac{1}{2} r \sin \theta \end{cases}$, Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \frac{1}{3} \cos \theta & -\frac{1}{3} r \sin \theta \\ \frac{1}{2} \sin \theta & \frac{1}{2} r \cos \theta \end{vmatrix} = \frac{1}{6} r$. 變數變換 $(x, y) \rightarrow (r, \theta)$ 後 Ω 由 $r = 0$, $r = 1$, $\theta = 0$, $\theta = \frac{\pi}{2}$ 圍成, 亦即 $\Omega = \{0 \leq r \leq 1, 0 \leq \theta \leq \frac{\pi}{2}\}$, 故 $\int_{\Omega} \sin(9x^2 + 4y^2) dA = \int_0^{\frac{\pi}{2}} \int_0^1 \sin r^2 \left| \frac{1}{6} r \right| dr d\theta = \frac{1}{6} \int_0^{\frac{\pi}{2}} d\theta \int_0^1 r \sin r^2 dr = \frac{\pi}{24} (1 - \cos 1)$.

例. 求 $\int_0^1 \int_0^{1-x} \sqrt{x+y} (y-2x)^2 dy dx$.

解. 令 $\begin{cases} u = x + y \\ v = y - 2x \end{cases}$, 則 $\begin{cases} x = \frac{1}{3}u - \frac{1}{3}v \\ y = \frac{2}{3}u + \frac{1}{3}v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{1}{3} \end{vmatrix} = \frac{1}{3}$. 變數變換前積分區域由 $x + y = 0$, $x + y = 1$, $x = 0$, $y = 0$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後積分區域由 $u = 0$, $u = 1$, $u - v = 0$, $2u + v = 0$ 圍成, 故 $\int_0^1 \int_0^{1-x} \sqrt{x+y} (y-2x)^2 dy dx = \int_0^1 \int_{-2u}^u \sqrt{u} v^2 \left| \frac{1}{3} \right| dv du = \frac{1}{3} \int_0^1 \sqrt{u} \left(\frac{v^3}{3} \Big|_{v=-2u}^{v=u} \right) du = \frac{2}{9}$.

例. 若 Ω 為頂點 $(0, 0)$, $(1, 0)$, $(0, 1)$ 之三角形, f 為可積, 證明 $\int_{\Omega} f(x+y) dA = \int_0^1 u f(u) du$.

解. 令 $\begin{cases} u = x + y \\ v = x \end{cases}$, 則 $\begin{cases} x = v \\ y = u - v \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ 1 & -1 \end{vmatrix} = -1$. 變數變換前 Ω 由 $x + y = 0$, $x + y = 1$, $x = 0$, $y = 0$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後 Ω 由 $u = 0$, $u = 1$, $v = 0$, $u - v = 0$ 圍成, 故 $\int_{\Omega} f(x+y) dA = \int_0^1 \int_0^u f(u) |-1| dv du = \int_0^1 u f(u) du$.

例. 1. 證明 $\int_0^1 \int_0^1 \frac{1}{1-xy} dx dy = \sum_{n=1}^{\infty} \frac{1}{n^2}$. 2. 以變數變換證明 $\int_0^1 \int_0^1 \frac{1}{1-xy} dx dy = \frac{\pi^2}{6}$.

解.

1. $|xy| < 1$ 時 $\frac{1}{1-xy} = \sum_{n=0}^{\infty} (xy)^n$ 且可逐項積分; 代入 $\int_0^1 \int_0^1 \frac{1}{1-xy} dx dy = \int_0^1 \int_0^1 \sum_{n=0}^{\infty} (xy)^n dx dy = \sum_{n=0}^{\infty} \int_0^1 x^n dx \int_0^1 y^n dy = \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} = \sum_{n=1}^{\infty} \frac{1}{n^2}$.

2. 令 $\begin{cases} x = \frac{u-v}{\sqrt{2}} \\ y = \frac{u+v}{\sqrt{2}} \end{cases}$; Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{vmatrix} = 1$. 變數變換前積分區域由 $x = 0$, $x = 1$, $y = 0$, $y = 1$ 所圍成, 變數變換 $(x, y) \rightarrow (u, v)$ 後積分區域由 $u - v = 0$, $u - v = \sqrt{2}$, $u + v = 0$, $u + v = \sqrt{2}$ 圍成, 故 $\int_0^1 \int_0^1 \frac{1}{1-xy} dx dy = \int_0^{\frac{\sqrt{2}}{2}} \int_{-u}^u \frac{dv du}{1 - \frac{u-v}{\sqrt{2}} \frac{u+v}{\sqrt{2}}} + \int_{\frac{\sqrt{2}}{2}}^{\sqrt{2}} \int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{dv du}{1 - \frac{u-v}{\sqrt{2}} \frac{u+v}{\sqrt{2}}} = 2 \left(\int_0^{\frac{\sqrt{2}}{2}} \int_{-u}^u \frac{dv du}{2 - u^2 + v^2} + \int_{\frac{\sqrt{2}}{2}}^{\sqrt{2}} \int_{-\sqrt{2}+u}^{\sqrt{2}-u} \frac{dv du}{2 - u^2 + v^2} \right) = 2 \left(\int_0^{\frac{\sqrt{2}}{2}} \frac{1}{\sqrt{2-u^2}} \left(\tan^{-1} \frac{v}{\sqrt{2-u^2}} \Big|_{v=-u}^{v=u} \right) du + \int_{\frac{\sqrt{2}}{2}}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}} \left(\tan^{-1} \frac{v}{\sqrt{2-u^2}} \Big|_{v=-\sqrt{2}+u}^{v=\sqrt{2}-u} \right) du \right) = 4 \left(\int_0^{\frac{\sqrt{2}}{2}} \frac{1}{\sqrt{2-u^2}} \tan^{-1} \frac{u}{\sqrt{2-u^2}} du + \int_{\frac{\sqrt{2}}{2}}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}} \tan^{-1} \frac{\sqrt{2}-u}{\sqrt{2-u^2}} du \right)$. 令 $u = \sqrt{2} \sin \theta$, 則 $du = \sqrt{2} \cos \theta d\theta$; 化簡 $4 \left(\int_0^{\frac{\sqrt{2}}{2}} \frac{1}{\sqrt{2-u^2}} \tan^{-1} \frac{u}{\sqrt{2-u^2}} du + \int_{\frac{\sqrt{2}}{2}}^{\sqrt{2}} \frac{1}{\sqrt{2-u^2}} \tan^{-1} \frac{\sqrt{2}-u}{\sqrt{2-u^2}} du \right) = 4 \left(\int_0^{\frac{\pi}{6}} \tan^{-1}(\tan \theta) d\theta + \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \tan^{-1} \left(\frac{1 - \sin \theta}{\cos \theta} \right) d\theta \right) = 4 \left(\int_0^{\frac{\pi}{6}} \theta d\theta + \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{1}{2} \left(\frac{\pi}{2} - \theta \right) d\theta \right) = \frac{\pi^2}{6}$; 其中 $\frac{1 - \sin \theta}{\cos \theta} = \frac{1 - \cos(\frac{\pi}{2} - \theta)}{\sin(\frac{\pi}{2} - \theta)} = \frac{2 \sin^2 \frac{1}{2}(\frac{\pi}{2} - \theta)}{2 \sin \frac{1}{2}(\frac{\pi}{2} - \theta) \cos \frac{1}{2}(\frac{\pi}{2} - \theta)} = \tan \frac{1}{2}(\frac{\pi}{2} - \theta)$.

極座標二重積分

令 $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$, Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r$.

例. 求 $\int_{\Omega} (3x + 4y^2) \, dA$, Ω 為 $x^2 + y^2 = 1$, $x^2 + y^2 = 4$ 與 $y \geq 0$ 所圍成之區域.

解. 變數變換 $(x, y) \rightarrow (r, \theta)$ 後 Ω 由 $r = 1$, $r = 2$, $\theta = 0$, $\theta = \pi$ 圍成, 故 $\int_{\Omega} (3x + 4y^2) \, dA = \int_0^{\pi} \int_1^2 (3r \cos \theta + 4r^2 \sin^2 \theta) r \, dr \, d\theta = \int_0^{\pi} (7 \cos \theta + 15 \sin^2 \theta) \, d\theta = \frac{15\pi}{2}$.

例. 求 $z = 1 - x^2 - y^2$ 與 $z \geq 0$ 所圍成之體積.

解. 體積為 $\int_{\Omega} (1 - x^2 - y^2) \, dA$, Ω 為單位圓 $x^2 + y^2 = 1$. 變數變換 $(x, y) \rightarrow (r, \theta)$ 後 Ω 由 $r = 0$, $r = 1$, $\theta = 0$, $\theta = 2\pi$ 圍成, 故 $\int_{\Omega} (1 - x^2 - y^2) \, dA = \int_0^{2\pi} \int_0^1 (1 - r^2) r \, dr \, d\theta = \frac{\pi}{2}$.

例. 求 $\int_{\Omega} \frac{y^2}{x^2} \, dA$, Ω 為 $0 < a^2 \leq x^2 + y^2 \leq b^2$ 與 $x \geq y \geq 0$ 所圍成之區域.

解. 變數變換 $(x, y) \rightarrow (r, \theta)$ 後 Ω 由 $r = a$, $r = b$, $\theta = 0$, $\theta = \frac{\pi}{4}$ 圍成, 故 $\int_{\Omega} \frac{y^2}{x^2} \, dA = \int_0^{\frac{\pi}{4}} \int_a^b \tan^2 \theta \cdot r \, dr \, d\theta = \frac{b^2 - a^2}{2} \int_0^{\frac{\pi}{4}} \tan^2 \theta \, d\theta = \frac{b^2 - a^2}{2} \int_0^{\frac{\pi}{4}} (\sec^2 \theta - 1) \, d\theta = \frac{b^2 - a^2}{2} (\tan \theta - \theta) \Big|_0^{\frac{\pi}{4}} = \frac{b^2 - a^2}{2} \left(1 - \frac{\pi}{4}\right)$.

例. 反變數變換: $\int_0^{\frac{\pi}{2}} \int_0^a \frac{r}{\sqrt{a^2 - r^2 \cos^2 \theta}} \, dr \, d\theta = \int_0^a \int_0^{\sqrt{a^2 - x^2}} \frac{1}{\sqrt{a^2 - x^2}} \, dy \, dx = \int_0^a \frac{\sqrt{a^2 - x^2}}{\sqrt{a^2 - x^2}} \, dx = a$.

曲面表面積

結論. 若曲面 S 由 $z = f(x, y)$, $(x, y) \in \Omega$ 所定義, 則 S 的表面積為 $\int_{\Omega} \sqrt{1 + f_x^2 + f_y^2} \, dA$.

證 (sketch). 以切平面近似曲面 S ; S 參數式為 $\mathbf{r} = (x, y, f(x, y))$, x, y 方向切向量分別為 $\mathbf{r}_x = (1, 0, f_x)$, $\mathbf{r}_y = (0, 1, f_y)$; $|\mathbf{r}_x \times \mathbf{r}_y| = |(-f_x, -f_y, 1)| = \sqrt{1 + f_x^2 + f_y^2}$.

例. 求半球面 $z = \sqrt{a^2 - x^2 - y^2}$ 與圓柱體 $(x - \frac{a}{2})^2 + y^2 = (\frac{a}{2})^2$ 相交區域之表面積.

解. 由 $z = f(x, y) = \sqrt{a^2 - x^2 - y^2}$, $f_x = \frac{-x}{\sqrt{a^2 - x^2 - y^2}}$, $f_y = \frac{-y}{\sqrt{a^2 - x^2 - y^2}}$, $dS = \sqrt{1 + f_x^2 + f_y^2} \, dx \, dy = \sqrt{1 + \frac{x^2}{a^2 - x^2 - y^2} + \frac{y^2}{a^2 - x^2 - y^2}} \, dx \, dy = \sqrt{\frac{a^2}{a^2 - x^2 - y^2}} \, dx \, dy$. 所求表面積為 $\int_{\Omega} dS$, $\Omega = \{(x, y) \mid (x - \frac{a}{2})^2 + y^2 \leq (\frac{a}{2})^2\}$; $\int_{\Omega} dS = \int_{\Omega} \sqrt{\frac{a^2}{a^2 - x^2 - y^2}} \, dx \, dy = 2 \int_0^{\frac{\pi}{2}} \int_0^{a \cos \theta} \sqrt{\frac{a^2}{a^2 - r^2}} r \, dr \, d\theta = 2a \int_0^{\frac{\pi}{2}} \left(-\sqrt{a^2 - r^2} \Big|_{r=0}^{r=a \cos \theta}\right) \, d\theta = 2a^2 \int_0^{\frac{\pi}{2}} (a - \sin \theta) \, d\theta = a^2(\pi - 2)$.

7.2 三重積分

一般區域積分

例. 若 E 為中心 0, 半徑 a 之球體, 寫出 $\int_E f(x, y, z) \, dV$.

解. $\int_E f(x, y, z) dV$

$$= \int_{-a}^a \int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} \int_{-\sqrt{a^2-x^2-y^2}}^{\sqrt{a^2-x^2-y^2}} f(x, y, z) dz dy dx = \int_{-a}^a \int_{-\sqrt{a^2-y^2}}^{\sqrt{a^2-y^2}} \int_{-\sqrt{a^2-x^2-y^2}}^{\sqrt{a^2-x^2-y^2}} f(x, y, z) dz dx dy$$

$$= \int_{-a}^a \int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} \int_{-\sqrt{a^2-x^2-z^2}}^{\sqrt{a^2-x^2-z^2}} f(x, y, z) dy dz dx = \int_{-a}^a \int_{-\sqrt{a^2-z^2}}^{\sqrt{a^2-z^2}} \int_{-\sqrt{a^2-x^2-z^2}}^{\sqrt{a^2-x^2-z^2}} f(x, y, z) dy dx dz$$

$$= \int_{-a}^a \int_{-\sqrt{a^2-y^2}}^{\sqrt{a^2-y^2}} \int_{-\sqrt{a^2-y^2-z^2}}^{\sqrt{a^2-y^2-z^2}} f(x, y, z) dx dz dy = \int_{-a}^a \int_{-\sqrt{a^2-z^2}}^{\sqrt{a^2-z^2}} \int_{-\sqrt{a^2-y^2-z^2}}^{\sqrt{a^2-y^2-z^2}} f(x, y, z) dx dy dz$$

例. 若 E 為第一卦限中 $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} \leq 1$ 圍成之區域, $a, b, c > 0$, 寫出 $\int_E f(x, y, z) dV$.


解. $\int_E f(x, y, z) dV$

$$= \int_0^a \int_0^{b-\frac{bx}{a}} \int_0^{c-\frac{cx}{a}-\frac{cy}{b}} f(x, y, z) dz dy dx = \int_0^b \int_0^{a-\frac{ay}{b}} \int_0^{c-\frac{cx}{a}-\frac{cy}{b}} f(x, y, z) dz dx dy$$

$$= \int_0^c \int_0^{a-\frac{ax}{c}} \int_0^{b-\frac{bx}{a}-\frac{bz}{c}} f(x, y, z) dy dz dx = \int_0^c \int_0^{b-\frac{bz}{c}} \int_0^{a-\frac{ay}{b}-\frac{az}{c}} f(x, y, z) dx dy dz$$

$$= \int_0^b \int_0^{c-\frac{cy}{b}} \int_0^{a-\frac{ax}{a}-\frac{az}{c}} f(x, y, z) dx dz dy = \int_0^c \int_0^{b-\frac{bz}{c}} \int_0^{a-\frac{ay}{b}-\frac{az}{c}} f(x, y, z) dx dy dz$$

例. 求 $x^2 + y^2 = a^2$ 與 $x^2 + z^2 = a^2$ 交集區域之表面積與體積.




$$(\text{表面積}) = 16 \int_0^a \int_0^{\sqrt{a^2-x^2}} \sqrt{1 + \left(\frac{-x}{\sqrt{a^2-x^2}}\right)^2} dy dx = 16 \int_0^a \int_0^{\sqrt{a^2-x^2}} \sqrt{\frac{a^2}{a^2-x^2}} dy dx$$

$$= 16a \int_0^a \frac{\sqrt{a^2-x^2}}{\sqrt{a^2-x^2}} dx = 16a^2.$$

$$(\text{體積}) = 8 \int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2}} dz dy dx = 8 \int_0^a (a^2 - x^2) dx = \frac{16a^3}{3}.$$

例. 若 E 為 $x = 1, y = 1, z = 1, x + y + z = 2$ 圍成之四面體, 求 $\int_E x dV$.




$$\int_E x dV = \int_0^1 \int_{1-x}^1 \int_{2-x-y}^1 x dz dy dx = \int_0^1 \int_{1-x}^1 x(x+y-1) dy dx$$

$$= \int_0^1 \int_{1-x}^1 (x^2 - x + xy) dy dx = \int_0^1 \left((x^2 - x)x + x \frac{1 - (1-x)^2}{2} \right) dx$$

$$= \int_0^1 \left(x^3 - x^2 + x \frac{2x - x^2}{2} \right) dx = \int_0^1 \frac{x^3}{2} dx = \frac{1}{8}$$

例. 若 E 為第一卦限中 $y^2 + z^2 = 1, x + y = 2, y = 0, z = 0$ 圍成之區域, 求 $\int_E z dV$.


解.



$$\begin{aligned}\int_E z \, dV &= \int_0^1 \int_0^{2-y} \int_0^{\sqrt{1-y^2}} z \, dz \, dx \, dy = \int_0^1 \frac{1}{2} (1-y^2)(2-y) \, dy \\ &= \frac{1}{2} \int_0^1 (2-y-2y^2+y^3) \, dy = \frac{1}{2} \left(2 - \frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) = \frac{1}{2} \cdot \frac{13}{12} = \frac{13}{24}\end{aligned}$$

例. 若 E 為第一卦限中 $y = x^2$, $y + z = 1$ 圍成之區域, 求 $\int_E x \, dV$.


解.



$$\begin{aligned}\int_E x \, dV &= \int_0^1 \int_{x^2}^1 \int_0^{1-y} x \, dz \, dy \, dx \\ &= \int_0^1 \int_{x^2}^1 x(1-y) \, dy \, dx \\ &= \int_0^1 \left(x(1-x^2) - x \left(\frac{y^2}{2} \Big|_{y=x^2}^{y=1} \right) \right) dx \\ &= \int_0^1 \left(x - x^3 + \frac{x^5}{2} - \frac{x}{2} \right) dx = \frac{1}{12}\end{aligned}$$

例. 若 E 為第一卦限中 $z = y^2$, $x + y = 1$ 圍成之區域, 求 $\int_E z \, dV$.


解.



$$\begin{aligned}\int_E z \, dV &= \int_0^1 \int_0^{1-x} \int_0^{y^2} z \, dz \, dy \, dx \\ &= \int_0^1 \int_0^{1-x} \frac{y^4}{2} \, dy \, dx \\ &= \int_0^1 \frac{(1-x)^5}{10} \, dx = \frac{1}{60}\end{aligned}$$

例. 若 E 為 $z = 1 - x^2$, $y = z$, $y = 0$, $z = 0$ 圍成之區域, 求 $\int_E dV$.


解.



$$\begin{aligned}\int_E dV &= \int_{-1}^1 \int_0^{1-x^2} \int_y^{1-x^2} dz \, dy \, dx \\ &= \int_{-1}^1 \frac{(1-x^2)^2}{2} \, dx = \frac{8}{15}\end{aligned}$$

例. 若 E 為 $x = 0$, $y = 0$, $z = 0$, $x + y + z = 2$, $x^2 + z = 1$ 圍成之區域, 求 $\int_E x \, dV$.

解.



$$\begin{aligned}\int_E x \, dV &= \int_0^1 \int_0^{1-x^2} \int_0^{2-x-z} x \, dy \, dz \, dx = \int_0^1 \int_0^{1-x^2} (2-x-z) x \, dz \, dx \\ &= \int_0^1 \left((2-x)x(1-x^2) - x \frac{(1-x^2)^2}{2} \right) dx \\ &= \frac{1}{2} \int_0^1 (-x^5 + 2x^4 - 2x^3 - 2x^2 + 3x) dx \\ &= \frac{1}{2} \left(-\frac{1}{6} + \frac{2}{5} - \frac{1}{2} - \frac{2}{3} + \frac{3}{2} \right) = \frac{17}{60}\end{aligned}$$

例. 若 E 為 $x + y + z = 1$, $x = 0$, $y = 0$, $z = 0$ 圍成之區域, 求 $\int_E z \, dV$.


$$\begin{aligned}\text{解. } \int_E z \, dV &= \int_0^1 \int_0^{1-x} \int_0^{1-x-y} z \, dz \, dy \, dx = \int_0^1 \int_0^{1-x} \frac{(1-x-y)^2}{2} dy \, dx = \int_0^1 -\frac{(1-x-y)^3}{6} \Big|_{y=0}^{y=1-x} dx \\ &= \int_0^1 \frac{1}{6}(1-x)^3 dx = -\frac{1}{24}(1-x)^4 \Big|_0^1 = \frac{1}{24}.\end{aligned}$$

例. 若 E 為 $x + 2y + z = 2$, $x = 2y$, $x = 0$, $z = 0$ 圍成之區域, 求 $\int_E y \, dV$.

$$\begin{aligned}\text{解. } \int_E y \, dV &= \int_0^1 \int_{\frac{x}{2}}^{\frac{2-x}{2}} \int_0^{2-x-2y} y \, dz \, dy \, dx = \int_0^1 \int_{\frac{x}{2}}^{\frac{2-x}{2}} (2-x-2y) y \, dy \, dx = \int_0^1 \left(\frac{(2-x)y^2}{2} - \frac{2y^3}{3} \right) \Big|_{\frac{x}{2}}^{\frac{2-x}{2}} dx \\ &= \int_0^1 \left(\frac{2-x}{2} \left(\left(\frac{2-x}{2} \right)^2 - \left(\frac{x}{2} \right)^2 \right) - \frac{2}{3} \left(\left(\frac{2-x}{2} \right)^3 - \left(\frac{x}{2} \right)^3 \right) \right) dx = \int_0^1 \left(\frac{2-x}{2} (1-x) - \frac{2}{3} (1-x) \left(\left(\frac{2-x}{2} \right)^2 + \frac{2-x}{2} \cdot \frac{x}{2} + \left(\frac{x}{2} \right)^2 \right) \right) dx \\ &= \frac{1}{6} \int_0^1 (x^3 - 3x + 2) dx = \frac{1}{6} \left(\frac{1}{4} - \frac{3}{2} + 2 \right) = \frac{1}{8}.\end{aligned}$$

變數變換


柱面座標



$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases} \iff \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1} \frac{y}{x} \\ z = z \end{cases}$$

$$\text{Jacobian } J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial z} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial z} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial z} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$$


球面座標



$$\begin{cases} x = \rho \sin \varphi \cos \theta \\ y = \rho \sin \varphi \sin \theta \\ z = \rho \cos \varphi \end{cases} \iff \begin{cases} \rho = \sqrt{x^2 + y^2 + z^2} \\ \theta = \tan^{-1} \frac{y}{x} \\ \varphi = \tan^{-1} \frac{\sqrt{x^2 + y^2}}{z} \end{cases}$$


$$\text{Jacobian } J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial \varphi} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial \varphi} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial \varphi} \end{vmatrix} = \begin{vmatrix} \sin \varphi \cos \theta & -\rho \sin \varphi \sin \theta & \rho \cos \varphi \cos \theta \\ \sin \varphi \sin \theta & \rho \sin \varphi \cos \theta & \rho \cos \varphi \sin \theta \\ \cos \varphi & 0 & -\rho \sin \varphi \end{vmatrix} = -\rho^2 \sin \varphi$$

例. 半徑為 a 之球中心對稱鑽半徑為 b 之圓孔, $a > b > 0$, 求球剩下的體積.




- 柱面座標: $\int_E dV = 8 \int_0^{\frac{\pi}{2}} \int_b^a \int_0^{\sqrt{a^2 - r^2}} r \, dz \, dr \, d\theta = \frac{4\pi}{3}(a^2 - b^2)^{\frac{3}{2}}$
- 球面座標: $\int_E dV = 2 \int_{\sin^{-1} \frac{b}{a}}^{\frac{\pi}{2}} \int_0^{2\pi} \int_{\frac{b}{\sin \varphi}}^a \rho^2 \sin \varphi \, d\rho \, d\theta \, d\varphi = \frac{4\pi}{3}(a^2 - b^2)^{\frac{3}{2}}$

例. 若 E 為 $x^2 + y^2 \leq z^2$, $x^2 + y^2 + z^2 \leq 1$ 與 $z \geq 0$ 圍成之區域, 求 $\int_E \sqrt{x^2 + y^2 + z^2} \, dV$.



- 柱面座標: $\int_E \sqrt{x^2 + y^2 + z^2} \, dV = \int_0^{2\pi} \int_{\frac{1}{\sqrt{2}}}^1 \int_r^{\sqrt{1-r^2}} \sqrt{r^2 + z^2} \cdot r \, dz \, dr \, d\theta$
- 球面座標: $\int_E \sqrt{x^2 + y^2 + z^2} \, dV = \int_0^{\frac{\pi}{4}} \int_0^{2\pi} \int_0^1 \rho \cdot \rho^2 \sin \varphi \, d\rho \, d\theta \, d\varphi = \frac{\pi}{2} \left(1 - \frac{1}{\sqrt{2}}\right)$


例. 若 E 為 $0 \leq z \leq \sqrt{x^2 + y^2}$ 與 $x^2 + y^2 \leq 1$ 圍成之區域, 求 $\int_E z \sqrt{x^2 + y^2 + z^2} \, dV$.



- 柱面座標: $\int_E z \sqrt{x^2 + y^2 + z^2} \, dV = \int_0^{2\pi} \int_0^1 \int_0^r z \sqrt{r^2 + z^2} \cdot r \, dz \, dr \, d\theta = 2\pi \int_0^1 \frac{r}{3} (r^2 + z^2)^{\frac{3}{2}} \Big|_{z=0}^{z=r} \, dr = \frac{2\pi}{3} \int_0^1 r \cdot (2^{\frac{3}{2}} - 1) r^3 \, dr = \frac{2\pi (2^{\frac{3}{2}} - 1)}{15}$
- 球面座標: $\int_E z \sqrt{x^2 + y^2 + z^2} \, dV = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^{2\pi} \int_0^{\frac{1}{\sin \varphi}} \rho \cos \varphi \cdot \rho \cdot \rho^2 \sin \varphi \, d\rho \, d\theta \, d\varphi$

例. 若 E 為 $z = x^2 + y^2$ 與 $z \leq \sqrt{x^2 + y^2}$ 圍成之區域, 求 $\int_E (x^2 + y^2 + z^2) \, dV$.

解.



- 柱面座標: $\int_E z(x^2 + y^2 + z^2) dV = \int_0^{2\pi} \int_0^1 \int_{r^2}^r z(r^2 + z^2) \cdot r dz dr d\theta$
 $= 2\pi \int_0^1 \int_{r^2}^r (r^3 z + r z^3) dz dr = 2\pi \int_0^1 \left(\frac{r^3}{2}(r^2 - r^4) + \frac{r}{4}(r^4 - r^8) \right) dr = \frac{3\pi}{40}$
- 球面座標: $z = x^2 + y^2 \implies \rho \cos \varphi = \rho^2 \sin^2 \varphi \implies \rho = \frac{\cos \varphi}{\sin^2 \varphi}$;
 $z = \sqrt{x^2 + y^2} \implies \rho \cos \varphi = \rho \sin \varphi \implies \tan \varphi = 1$;
 $\int_E z(x^2 + y^2 + z^2) dV = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \int_0^{2\pi} \int_{\frac{\cos \varphi}{\sin^2 \varphi}}^{\frac{\cos \varphi}{\sin^2 \varphi}} \rho \cos \varphi \cdot \rho^2 \cdot \rho^2 \sin \varphi d\rho d\theta d\varphi$


例. 若 E 為 $y = x^2 + z^2$ 與 $y = 4$ 圍成之區域, 求 $\int_E \sqrt{x^2 + z^2} dV$.

解. 令 $\begin{cases} x = r \cos \theta \\ z = r \sin \theta \end{cases}$, 投影之 Ω 為 $x^2 + z^2 \leq 4$, 則 $\int_E \sqrt{x^2 + z^2} dV = \int_{\Omega} \int_{x^2+z^2}^4 \sqrt{x^2 + z^2} dy dA$
 $= \int_{\Omega} (4 - (x^2 + z^2)) \sqrt{x^2 + z^2} dA = \int_0^{2\pi} \int_0^2 (4 - r^2) r \cdot r dr d\theta = 2\pi \int_0^2 (4r^2 - r^4) dr = 2\pi \cdot \left(\frac{4 \cdot 2^3}{3} - \frac{2^5}{5} \right) = \frac{128\pi}{15}$.

例. 若 $E = \{(x, y, z) \mid x^2 + y^2 + (z - 1)^2 \leq 1\}$, 求 $\int_E (x^2 + y^2 + z^2)^{\frac{5}{2}} dV$.

解. 代入球面座標於 $x^2 + y^2 + (z - 1)^2 \leq 1 \implies (\rho \sin \varphi \cos \theta)^2 + (\rho \sin \varphi \sin \theta)^2 + (\rho \cos \varphi - 1)^2 \leq 1 \implies$
 $\rho^2 \sin^2 \varphi + \rho^2 \cos^2 \varphi - 2\rho \cos \varphi + 1 \leq 1 \implies \rho^2 \leq 2\rho \cos \varphi \implies \rho \leq 2 \cos \varphi$, 故 $\int_E (x^2 + y^2 + z^2)^{\frac{5}{2}} dV =$
 $\int_0^{\frac{\pi}{2}} \int_0^{2\pi} \int_0^{2 \cos \varphi} \rho^5 \cdot \rho^2 \sin \varphi d\rho d\theta d\varphi = 2\pi \int_0^{\frac{\pi}{2}} \frac{(2 \cos \varphi)^8}{8} \sin \varphi d\varphi = \frac{64\pi}{9} (-\cos^9 \varphi) \Big|_0^{\frac{\pi}{2}} = \frac{64\pi}{9}$.

例. 若 E 為 $z = \sqrt{x^2 + y^2}$ 與 $x^2 + y^2 + (z - 1)^2 = 1$ 於第一卦限圍成之區域, 求其體積.



- 柱面座標: $\int_E dV = \int_0^{\frac{\pi}{2}} \int_0^{2\pi} \int_r^{1+\sqrt{1-r^2}} r dz dr d\theta = \frac{\pi}{4}$

- 球面座標: $\int_E dV = \int_0^{\frac{\pi}{4}} \int_0^{2\pi} \int_0^{2 \cos \varphi} \rho^2 \sin \varphi d\rho d\theta d\varphi = \frac{\pi}{4}$

7.3 綜合問題

例. 已知 $\forall 0 < \alpha < 1$, $\int_0^{\infty} \frac{x^{\alpha-1}}{1+x} dx = \frac{\pi}{\sin \alpha \pi}$, 證明 $\Gamma(\alpha) \Gamma(1-\alpha) = \frac{\pi}{\sin \alpha \pi}$.

解. $\Gamma(\alpha) \Gamma(1-\alpha) = \int_0^{\infty} e^{-t} t^{\alpha-1} dt \int_0^{\infty} e^{-s} s^{-\alpha} ds = \int_0^{\infty} \int_0^{\infty} e^{-(t+s)} t^{\alpha-1} s^{-\alpha} ds dt$. 令 $\begin{cases} u = s+t \\ v = \frac{t}{s} \end{cases}$, 則 $\begin{cases} s = \frac{u}{1+v} \\ t = \frac{uv}{1+v} \end{cases}$;

Jacobian $J_{\mathbf{x}}(\mathbf{u}) = \frac{\partial \mathbf{x}}{\partial \mathbf{u}}(\mathbf{u}) = \begin{vmatrix} \frac{\partial s}{\partial u} & \frac{\partial s}{\partial v} \\ \frac{\partial t}{\partial u} & \frac{\partial t}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{1+v} & \frac{-u}{(1+v)^2} \\ \frac{v}{1+v} & \frac{u}{(1+v)^2} \end{vmatrix} = \frac{u}{(1+v)^2}$. 變數變換 $(s, t) \rightarrow (u, v)$ 後積分

範圍仍為 $0 < u < \infty$, $0 < v < \infty$, 故 $\int_0^{\infty} \int_0^{\infty} e^{-(t+s)} t^{\alpha-1} s^{-\alpha} ds dt = \int_0^{\infty} \int_0^{\infty} e^{-u} \left(\frac{t}{s} \right)^{\alpha} t^{-1} ds dt =$
 $\int_0^{\infty} \int_0^{\infty} e^{-u} v^{\alpha} \frac{1+v}{uv} \frac{u}{(1+v)^2} du dv = \int_0^{\infty} \int_0^{\infty} e^{-u} \frac{v^{\alpha-1}}{1+v} du dv = \int_0^{\infty} e^{-u} du \int_0^{\infty} \frac{v^{\alpha-1}}{1+v} dv = \int_0^{\infty} \frac{v^{\alpha-1}}{1+v} dv = \frac{\pi}{\sin \alpha \pi}$.

例. Beta 函數定義為 $B(m, n) = \int_0^1 t^{m-1}(1-t)^{n-1} dt \quad \forall m, n > 0$. 證明 $B(m, n) = B(n, m) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$.

證. $\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt$; 令 $t = u^2$, 則 $\Gamma(x) = 2 \int_0^\infty e^{-u^2} u^{2x-1} du$.

$$\begin{aligned} \Gamma(m)\Gamma(n) &= \left(2 \int_0^\infty e^{-u^2} u^{2m-1} du\right) \left(2 \int_0^\infty e^{-v^2} v^{2n-1} dv\right) = 4 \int_0^\infty \int_0^\infty e^{-(u^2+v^2)} u^{2m-1} v^{2n-1} du dv \\ &= 4 \int_0^{\frac{\pi}{2}} \int_0^\infty e^{-r^2} (r \cos \theta)^{2m-1} (r \sin \theta)^{2n-1} r dr d\theta = \left(2 \int_0^\infty e^{-r^2} r^{2(m+n)-1} dr\right) \left(2 \int_0^{\frac{\pi}{2}} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta\right) \\ &= \Gamma(m+n) \cdot 2 \int_0^{\frac{\pi}{2}} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta. \text{ 令 } t = \cos^2 \theta, \text{ 則 } dt = -2 \cos \theta \sin \theta d\theta; \text{ 故 } 2 \int_0^{\frac{\pi}{2}} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta \\ &= \int_0^{\frac{\pi}{2}} \cos^{2m-2} \theta \sin^{2n-2} \theta (-2 \cos \theta \sin \theta d\theta) = - \int_1^0 t^{m-1} (1-t)^{n-1} dt = \int_0^1 t^{m-1} (1-t)^{n-1} dt = B(m, n). \end{aligned}$$

例. 設 $V_n(a)$ 為半徑 a 之 n 維球體積, $n \geq 1, a > 0$; 證明 $V_n(1) = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2} + 1)}$.

證. $V_n(a) = \int_{x_1^2 + x_2^2 + \dots + x_n^2 \leq a^2} dx_1 dx_2 \dots dx_n$. 變數變換 $x_i = au_i \quad \forall i = 1, 2, \dots, n$, 則 Jacobian 為 a^n , 積分範圍變為 $u_1^2 + u_2^2 + \dots + u_n^2 \leq 1$, 故 $V_n(a) = a^n V_n(1)$. $V_n(1) = \int_{u_1^2 + u_2^2 + \dots + u_n^2 \leq 1} du_1 du_2 \dots du_n$

$$\begin{aligned} &= \int_{u_n^2 \leq 1} \left(\int_{u_1^2 + u_2^2 + \dots + u_{n-1}^2 \leq 1 - u_n^2} du_1 du_2 \dots du_{n-1} \right) du_n = \int_{-1}^1 V_{n-1}(\sqrt{1 - u_n^2}) du_n = V_{n-1}(1) \cdot \int_{-1}^1 (1 - u_n^2)^{\frac{n-1}{2}} du_n \\ &= V_{n-1}(1) \cdot 2 \int_0^1 (1 - u_n^2)^{\frac{n-1}{2}} du_n. \text{ 令 } t = u_n^2 \Rightarrow u_n = \sqrt{t} \Rightarrow du_n = \frac{dt}{2\sqrt{t}}, \text{ 則 } 2 \int_0^1 (1 - u_n^2)^{\frac{n-1}{2}} du_n = \\ &= \int_0^1 (1 - t)^{\frac{n-1}{2}} t^{-\frac{1}{2}} dt = B\left(\frac{n+1}{2}, \frac{1}{2}\right) = \frac{\Gamma(\frac{n+1}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+2}{2})}. \text{ 故 } V_n(1) = \frac{\Gamma(\frac{n+1}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+2}{2})} V_{n-1}(1) = \frac{\Gamma(\frac{n+1}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+2}{2})} \\ &= \frac{\Gamma(\frac{n}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+1}{2})} V_{n-2}(1) = \frac{\Gamma(\frac{n+1}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+2}{2})} \cdot \frac{\Gamma(\frac{n}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n+1}{2})} \cdot \frac{\Gamma(\frac{n-1}{2}) \cdot \Gamma(\frac{1}{2})}{\Gamma(\frac{n}{2})} V_{n-3}(1) = \dots = \frac{\Gamma(\frac{3}{2}) \cdot (\Gamma(\frac{1}{2}))^{n-1}}{\Gamma(\frac{n+2}{2})} V_1(1) = \\ &= \frac{\frac{1}{2}\sqrt{\pi} \cdot (\sqrt{\pi})^{n-1}}{\Gamma(\frac{n}{2} + 1)} \cdot 2 = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2} + 1)}. \end{aligned}$$

例. 設 $V_n(a)$ 為 n 維區域 $|x_1| + |x_2| + \dots + |x_n| \leq a$ 之體積, $n \geq 1, a > 0$; 證明 $V_n(a) = a^n \frac{2^n}{n!}$.

證. $V_n(a) = \int_{|x_1| + |x_2| + \dots + |x_n| \leq a} dx_1 dx_2 \dots dx_n$. 變數變換 $x_i = au_i \quad \forall i = 1, 2, \dots, n$, 則 Jacobian 為 a^n , 積分範圍變為 $|u_1| + |u_2| + \dots + |u_n| \leq 1$, 故 $V_n(a) = a^n V_n(1)$. $V_n(1) = \int_{|u_1| + |u_2| + \dots + |u_n| \leq 1} du_1 du_2 \dots du_n$

$$\begin{aligned} &= \int_{|u_n| \leq 1} \left(\int_{|u_1| + |u_2| + \dots + |u_{n-1}| \leq 1 - |u_n|} du_1 du_2 \dots du_{n-1} \right) du_n = \int_{-1}^1 V_{n-1}(1 - |u_n|) du_n = V_{n-1}(1) \int_{-1}^1 (1 - |u_n|)^{n-1} du_n \\ &= V_{n-1}(1) \left(\int_{-1}^0 (1 + u_n)^{n-1} du_n + \int_0^1 (1 - u_n)^{n-1} du_n \right) = \frac{2}{n} V_{n-1}(1). \text{ 故 } V_n(1) = \frac{2^{n-1}}{n \cdot (n-1) \dots 2} V_1(1) = \\ &= \frac{2^{n-1}}{n!} \cdot 2 = \frac{2^n}{n!}, V_n(a) = a^n \frac{2^n}{n!}. \end{aligned}$$