

# Introduction to Financial Models

## Lecture 03: Surprises & Paradoxes III

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4
  - If heads appears on the third flip, win \$8

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4
  - If heads appears on the third flip, win \$8
  - In general, if heads appears on the  $n$ -th flip, win  $\$2^n$



# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4
  - If heads appears on the third flip, win \$8
  - In general, if heads appears on the  $n$ -th flip, win  $\$2^n$
- The expected value calculation:

$$E X = \sum_{n=1}^{\infty} 2^n \cdot P(\text{heads on flip } n) = \sum_{n=1}^{\infty} 2^n \cdot \frac{1}{2^n} = \sum_{n=1}^{\infty} 1 = \infty$$

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4
  - If heads appears on the third flip, win \$8
  - In general, if heads appears on the  $n$ -th flip, win  $\$2^n$
- The expected value calculation:

$$E X = \sum_{n=1}^{\infty} 2^n \cdot P(\text{heads on flip } n) = \sum_{n=1}^{\infty} 2^n \cdot \frac{1}{2^n} = \sum_{n=1}^{\infty} 1 = \infty$$

- The paradox: The game has infinite expected value, but most people would only pay a small amount to play

# St. Petersburg Paradox

- Proposed by Nicolas Bernoulli in 1713, analyzed by Daniel Bernoulli in 1738
- The game:
  - Flip a fair coin until it shows heads
  - If heads appears on the first flip, win \$2
  - If heads appears on the second flip, win \$4
  - If heads appears on the third flip, win \$8
  - In general, if heads appears on the  $n$ -th flip, win  $\$2^n$
- The expected value calculation:

$$E X = \sum_{n=1}^{\infty} 2^n \cdot P(\text{heads on flip } n) = \sum_{n=1}^{\infty} 2^n \cdot \frac{1}{2^n} = \sum_{n=1}^{\infty} 1 = \infty$$

- The paradox: The game has infinite expected value, but most people would only pay a small amount to play
- If people maximize expected value, they should be willing to pay any finite amount to play

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1 \text{ million}$ ,  $E X \approx \$21$

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1$  million,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1 \text{ million}$ ,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"



# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1 \text{ million}$ ,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"
  - Proposed logarithmic utility function:  $U(w) = \log(w)$

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1$  million,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"
  - Proposed logarithmic utility function:  $U(w) = \log(w)$
  - Diminishing marginal utility: Each extra dollar adds less utility

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1$  million,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"
  - Proposed logarithmic utility function:  $U(w) = \log(w)$
  - Diminishing marginal utility: Each extra dollar adds less utility
- Gabriel Cramer (1728) suggested:  $U(w) = \sqrt{w}$

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1$  million,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"
  - Proposed logarithmic utility function:  $U(w) = \log(w)$
  - Diminishing marginal utility: Each extra dollar adds less utility
- Gabriel Cramer (1728) suggested:  $U(w) = \sqrt{w}$
- With logarithmic utility, expected utility is finite:

$$E U(X) = \sum_{n=1}^{\infty} \log(2^n) \cdot \frac{1}{2^n} = \sum_{n=1}^{\infty} \frac{n \log(2)}{2^n} = 2 \log(2) < \infty$$

# Early Solutions to the Paradox

- Practical resolution: No casino has infinite resources
  - With a capped maximum payout of  $M$ ,  $E X = \log_2(M) + \frac{M}{2^{\lfloor \log_2 M \rfloor}}$
  - For  $M = 2^{20} \approx \$1$  million,  $E X \approx \$21$
- Daniel Bernoulli's insight (1738): People value money differently
  - "The value of an item must not be based on its price, but rather on the utility it yields"
  - Proposed logarithmic utility function:  $U(w) = \log(w)$
  - Diminishing marginal utility: Each extra dollar adds less utility
- Gabriel Cramer (1728) suggested:  $U(w) = \sqrt{w}$
- With logarithmic utility, expected utility is finite:

$$E U(X) = \sum_{n=1}^{\infty} \log(2^n) \cdot \frac{1}{2^n} = \sum_{n=1}^{\infty} \frac{n \log(2)}{2^n} = 2 \log(2) < \infty$$

- This amounts to  $E U(X) \approx \$1.39$ , explaining why people would only pay a small amount

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty
- It seeks to address paradoxes like St. Petersburg by incorporating risk preferences



# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty
- It seeks to address paradoxes like St. Petersburg by incorporating risk preferences
- Key insight: people care about the utility of outcomes, not just monetary values

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty
- It seeks to address paradoxes like St. Petersburg by incorporating risk preferences
- Key insight: people care about the utility of outcomes, not just monetary values
- Decision-making is based on:

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty
- It seeks to address paradoxes like St. Petersburg by incorporating risk preferences
- Key insight: people care about the utility of outcomes, not just monetary values
- Decision-making is based on:
  - ① Probabilities of different outcomes

# The Expected Utility Hypothesis

## Mathematical Formulation

The agent prefers the r.v.  $X$  to r.v.  $Y$  if and only if  $E U(X) > E U(Y)$ , where  $U: \mathbb{R} \mapsto \mathbb{R}$  is the agent's utility function.

- Expected utility theory attempts to explain how people make decisions under uncertainty
- It seeks to address paradoxes like St. Petersburg by incorporating risk preferences
- Key insight: people care about the utility of outcomes, not just monetary values
- Decision-making is based on:
  - 1 Probabilities of different outcomes
  - 2 Subjective valuation (utility) of those outcomes

# Properties of Utility Functions

- Three common assumptions about utility functions:

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases



# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$
  - Power utility:  $U(w) = \frac{w^{1-\gamma}-1}{1-\gamma}$  for  $\gamma > 0, \gamma \neq 1$

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$
  - Power utility:  $U(w) = \frac{w^{1-\gamma}-1}{1-\gamma}$  for  $\gamma > 0, \gamma \neq 1$
  - Exponential utility:  $U(w) = -e^{-\gamma w}$  for  $\gamma > 0$

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$
  - Power utility:  $U(w) = \frac{w^{1-\gamma}-1}{1-\gamma}$  for  $\gamma > 0, \gamma \neq 1$
  - Exponential utility:  $U(w) = -e^{-\gamma w}$  for  $\gamma > 0$
  - Quadratic utility:  $U(w) = w - \frac{\gamma}{2} w^2$  for  $w < \frac{1}{\gamma}$

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$
  - Power utility:  $U(w) = \frac{w^{1-\gamma}-1}{1-\gamma}$  for  $\gamma > 0, \gamma \neq 1$
  - Exponential utility:  $U(w) = -e^{-\gamma w}$  for  $\gamma > 0$
  - Quadratic utility:  $U(w) = w - \frac{\gamma}{2} w^2$  for  $w < \frac{1}{\gamma}$
- The parameter  $\gamma$  reflects the degree of risk aversion

# Properties of Utility Functions

- Three common assumptions about utility functions:
  - **More is better:**  $U'(w) > 0$  (monotonically increasing)
  - **Risk aversion:**  $U''(w) < 0$  (concave function)
  - **Decreasing absolute risk aversion:**  $-\frac{U''(w)}{U'(w)}$  decreases as  $w$  increases
- Commonly used utility functions:
  - Logarithmic:  $U(w) = \log(w)$
  - Power utility:  $U(w) = \frac{w^{1-\gamma}-1}{1-\gamma}$  for  $\gamma > 0, \gamma \neq 1$
  - Exponential utility:  $U(w) = -e^{-\gamma w}$  for  $\gamma > 0$
  - Quadratic utility:  $U(w) = w - \frac{\gamma}{2} w^2$  for  $w < \frac{1}{\gamma}$
- The parameter  $\gamma$  reflects the degree of risk aversion
- For power utility,  $\gamma = 1$  corresponds to logarithmic utility (by L'Hôpital's rule)

# Risk Aversion and Risk Premium I

- Risk Aversion



# Risk Aversion and Risk Premium I

- Risk Aversion
  - The **risk aversion** is the preference for a certain amount over a gamble with the same expected value

# Risk Aversion and Risk Premium I

- Risk Aversion
  - The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
  - Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)

# Risk Aversion and Risk Premium I

- Risk Aversion
  - The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
  - Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)
  - Mathematically represented by a concave utility function:  $U''(w) < 0$

# Risk Aversion and Risk Premium I

- Risk Aversion
  - The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
  - Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)
  - Mathematically represented by a concave utility function:  $U''(w) < 0$
- Risk Premium

# Risk Aversion and Risk Premium I

- Risk Aversion
  - The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
  - Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)
  - Mathematically represented by a concave utility function:  $U''(w) < 0$
- Risk Premium
  - Let  $\tilde{X}$  be a random variable with  $E \tilde{X} = 0$  (a fair gamble)

# Risk Aversion and Risk Premium I

- Risk Aversion

- The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
- Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)
- Mathematically represented by a concave utility function:  $U''(w) < 0$

- Risk Premium

- Let  $\tilde{X}$  be a random variable with  $E \tilde{X} = 0$  (a fair gamble)
- A risk-averse person would pay to avoid this gamble

# Risk Aversion and Risk Premium I

- Risk Aversion

- The **risk aversion** is the preference for a certain amount over a gamble with the same expected value
- Example: Preferring \$50 with certainty over a 50% chance of \$100 (and 50% chance of \$0)
- Mathematically represented by a concave utility function:  $U''(w) < 0$

- Risk Premium

- Let  $\tilde{X}$  be a random variable with  $E \tilde{X} = 0$  (a fair gamble)
- A risk-averse person would pay to avoid this gamble
- The **risk premium**  $\pi$  is the maximum amount they would pay:

$$U(w - \pi) = E U(w + \tilde{X}) \quad (1)$$

# Risk Aversion and Risk Premium II

- For small risks, we can use Taylor expansion

$$U(w + \tilde{X}) \approx U(w) + U'(w)\tilde{X} + \frac{1}{2}U''(w)\tilde{X}^2$$



# Risk Aversion and Risk Premium II

- For small risks, we can use Taylor expansion

$$U(w + \tilde{X}) \approx U(w) + U'(w)\tilde{X} + \frac{1}{2}U''(w)\tilde{X}^2$$

- By  $E\tilde{X} = 0$  and  $\text{var } \tilde{X} = E\tilde{X}^2 - (E\tilde{X})^2 = E\tilde{X}^2$ ,

$$E U(w + \tilde{X}) \approx U(w) + U'(w) E\tilde{X} + \frac{1}{2} U''(w) E\tilde{X}^2 = U(w) + \frac{1}{2} U''(w) \text{var } \tilde{X}$$

# Risk Aversion and Risk Premium II

- For small risks, we can use Taylor expansion

$$U(w + \tilde{X}) \approx U(w) + U'(w)\tilde{X} + \frac{1}{2}U''(w)\tilde{X}^2$$

- By  $E\tilde{X} = 0$  and  $\text{var } \tilde{X} = E\tilde{X}^2 - (E\tilde{X})^2 = E\tilde{X}^2$ ,

$$E U(w + \tilde{X}) \approx U(w) + U'(w)E\tilde{X} + \frac{1}{2}U''(w)E\tilde{X}^2 = U(w) + \frac{1}{2}U''(w)\text{var } \tilde{X}$$

- Similarly, for small premium, one can use Taylor expansion

$$U(w - \pi) \approx U(w) - \pi U'(w)$$

# Risk Aversion and Risk Premium II

- For small risks, we can use Taylor expansion

$$U(w + \tilde{X}) \approx U(w) + U'(w)\tilde{X} + \frac{1}{2}U''(w)\tilde{X}^2$$

- By  $E\tilde{X} = 0$  and  $\text{var } \tilde{X} = E\tilde{X}^2 - (E\tilde{X})^2 = E\tilde{X}^2$ ,

$$E U(w + \tilde{X}) \approx U(w) + U'(w)E\tilde{X} + \frac{1}{2}U''(w)E\tilde{X}^2 = U(w) + \frac{1}{2}U''(w)\text{var } \tilde{X}$$

- Similarly, for small premium, one can use Taylor expansion

$$U(w - \pi) \approx U(w) - \pi U'(w)$$

- Substitute into the risk premium formula (1)  $U(w - \pi) = E U(w + \tilde{X})$ ,

$$U(w) - \pi U'(w) = U(w) + \frac{1}{2}U''(w)\text{var } \tilde{X} \implies \pi = \frac{1}{2} \left( -\frac{U''(w)}{U'(w)} \right) \text{var } \tilde{X}$$

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)



# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)
- The term  $-\frac{U''(w)}{U'(w)} \cdot w$  is the coefficient of relative risk aversion (RRA)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)
- The term  $-\frac{U''(w)}{U'(w)} \cdot w$  is the coefficient of relative risk aversion (RRA)
  - Measures risk aversion relative to wealth level

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)
- The term  $-\frac{U''(w)}{U'(w)} \cdot w$  is the coefficient of relative risk aversion (RRA)
  - Measures risk aversion relative to wealth level
  - Logarithmic utility:  $RRA = 1$  (constant)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)
- The term  $-\frac{U''(w)}{U'(w)} \cdot w$  is the coefficient of relative risk aversion (RRA)
  - Measures risk aversion relative to wealth level
  - Logarithmic utility:  $RRA = 1$  (constant)
  - Power utility:  $RRA = \gamma$  (constant)

# Risk Aversion and Risk Premium - III

- The term  $-\frac{U''(w)}{U'(w)}$  is the coefficient of absolute risk aversion (ARA)
  - Measures risk aversion in absolute dollar terms
  - Logarithmic utility:  $ARA = \frac{1}{w}$  (decreasing with wealth)
  - Power utility:  $ARA = \frac{\gamma}{w}$  (decreasing with wealth)
  - Exponential utility:  $ARA = \gamma$  (constant regardless of wealth)
- The term  $-\frac{U''(w)}{U'(w)} \cdot w$  is the coefficient of relative risk aversion (RRA)
  - Measures risk aversion relative to wealth level
  - Logarithmic utility:  $RRA = 1$  (constant)
  - Power utility:  $RRA = \gamma$  (constant)
  - Exponential utility:  $RRA = \gamma w$  (increasing with wealth)

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both



# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$
  - **Independence:** For any lotteries  $L_1, L_2, L_3$  and any probability  $p \in (0, 1]$ ,  $L_1 \succeq L_2$  if and only if  $pL_1 + (1 - p)L_3 \succeq pL_2 + (1 - p)L_3$

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$
  - **Independence:** For any lotteries  $L_1, L_2, L_3$  and any probability  $p \in (0, 1]$ ,  $L_1 \succeq L_2$  if and only if  $pL_1 + (1 - p)L_3 \succeq pL_2 + (1 - p)L_3$
- These axioms lead to the expected utility representation:

$$L_1 \succeq L_2 \iff E_{L_1}[U(x)] \geq E_{L_2}[U(x)]$$

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$
  - **Independence:** For any lotteries  $L_1, L_2, L_3$  and any probability  $p \in (0, 1]$ ,  $L_1 \succeq L_2$  if and only if  $pL_1 + (1 - p)L_3 \succeq pL_2 + (1 - p)L_3$
- These axioms lead to the expected utility representation:

$$L_1 \succeq L_2 \iff E_{L_1}[U(x)] \geq E_{L_2}[U(x)]$$

- The independence axiom is particularly important and controversial

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$
  - **Independence:** For any lotteries  $L_1, L_2, L_3$  and any probability  $p \in (0, 1]$ ,  $L_1 \succeq L_2$  if and only if  $pL_1 + (1 - p)L_3 \succeq pL_2 + (1 - p)L_3$
- These axioms lead to the expected utility representation:

$$L_1 \succeq L_2 \iff E_{L_1}[U(x)] \geq E_{L_2}[U(x)]$$

- The independence axiom is particularly important and controversial
  - It states that preferences between lotteries should not be affected by mixing them with a third lottery

# Axiomatic Foundation of Expected Utility

- John von Neumann and Oskar Morgenstern (1947) provided axioms for expected utility
- Four key axioms:
  - **Completeness:** For any lotteries  $L_1$  and  $L_2$ , either  $L_1 \succeq L_2$  or  $L_2 \succeq L_1$  or both
  - **Transitivity:** If  $L_1 \succeq L_2$  and  $L_2 \succeq L_3$ , then  $L_1 \succeq L_3$
  - **Continuity:** If  $L_1 \succeq L_2 \succeq L_3$ , then there exists a probability  $p \in [0, 1]$  such that  $L_2 \sim pL_1 + (1 - p)L_3$
  - **Independence:** For any lotteries  $L_1, L_2, L_3$  and any probability  $p \in (0, 1]$ ,  $L_1 \succeq L_2$  if and only if  $pL_1 + (1 - p)L_3 \succeq pL_2 + (1 - p)L_3$
- These axioms lead to the expected utility representation:

$$L_1 \succeq L_2 \iff E_{L_1}[U(x)] \geq E_{L_2}[U(x)]$$

- The independence axiom is particularly important and controversial
  - It states that preferences between lotteries should not be affected by mixing them with a third lottery
  - This axiom is violated in several famous paradoxes

# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$



# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} U(100) &> 0.33 \cdot U(101) + 0.66 \cdot U(100) + 0.01 \cdot U(0) \\ \implies 0.34 \cdot U(100) &> 0.33 \cdot U(101) + 0.01 \cdot U(0) \quad (2) \end{aligned}$$

# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} U(100) &> 0.33 \cdot U(101) + 0.66 \cdot U(100) + 0.01 \cdot U(0) \\ \implies 0.34 \cdot U(100) &> 0.33 \cdot U(101) + 0.01 \cdot U(0) \quad (2) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{prob. } 0.34 \\ 0 & \text{prob. } 0.66 \end{cases} \quad Y = \begin{cases} 101 & \text{prob. } 0.33 \\ 0 & \text{prob. } 0.67 \end{cases}$$

# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} U(100) &> 0.33 \cdot U(101) + 0.66 \cdot U(100) + 0.01 \cdot U(0) \\ \implies 0.34 \cdot U(100) &> 0.33 \cdot U(101) + 0.01 \cdot U(0) \quad (2) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{prob. } 0.34 \\ 0 & \text{prob. } 0.66 \end{cases} \quad Y = \begin{cases} 101 & \text{prob. } 0.33 \\ 0 & \text{prob. } 0.67 \end{cases}$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

# Allais Paradox

- Game A

$$X = \begin{cases} 101 & \text{prob. } 0.33 \\ 100 & \text{prob. } 0.66 \\ 0 & \text{prob. } 0.01 \end{cases} \quad Y = 100 \text{ with prob. } 1$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} U(100) &> 0.33 \cdot U(101) + 0.66 \cdot U(100) + 0.01 \cdot U(0) \\ \implies 0.34 \cdot U(100) &> 0.33 \cdot U(101) + 0.01 \cdot U(0) \quad (2) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{prob. } 0.34 \\ 0 & \text{prob. } 0.66 \end{cases} \quad Y = \begin{cases} 101 & \text{prob. } 0.33 \\ 0 & \text{prob. } 0.67 \end{cases}$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} 0.33 \cdot U(101) + 0.67 \cdot U(0) &> 0.34 \cdot U(100) + 0.66 \cdot U(0) \\ \implies 0.33 \cdot U(101) + 0.01 \cdot U(0) &> 0.34 \cdot U(100) \quad (3) \end{aligned}$$

# Ellsberg Paradox

- Given an urn with 30 balls of colors red, yellow, and black

# Ellsberg Paradox

- Given an urn with 30 balls of colors red, yellow, and black
- There are 10 red balls; total 20 yellow / black balls, but the number of each type unknown

# Ellsberg Paradox

- Given an urn with 30 balls of colors red, yellow, and black
- There are 10 red balls; total 20 yellow / black balls, but the number of each type unknown
- The agent estimates the probability of drawing yellow as  $p$  where  $0 < p < \frac{2}{3}$



# Ellsberg Paradox

- Given an urn with 30 balls of colors red, yellow, and black
- There are 10 red balls; total 20 yellow / black balls, but the number of each type unknown
- The agent estimates the probability of drawing yellow as  $p$  where  $0 < p < \frac{2}{3}$
- A single ball is drawn from the urn

# Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases}$$

$$Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

# Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

Mostly prefer  $X$  to  $Y$ : from the Expected Utility Hypothesis

# Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

Mostly prefer  $X$  to  $Y$ : from the Expected Utility Hypothesis

$$\begin{aligned} \frac{1}{3} \cdot U(100) + \frac{2}{3} \cdot U(0) &> p \cdot U(100) + (1 - p) \cdot U(0) \\ \implies \left(\frac{1}{3} - p\right) \cdot U(100) &> \left(\frac{1}{3} - p\right) \cdot U(0) \quad (4) \end{aligned}$$

# Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

Mostly prefer  $X$  to  $Y$ : from the Expected Utility Hypothesis

$$\begin{aligned} \frac{1}{3} \cdot U(100) + \frac{2}{3} \cdot U(0) &> p \cdot U(100) + (1 - p) \cdot U(0) \\ \implies \left(\frac{1}{3} - p\right) \cdot U(100) &> \left(\frac{1}{3} - p\right) \cdot U(0) \quad (4) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{if red or black} \\ 0 & \text{if yellow} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow or black} \\ 0 & \text{if red} \end{cases}$$

## Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

Mostly prefer  $X$  to  $Y$ : from the Expected Utility Hypothesis

$$\begin{aligned} \frac{1}{3} \cdot U(100) + \frac{2}{3} \cdot U(0) &> p \cdot U(100) + (1-p) \cdot U(0) \\ \implies \left(\frac{1}{3} - p\right) \cdot U(100) &> \left(\frac{1}{3} - p\right) \cdot U(0) \quad (4) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{if red or black} \\ 0 & \text{if yellow} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow or black} \\ 0 & \text{if red} \end{cases}$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

# Ellsberg Paradox (Cont'd)

- Game A

$$X = \begin{cases} 100 & \text{if red} \\ 0 & \text{if yellow or black} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow} \\ 0 & \text{if red or black} \end{cases}$$

Mostly prefer  $X$  to  $Y$ : from the Expected Utility Hypothesis

$$\begin{aligned} \frac{1}{3} \cdot U(100) + \frac{2}{3} \cdot U(0) &> p \cdot U(100) + (1-p) \cdot U(0) \\ \implies \left(\frac{1}{3} - p\right) \cdot U(100) &> \left(\frac{1}{3} - p\right) \cdot U(0) \quad (4) \end{aligned}$$

- Game B

$$X = \begin{cases} 100 & \text{if red or black} \\ 0 & \text{if yellow} \end{cases} \quad Y = \begin{cases} 100 & \text{if yellow or black} \\ 0 & \text{if red} \end{cases}$$

Mostly prefer  $Y$  to  $X$ : from the Expected Utility Hypothesis

$$\begin{aligned} \frac{2}{3} \cdot U(100) + \frac{1}{3} \cdot U(0) &> (1-p) \cdot U(100) + p \cdot U(0) \\ \implies \left(\frac{1}{3} - p\right) \cdot U(0) &> \left(\frac{1}{3} - p\right) \cdot U(100) \quad (5) \end{aligned}$$