

Recall: Mixed logit choice probability requires an integral

$$P_{ij}(X, Z; \beta, \mu, \sigma) = \int \frac{\exp(X_i(\beta_j - \beta_J) + \gamma_i(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma_i(Z_{ik} - Z_{iJ}))} f(\gamma_i; \mu, \sigma) d\gamma_i$$

Recall: Mixed logit choice probability requires an integral

$$P_{ij}(X, Z; \beta, \mu, \sigma) = \int \frac{\exp(X_i(\beta_j - \beta_J) + \gamma_i(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma_i(Z_{ik} - Z_{iJ}))} f(\gamma_i; \mu, \sigma) d\gamma_i$$

This integral has no closed form  $\rightarrow$  must approximate numerically

Recall: Mixed logit choice probability requires an integral

$$P_{ij}(X, Z; \beta, \mu, \sigma) = \int \frac{\exp(X_i(\beta_j - \beta_J) + \gamma_i(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma_i(Z_{ik} - Z_{iJ}))} f(\gamma_i; \mu, \sigma) d\gamma_i$$

This integral has no closed form  $\rightarrow$  must approximate numerically

**Classical approach:** Quadrature or simulation-based methods

Recall: Mixed logit choice probability requires an integral

$$P_{ij}(X, Z; \beta, \mu, \sigma) = \int \frac{\exp(X_i(\beta_j - \beta_J) + \gamma_i(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma_i(Z_{ik} - Z_{iJ}))} f(\gamma_i; \mu, \sigma) d\gamma_i$$

This integral has no closed form  $\rightarrow$  must approximate numerically

**Classical approach:** Quadrature or simulation-based methods

**Problem:** Maximization can be difficult (local maxima, slow convergence, ...)

Bayesian approach avoids maximization entirely

Instead of finding  $\max_{\theta} \ell(\theta)$ , calculate  $E[\theta|\text{data}]$

Bayesian approach avoids maximization entirely

Instead of finding  $\max_{\theta} \ell(\theta)$ , calculate  $E[\theta|\text{data}]$

**Key insight:** Treat individual  $\gamma_i$  as parameters alongside  $\mu, \sigma$

Bayesian approach avoids maximization entirely

Instead of finding  $\max_{\theta} \ell(\theta)$ , calculate  $E[\theta|\text{data}]$

**Key insight:** Treat individual  $\gamma_i$  as parameters alongside  $\mu, \sigma$

Use data augmentation via posterior distribution instead of integrating:

- Classical:  $\int L(y_i|\gamma_i)dF(\gamma_i; \mu, \sigma)$  inside likelihood
- Bayesian: Draw  $\gamma_i$  directly, condition on it

Bayesian approach avoids maximization entirely

Instead of finding  $\max_{\theta} \ell(\theta)$ , calculate  $E[\theta|\text{data}]$

**Key insight:** Treat individual  $\gamma_i$  as parameters alongside  $\mu, \sigma$

Use data augmentation via posterior distribution instead of integrating:

- Classical:  $\int L(y_i|\gamma_i)dF(\gamma_i; \mu, \sigma)$  inside likelihood
- Bayesian: Draw  $\gamma_i$  directly, condition on it

No simulation of choice probabilities needed  $\rightarrow$  just evaluate logit formula at drawn  $\gamma_i$



Bayesian Gibbs sampling is like EM with continuous types

Both treat  $\gamma_i$  as latent  $\rightarrow$  fill in  $\gamma_i$  instead of integrating it out

EM (you know this):

E-step:  $E[\gamma_i | y_i, \mu^{(m)}, \sigma^{(m)}]$

M-step: Maximize using expectations

$\rightarrow$  MLE

Bayesian Gibbs (new):

Draw  $\gamma_i \sim p(\gamma_i | y_i, \mu, \sigma)$

Draw  $\mu, \sigma \sim p(\mu, \sigma | \{\gamma_i\})$

$\rightarrow$  Posterior mean  $\approx$  MLE in limit

Bayesian Gibbs sampling is like EM with continuous types

Both treat  $\gamma_i$  as latent  $\rightarrow$  fill in  $\gamma_i$  instead of integrating it out

EM (you know this):

E-step:  $E[\gamma_i | y_i, \mu^{(m)}, \sigma^{(m)}]$

M-step: Maximize using expectations

$\rightarrow$  MLE

Bayesian Gibbs (new):

Draw  $\gamma_i \sim p(\gamma_i | y_i, \mu, \sigma)$

Draw  $\mu, \sigma \sim p(\mu, \sigma | \{\gamma_i\})$

$\rightarrow$  Posterior mean  $\approx$  MLE in limit

**Difference:** EM integrates analytically, Gibbs samples via MCMC

Bayesian procedure for mixed logit:

Gibbs sampling - iteratively draw from conditional posteriors:

Bayesian procedure for mixed logit:

Gibbs sampling - iteratively draw from conditional posteriors:

1. Draw each  $\gamma_i$  from posterior distribution  $p(\gamma_i | \mu, \sigma, y_i)$ 
  - Just evaluate logit likelihood at current  $\mu, \sigma$
  - No integral needed

Bayesian procedure for mixed logit:

Gibbs sampling - iteratively draw from conditional posteriors:

1. Draw each  $\gamma_i$  from posterior distribution  $p(\gamma_i | \mu, \sigma, y_i)$ 
  - Just evaluate logit likelihood at current  $\mu, \sigma$
  - No integral needed
2. Draw  $\mu$  from  $p(\mu | \sigma, \{\gamma_i\})$ 
  - Essentially: average of the  $\gamma_i$ 's

Bayesian procedure for mixed logit:

Gibbs sampling - iteratively draw from conditional posteriors:

1. Draw each  $\gamma_i$  from posterior distribution  $p(\gamma_i | \mu, \sigma, y_i)$ 
  - Just evaluate logit likelihood at current  $\mu, \sigma$
  - No integral needed
2. Draw  $\mu$  from  $p(\mu | \sigma, \{\gamma_i\})$ 
  - Essentially: average of the  $\gamma_i$ 's
3. Draw  $\sigma$  from  $p(\sigma | \mu, \{\gamma_i\})$ 
  - Essentially: variance of the  $\gamma_i$ 's around  $\mu$

Bayesian procedure for mixed logit:

Gibbs sampling - iteratively draw from conditional posteriors:

1. Draw each  $\gamma_i$  from posterior distribution  $p(\gamma_i|\mu, \sigma, y_i)$ 
  - Just evaluate logit likelihood at current  $\mu, \sigma$
  - No integral needed
2. Draw  $\mu$  from  $p(\mu|\sigma, \{\gamma_i\})$ 
  - Essentially: average of the  $\gamma_i$ 's
3. Draw  $\sigma$  from  $p(\sigma|\mu, \{\gamma_i\})$ 
  - Essentially: variance of the  $\gamma_i$ 's around  $\mu$

Repeat until convergence  $\rightarrow$  posterior mean of draws = estimates

How do we know the posterior distributions?

Bayes' rule:

$$p(\theta|\text{data}) \propto L(\text{data}|\theta) \times \pi(\theta)$$



How do we know the posterior distributions?

Bayes' rule:

$$p(\theta|\text{data}) \propto L(\text{data}|\theta) \times \pi(\theta)$$

For  $\gamma_i$ :

- $p(\gamma_i|\mu, \sigma, y_i) \propto L(y_i|\gamma_i) \times \pi(\gamma_i|\mu, \sigma)$
- No closed form  $\rightarrow$  sample via Metropolis-Hastings (accept/reject)

How do we know the posterior distributions?

Bayes' rule:

$$p(\theta|\text{data}) \propto L(\text{data}|\theta) \times \pi(\theta)$$

For  $\gamma_i$ :

- $p(\gamma_i|\mu, \sigma, y_i) \propto L(y_i|\gamma_i) \times \pi(\gamma_i|\mu, \sigma)$
- No closed form  $\rightarrow$  sample via Metropolis-Hastings (accept/reject)

For  $\mu, \sigma$ :

- Conjugate priors  $\rightarrow$  closed form posteriors
- $p(\mu|\sigma, \{\gamma_i\}) \sim \text{Normal}$
- $p(\sigma|\mu, \{\gamma_i\}) \sim \text{Inverse Wishart}$
- Direct sampling (no M-H needed)
- (though usually use M-H for all parameters)

Advantages of Bayesian approach:

Advantages of Bayesian approach:

- No maximization  $\rightarrow$  no convergence issues, starting values less critical

Advantages of Bayesian approach:

- No maximization  $\rightarrow$  no convergence issues, starting values less critical
- Number of simulation draws  $R$  need not grow with sample size

## Advantages of Bayesian approach:

- No maximization  $\rightarrow$  no convergence issues, starting values less critical
- Number of simulation draws  $R$  need not grow with sample size
- Naturally provides full posterior distribution of  $\gamma_i$  (not just point estimates)

## Advantages of Bayesian approach:

- No maximization  $\rightarrow$  no convergence issues, starting values less critical
- Number of simulation draws  $R$  need not grow with sample size
- Naturally provides full posterior distribution of  $\gamma_i$  (not just point estimates)
- Asymptotically equivalent to MLE (Bernstein-von Mises theorem)

## Advantages of Bayesian approach:

- No maximization  $\rightarrow$  no convergence issues, starting values less critical
- Number of simulation draws  $R$  need not grow with sample size
- Naturally provides full posterior distribution of  $\gamma_i$  (not just point estimates)
- Asymptotically equivalent to MLE (Bernstein-von Mises theorem)

## Disadvantages:

- Different convergence issue: to posterior distribution (burn-in)
- Slower for some specifications (fixed coefficients, bounded distributions)