

Stated probabilistic choice models

## Stated probabilistic choice models

- Now, we observe a probability of choosing each alternative

## Stated probabilistic choice models

- Now, we observe a probability of choosing each alternative
- What does this information represent? It represents the person's uncertainty

## Stated probabilistic choice models

- Now, we observe a probability of choosing each alternative
- What does this information represent? It represents the person's uncertainty
- Specifically, it represents **resolvable uncertainty**:

## Stated probabilistic choice models

- Now, we observe a probability of choosing each alternative
- What does this information represent? It represents the person's uncertainty
- Specifically, it represents **resolvable uncertainty**:
  - uncertainty about unspecified attributes or states of the world in which choices will ultimately be made

Example of resolvable uncertainty:

Example of resolvable uncertainty:

- Q. “Will you go for Mexican or Thai food on Friday night?”

Example of resolvable uncertainty:

- Q. “Will you go for Mexican or Thai food on Friday night?”
- A. “Well, I’m not sure what mood I’ll be in that day, but probably Mexican”



Example of resolvable uncertainty:

- Q. “Will you go for Mexican or Thai food on Friday night?”
- A. “Well, I’m not sure what mood I’ll be in that day, but probably Mexican”
- Q. “What do you mean by ‘probably’?”

Example of resolvable uncertainty:

- Q. "Will you go for Mexican or Thai food on Friday night?"
- A. "Well, I'm not sure what mood I'll be in that day, but probably Mexican"
- Q. "What do you mean by 'probably'?"
- A. "A 78% chance"

Example of resolvable uncertainty:

- Q. “Will you go for Mexican or Thai food on Friday night?”
- A. “Well, I’m not sure what mood I’ll be in that day, but probably Mexican”
- Q. “What do you mean by ‘probably’?”
- A. “A 78% chance”

If the person thinks there is no such uncertainty, then they can report  $p = 0$  or  $p = 1$

Estimating probabilistic choice models

Estimating probabilistic choice models

How do we proceed with estimation when  $y$  is itself a probability (not 0/1)?

Estimating probabilistic choice models

How do we proceed with estimation when  $y$  is itself a probability (not 0/1)?

We invert the logit formula

Estimating probabilistic choice models

How do we proceed with estimation when  $y$  is itself a probability (not 0/1)?

We invert the logit formula

Consider the binomial logit as an example:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2}) \gamma)}{1 + \exp((Z_{i1} - Z_{i2}) \gamma)}$$

Start with the logit probability:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2}) \gamma)}{1 + \exp((Z_{i1} - Z_{i2}) \gamma)}$$



Start with the logit probability:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2})\gamma)}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

And the complement:

$$1 - P_{i1} = \frac{1}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

Start with the logit probability:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2})\gamma)}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

And the complement:

$$1 - P_{i1} = \frac{1}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

Take the ratio (denominators cancel):

$$\frac{P_{i1}}{1 - P_{i1}} = \exp((Z_{i1} - Z_{i2})\gamma)$$

Start with the logit probability:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2})\gamma)}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

And the complement:

$$1 - P_{i1} = \frac{1}{1 + \exp((Z_{i1} - Z_{i2})\gamma)}$$

Take the ratio (denominators cancel):

$$\frac{P_{i1}}{1 - P_{i1}} = \exp((Z_{i1} - Z_{i2})\gamma)$$

Take logs:

$$\log\left(\frac{P_{i1}}{1 - P_{i1}}\right) = (Z_{i1} - Z_{i2})\gamma$$

Start with the logit probability:

$$P_{i1} = \frac{\exp((Z_{i1} - Z_{i2}) \gamma)}{1 + \exp((Z_{i1} - Z_{i2}) \gamma)}$$

And the complement:

$$1 - P_{i1} = \frac{1}{1 + \exp((Z_{i1} - Z_{i2}) \gamma)}$$

Take the ratio (denominators cancel):

$$\frac{P_{i1}}{1 - P_{i1}} = \exp((Z_{i1} - Z_{i2}) \gamma)$$

Take logs:

$$\log\left(\frac{P_{i1}}{1 - P_{i1}}\right) = (Z_{i1} - Z_{i2}) \gamma$$

Now we're in a world where we can (in principle) use OLS to estimate  $\gamma$ !

Stated probabilities provide more information than discrete choice or rank ordering

Stated probabilities provide more information than discrete choice or rank ordering

Consider the following responses for an individual:

Option	Discrete Choice	Rank Ordering	Stated Prob 1	Stated Prob 2
A	0	2	0.01	0.49
B	1	1	0.99	0.51
C	0	3	0.00	0.00

- The preference ordering is the same in each column

- The preference ordering is the same in each column
- But the implied preference intensity is much different in the last two columns



- The preference ordering is the same in each column
- But the implied preference intensity is much different in the last two columns
- A  $(0,1,0)$  discrete choice response corresponds to a  $(0,100,0)$  probability response

Measurement error

## Measurement error

- One concern with stated probabilities is measurement error

## Measurement error

- One concern with stated probabilities is measurement error
- Rather than report 99.5%, someone may just write 100%

## Measurement error

- One concern with stated probabilities is measurement error
- Rather than report 99.5%, someone may just write 100%
- But if  $p = 0$  or  $p = 1$ ,  $\log\left(\frac{P_{i1}}{1-P_{i1}}\right)$  is undefined!

## Measurement error

- One concern with stated probabilities is measurement error
- Rather than report 99.5%, someone may just write 100%
- But if  $p = 0$  or  $p = 1$ ,  $\log\left(\frac{P_{i1}}{1-P_{i1}}\right)$  is undefined!
- In this case, we have to recode 0s or 1s to be small values (e.g. .001, .999)

Then, to avoid contamination, we need to use LAD instead of OLS. New equation:

$$\log \left( \frac{\tilde{P}_{i1}}{1 - \tilde{P}_{i1}} \right) = (Z_{i1} - Z_{i2}) \gamma + \eta_{i1}$$

Then, to avoid contamination, we need to use LAD instead of OLS. New equation:

$$\log \left( \frac{\tilde{P}_{i1}}{1 - \tilde{P}_{i1}} \right) = (Z_{i1} - Z_{i2}) \gamma + \eta_{i1}$$

where  $\tilde{P}$  is the recoded probability and  $\eta_{i1}$  is the difference in measurement errors



Estimation with  $J > 2$

Estimation with  $J > 2$

- The above is the equation for a 2-option choice set

Estimation with  $J > 2$

- The above is the equation for a 2-option choice set
- With more than 2 options, we have more observations per scenario

Estimation with  $J > 2$

- The above is the equation for a 2-option choice set
- With more than 2 options, we have more observations per scenario

If  $J = 3$ , we have:

$$\log \left( \frac{\tilde{P}_{i1}}{\tilde{P}_{i2}} \right) = (Z_{i1} - Z_{i2}) \gamma + \eta_{i1}$$

$$\log \left( \frac{\tilde{P}_{i3}}{\tilde{P}_{i2}} \right) = (Z_{i3} - Z_{i2}) \gamma + \eta_{i3}$$

If we see  $N$  people each making  $T$  choices, then our data has  $NT(J - 1)$  rows:

If we see  $N$  people each making  $T$  choices, then our data has  $NT(J - 1)$  rows:

ID	Scenario	Alternative	Probability	$Z$	$Z_j - Z_B$	$\log(P_j/P_B)$
1	1	A	0.45	2.3	0.5	1.099
1	1	B	0.15	1.8	0.0	—
1	1	C	0.40	3.1	1.3	0.981
1	2	A	0.25	2.5	-0.4	-0.875
1	2	B	0.60	2.9	0.0	—
1	2	C	0.15	1.7	-1.2	-1.386
2	1	A	0.10	3.2	-0.9	-1.946
2	1	B	0.70	4.1	0.0	—
2	1	C	0.20	2.8	-1.3	-1.253

If we see  $N$  people each making  $T$  choices, then our data has  $NT(J - 1)$  rows:

ID	Scenario	Alternative	Probability	$Z$	$Z_j - Z_B$	$\log(P_j/P_B)$
1	1	A	0.45	2.3	0.5	1.099
1	1	B	0.15	1.8	0.0	—
1	1	C	0.40	3.1	1.3	0.981
1	2	A	0.25	2.5	-0.4	-0.875
1	2	B	0.60	2.9	0.0	—
1	2	C	0.15	1.7	-1.2	-1.386
2	1	A	0.10	3.2	-0.9	-1.946
2	1	B	0.70	4.1	0.0	—
2	1	C	0.20	2.8	-1.3	-1.253

Drop the rows for alternative B before estimating LAD regression models