$$\sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) = \sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) \quad \forall j, j'$$

$$\sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) = \sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) \quad \forall j, j'$$

In this case, choosing option k "renews" the state space, i.e.

$$\sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) = \sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) \quad \forall j, j'$$

In this case, choosing option k "renews" the state space, i.e.

Choosing k in t+1 removes any dependence of the period-t choice on the t+2 states

$$\sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) = \sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) \quad \forall j, j'$$

In this case, choosing option k "renews" the state space, i.e.

Choosing k in t+1 removes any dependence of the period-t choice on the t+2 states

We can use renewal, along with clever normalizations, to eliminate recursion

$$\sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) = \sum_{X_{t+1}} f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) \quad \forall j, j'$$

In this case, choosing option k "renews" the state space, i.e.

Choosing k in t+1 removes any dependence of the period-t choice on the t+2 states

We can use renewal, along with clever normalizations, to eliminate recursion

... Even when our choice set does not contain a terminal option

Why? Because CCPs don't have to correspond to optimal decisions

Normalize future value term relative to renewal action for choice j:

$$v_{jt}(X_t) = u_j(X_t) + \beta \sum_{X_{t+1}} \left[v_{kt+1}(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_j(X_{t+1}|X_t) + \beta c$$

Normalize future value term relative to renewal action for choice *j*:

$$v_{jt}(X_t) = u_j(X_t) + \beta \sum \left[v_{kt+1}(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_j(X_{t+1}|X_t) + \beta c$$

Expanding $v_{kt+1}(X_{t+1})$:

$$v_{kt+1}(X_{t+1}) = u_k(X_{t+1}) + \beta \sum_{t} V_{t+2}(X_{t+2}) f_k(X_{t+2}|X_{t+1})$$

Normalize future value term relative to renewal action for choice j:

$$v_{jt}(X_t) = u_j(X_t) + \beta \sum_{k} \left[v_{kt+1}(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_j(X_{t+1}|X_t) + \beta c$$

Expanding $v_{kt+1}(X_{t+1})$:

$$v_{kt+1}(X_{t+1}) = u_k(X_{t+1}) + \beta \sum_{x} V_{t+2}(X_{t+2}) f_k(X_{t+2}|X_{t+1})$$

Substituting in yields:

$$v_{jt}(X_t) = u_j(X_t) + \beta \sum_{X_{t+1}} \left[u_k(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_j(X_{t+1}|X_t) + \beta^2 \sum_{X_{t+1}} \sum_{X_{t+2}} V_{t+2}(X_{t+2}) f_k(X_{t+2}|X_{t+1}) f_j(X_{t+1}|X_t) + \beta c$$

For choice i', normalize FV term relative to choice k:

$$v_{j't}(X_t) = u_{j'}(X_t) + \beta \sum_{X_{t+1}} [u_k(X_{t+1}) - \log(p_{kt+1}(X_{t+1}))] f_{j'}(X_{t+1}|X_t) + \beta^2 \sum_{X_{t+2}} \sum_{X_{t+2}} V_{t+2}(X_{t+2}) f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) + \beta c$$

 $X_{t+1} X_{t+2}$

For choice j', normalize FV term relative to choice k:

$$\begin{aligned} v_{j't}(X_t) &= u_{j'}(X_t) + \beta \sum_{X_{t+1}} \left[u_k(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_{j'}(X_{t+1}|X_t) + \\ \beta^2 \sum_{X_{t+1}} \sum_{X_{t+2}} V_{t+2}(X_{t+2}) f_k(X_{t+2}|X_{t+1}) f_{j'}(X_{t+1}|X_t) + \beta c \end{aligned}$$

Renewal property: $V_{t+2}(X_{t+2})$ terms cancel out in differences

$$v_{jt}(X_t) - v_{j't}(X_t) = u_j(X_t) - u_{j'}(X_t)$$

$$+ \beta \sum_{X_{t+1}} \left[u_k(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_j(X_{t+1}|X_t)$$

$$- \beta \sum_{X_{t+1}} \left[u_k(X_{t+1}) - \log(p_{kt+1}(X_{t+1})) \right] f_{j'}(X_{t+1}|X_t)$$

Let's revisit the Rust (1987) bus engine model—two choices with flow payoffs:

$$u(X_t, d_t, \theta) = \begin{cases} -c(X_t, \theta) & \text{if } d_t = 0\\ -\overline{P} - P + c(0, \theta) & \text{if } d_t = 1 \end{cases}$$

Let's revisit the Rust (1987) bus engine model—two choices with flow payoffs:

$$u(X_t, d_t, \theta) = \begin{cases} -c(X_t, \theta) & \text{if } d_t = 0\\ -\overline{|P|} - P + c(0, \theta) & \text{if } d_t = 1 \end{cases}$$

Replacing the engine sets $X_{t+1}=0$ regardless of $X_t\implies j=1$ is renewal

Only need one-period-ahead replacement probability for future value

$$v_1(X) = u_1(X) + \beta [v_1(0) - \log(p_1(0))] + \beta c$$
 (replace)
 $v_0(X) = u_0(X) + \beta \sum_{X'} [v_1(X') - \log(p_1(X'))] f(X'|X) + \beta c$ (maintain)

$$v_1(X) = u_1(X) + \beta [v_1(0) - \log(p_1(0))] + \beta c$$
 (replace)
 $v_0(X) = u_0(X) + \beta \sum_{X'} [v_1(X') - \log(p_1(X'))] f(X'|X) + \beta c$ (maintain)

Key property: $v_1(X) = v_1(0)$ for all X because:

$$v_1(X) = u_1(X) + \beta [v_1(0) - \log(p_1(0))] + \beta c$$
 (replace)
 $v_0(X) = u_0(X) + \beta \sum_{X'} [v_1(X') - \log(p_1(X'))] f(X'|X) + \beta c$ (maintain)

Key property: $v_1(X) = v_1(0)$ for all X because:

1. Renewal: Next period's state is always 0 after replacement \implies continuation value $\beta[v_1(0) - \log(p_1(0))]$ is identical for all X

$$v_1(X) = u_1(X) + \beta \left[v_1(0) - \log(p_1(0)) \right] + \beta c$$
 (replace)
$$v_0(X) = u_0(X) + \beta \sum_{X'} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X) + \beta c$$
 (maintain)

Key property: $v_1(X) = v_1(0)$ for all X because:

- 1. Renewal: Next period's state is always 0 after replacement \implies continuation value $\beta[v_1(0) \log(p_1(0))]$ is identical for all X
- 2. Rust's specification: $u_1(X)$ is constant in X

$$v_1(X) = u_1(X) + \beta \left[v_1(0) - \log(p_1(0)) \right] + \beta c$$
 (replace)
 $v_0(X) = u_0(X) + \beta \sum_{X'} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X) + \beta c$ (maintain)

Key property: $v_1(X) = v_1(0)$ for all X because:

- 1. Renewal: Next period's state is always 0 after replacement \implies continuation value $\beta[v_1(0) \log(p_1(0))]$ is identical for all X
- 2. Rust's specification: $u_1(X)$ is constant in X

Together: both current flow utility and all future values are independent of X

$$\begin{aligned} v_1(X) - v_0(X) &= u_1(X) - u_0(X) \\ &+ \beta \left[v_1(0) - \log(p_1(0)) \right] \\ &- \beta \sum_{X'} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X) \end{aligned}$$

$$egin{aligned} v_1(X) - v_0(X) &= u_1(X) - u_0(X) \ &+ eta \left[v_1(0) - \log(p_1(0))
ight] \ &- eta \sum \left[\left. v_1(X') \right. - \log(p_1(X'))
ight] f(X'|X) \end{aligned}$$

 $= u_1(X) - u_0(X) + \beta v_1(0) - \beta \log(p_1(0))$

$$-\beta\sum_{X'}\left[\begin{array}{c}v_1(X')\\\end{array}-\log(p_1(X'))\right]f(X'|X)$$
 Since $v_1(X')=v_1(0)$ for all X' :

 $-\beta v_1(0) \sum_{X'} f(X'|X) + \beta \sum_{X'} \log(p_1(X')) f(X'|X)$

$$egin{aligned} v_1(X) - v_0(X) &= u_1(X) - u_0(X) \ &+ eta \left[v_1(0) - \log(p_1(0))
ight] \ &- eta \sum \left[v_1(X') - \log(p_1(X'))
ight] f(X'|X) \end{aligned}$$

$$-\beta \sum_{X'} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X)$$
 Since $v_1(X') = v_1(0)$ for all X' :

 $= u_1(X) - u_0(X) + \beta v_1(0) - \beta \log(p_1(0))$

 $-\beta v_1(0)\sum_{X'}f(X'|X)+\beta\sum_{X'}\log(p_1(X'))f(X'|X)$

$$v_1(X) - v_0(X) = u_1(X) - u_0(X)$$

 $+ \beta \left[v_1(0) - \log(p_1(0)) \right]$
 $- \beta \sum_{X \in \mathcal{X}} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X)$

Since $v_1(X') = v_1(0)$ for all X':

$$= u_1(X) - u_0(X) + \beta v_1(0) - \beta \log(p_1(0))$$
$$-\beta v_1(0) \sum_{X'} f(X'|X) + \beta \sum_{X'} \log(p_1(X')) f(X'|X)$$

The $\beta v_1(0)$ terms cancel (since $\sum_{X'} f(X'|X) = 1$):

$$v_1(X) - v_0(X) = u_1(X) - u_0(X) + \beta \sum_{X'} \left[\log(p_1(X')) - \log(p_1(0)) \right] f(X'|X)$$

$$v_1(X) - v_0(X) = u_1(X) - u_0(X) + \beta \left[v_1(0) - \log(p_1(0)) \right] - \beta \sum_{X'} \left[v_1(X') - \log(p_1(X')) \right] f(X'|X)$$

Since $v_1(X') = v_1(0)$ for all X':

$$= u_1(X) - u_0(X) + \beta v_1(0) - \beta \log(p_1(0)) - \beta v_1(0) \sum_{XX} f(X'|X) + \beta \sum_{XX} \log(p_1(X')) f(X'|X)$$

The $\beta v_1(0)$ terms cancel (since $\sum_{X'} f(X'|X) = 1$):

$$v_1(X) - v_0(X) = u_1(X) - u_0(X) + \beta \sum_{X'} \left[\log(p_1(X')) - \log(p_1(0)) \right] f(X'|X)$$

Estimation: logit with adjustment term, calculate p_1 and f(X'|X) in first stage