

Suppose our utility function is given by

$$X_i\beta_j + \gamma_i Z_{ij} + \epsilon_{ij}$$

where we assume that $\gamma_i \sim F$ with pdf f and distributional parameters μ and σ

Suppose our utility function is given by

$$X_i\beta_j + \gamma_i Z_{ij} + \epsilon_{ij}$$

where we assume that $\gamma_i \sim F$ with pdf f and distributional parameters μ and σ

Then the logit choice probabilities become:

$$P_{ij}(X, Z; \beta, \mu, \sigma) = \int \frac{\exp(X_i(\beta_j - \beta_J) + \gamma_i(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma_i(Z_{ik} - Z_{iJ}))} f(\gamma_i; \mu, \sigma) d\gamma_i$$

This is just like the expected value of a function of a random variable W :

$$\mathbb{E}[g(W)] = \int g(W)f(W; \mu, \sigma) dW$$

This is just like the expected value of a function of a random variable W :

$$\mathbb{E}[g(W)] = \int g(W) f(W; \mu, \sigma) dW$$

Annoyance: the log likelihood now has an integral inside the log!

$$\ell(X, Z; \beta, \gamma, \mu, \sigma) = \sum_{i=1}^N \log \left\{ \int \prod_j \left[\frac{\exp(X_i(\beta_j - \beta_J) + \gamma(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma(Z_{ik} - Z_{iJ}))} \right]^{d_{ij}} f(\gamma; \mu, \sigma) d\gamma \right\}$$

This is just like the expected value of a function of a random variable W :

$$\mathbb{E}[g(W)] = \int g(W)f(W; \mu, \sigma) dW$$

Annoyance: the log likelihood now has an integral inside the log!

$$\ell(X, Z; \beta, \gamma, \mu, \sigma) = \sum_{i=1}^N \log \left\{ \int \prod_j \left[\frac{\exp(X_i(\beta_j - \beta_J) + \gamma(Z_{ij} - Z_{iJ}))}{\sum_k \exp(X_i(\beta_k - \beta_J) + \gamma(Z_{ik} - Z_{iJ}))} \right]^{d_{ij}} f(\gamma; \mu, \sigma) d\gamma \right\}$$

Can't switch log and integral because of properties of logarithms

Common mixing distributions:

Common mixing distributions:

- Normal

Common mixing distributions:

- Normal
- Log-normal

Common mixing distributions:

- Normal
- Log-normal
- Uniform

Common mixing distributions:

- Normal
- Log-normal
- Uniform
- Triangular

Common mixing distributions:

- Normal
- Log-normal
- Uniform
- Triangular
- Can also specify a multivariate normal

Common mixing distributions:

- Normal
- Log-normal
- Uniform
- Triangular
- Can also specify a multivariate normal
 - This would allow, e.g., heterogeneity in γ to be correlated with β

With the integral inside the log, estimation of the mixed logit is intensive

With the integral inside the log, estimation of the mixed logit is intensive

To estimate the likelihood function, need to numerically approximate the integral

With the integral inside the log, estimation of the mixed logit is intensive

To estimate the likelihood function, need to numerically approximate the integral

The most common way of doing this is [quadrature](#)

With the integral inside the log, estimation of the mixed logit is intensive

To estimate the likelihood function, need to numerically approximate the integral

The most common way of doing this is [quadrature](#)

Another common way of doing this is by [simulation](#) (Monte Carlo integration)

With the integral inside the log, estimation of the mixed logit is intensive


To estimate the likelihood function, need to numerically approximate the integral

The most common way of doing this is **quadrature**


Another common way of doing this is by **simulation** (Monte Carlo integration)

Choice is often determined by specifics of the problem at hand (e.g. dimensionality)

Riemann Sum vs. Quadrature




Riemann Sum vs. Quadrature




Riemann sum: $w_k = \Delta\gamma$ for all k (uniform widths)

Quadrature: optimal weights w_k (not uniform, chosen for accuracy)

Monte Carlo Integration




Monte Carlo Integration



Monte Carlo: uniform weights $w_r = \frac{1}{R}$ for all random draws

Draw R random values γ_r from $F(\cdot)$, take average of function values

Monte Carlo Integration



Monte Carlo: uniform weights $w_r = \frac{1}{R}$ for all random draws

Draw R random values γ_r from $F(\cdot)$, take average of function values

R usually needs to be very large (min. 10,000)