

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

- Utility varies over time:  $u_{jt}(X_t)$

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

- Utility varies over time:  $u_{jt}(X_t)$
- State transitions vary over time:  $f_{jt}(X_{t+1}|X_t)$

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

- Utility varies over time:  $u_{jt}(X_t)$
- State transitions vary over time:  $f_{jt}(X_{t+1}|X_t)$
- Decisions made until period  $T$

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

- Utility varies over time:  $u_{jt}(X_t)$
- State transitions vary over time:  $f_{jt}(X_{t+1}|X_t)$
- Decisions made until period  $T$
- Observations only until period  $\mathcal{T} < T$

Key advantage of finite dependence: no need to make assumptions far into lifecycle

Setup:

- Utility varies over time:  $u_{jt}(X_t)$
- State transitions vary over time:  $f_{jt}(X_{t+1}|X_t)$
- Decisions made until period  $T$
- Observations only until period  $\mathcal{T} < T$

Question: What remains identifiable?

With terminal or renewal choices, can identify  $f_{jt}$ 's and  $u_{jt}$ 's until period  $\mathcal{T} - 1$

With terminal or renewal choices, can identify  $f_{jt}$ 's and  $u_{jt}$ 's until period  $\mathcal{T} - 1$

Extreme value case:

$$v_j(X_t) = u_{jt}(X_t) + \beta \sum_{X_{t+1}} \log \left( \sum_k \exp[v_{kt+1}(X_{t+1})] \right) f_{jt}(X_{t+1}|X_t) + \beta c$$



With terminal or renewal choices, can identify  $f_{jt}$ 's and  $u_{jt}$ 's until period  $\mathcal{T} - 1$

Extreme value case:

$$\begin{aligned} v_j(X_t) &= u_{jt}(X_t) + \beta \sum_{X_{t+1}} \log \left( \sum_k \exp[v_{kt+1}(X_{t+1})] \right) f_{jt}(X_{t+1}|X_t) + \beta c \\ &= u_{jt}(X_t) + \beta \sum_{X_{t+1}} [v_{1t+1}(X_{t+1}) - \log(p_{1t+1}(X_{t+1}))] f_{jt}(X_{t+1}|X_t) + \beta c \end{aligned}$$

With terminal or renewal choices, can identify  $f_{jt}$ 's and  $u_{jt}$ 's until period  $\mathcal{T} - 1$

Extreme value case:

$$\begin{aligned} v_j(X_t) &= u_{jt}(X_t) + \beta \sum_{X_{t+1}} \log \left( \sum_k \exp[v_{kt+1}(X_{t+1})] \right) f_{jt}(X_{t+1}|X_t) + \beta c \\ &= u_{jt}(X_t) + \beta \sum_{X_{t+1}} [v_{1t+1}(X_{t+1}) - \log(p_{1t+1}(X_{t+1}))] f_{jt}(X_{t+1}|X_t) + \beta c \end{aligned}$$

Last term differences out or is constant

Expression holds *regardless* of expectations at  $t + 2$

When finite dependence takes more than one period:

When finite dependence takes more than one period:

- Transition functions can vary over time

When finite dependence takes more than one period:

- Transition functions can vary over time
- Utility function needs to be stable

When finite dependence takes more than one period:

- Transition functions can vary over time
- Utility function needs to be stable
- Reason: future flow payoffs appear in expectation of future utility, but we lack corresponding data to recover them

When finite dependence takes more than one period:

- Transition functions can vary over time
- Utility function needs to be stable
- Reason: future flow payoffs appear in expectation of future utility, but we lack corresponding data to recover them
- Implication: possible to estimate non-stationary games

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$



If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$
- Recall:  $V_{t+1} = \log G$ , where  $G = \sum_k \exp(\cdot)$  for T1EV

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$
- Recall:  $V_{t+1} = \log G$ , where  $G = \sum_k \exp(\cdot)$  for T1EV
- For nested logit: formula involves nesting parameters ( $\lambda_r$ 's)

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$
- Recall:  $V_{t+1} = \log G$ , where  $G = \sum_k \exp(\cdot)$  for T1EV
- For nested logit: formula involves nesting parameters ( $\lambda_r$ 's)

If  $\epsilon$ 's are Normal:

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$
- Recall:  $V_{t+1} = \log G$ , where  $G = \sum_k \exp(\cdot)$  for T1EV
- For nested logit: formula involves nesting parameters ( $\lambda_r$ 's)

If  $\epsilon$ 's are Normal:

- No closed-form expression for  $V_{t+1}$

If  $\epsilon$ 's are GEV, can still express  $V_{t+1}$  as closed-form function of  $p_{jt+1}$

But it can get complicated:

- Math depends on form of  $G$
- Recall:  $V_{t+1} = \log G$ , where  $G = \sum_k \exp(\cdot)$  for T1EV
- For nested logit: formula involves nesting parameters ( $\lambda_r$ 's)

If  $\epsilon$ 's are Normal:

- No closed-form expression for  $V_{t+1}$
- Need simulation to compute  $\mathbb{E}$  max integral

$$G_t = \sum_r \left( \sum_{j \in J_r} \exp \left( \frac{v_j(X_t)}{\lambda_r} \right) \right)^{\lambda_r}$$



$$G_t = \sum_r \left( \sum_{j \in J_r} \exp \left( \frac{v_j(X_t)}{\lambda_r} \right) \right)^{\lambda_r}$$

Probability of choosing  $j$  in nest  $J_r$ :

$$p_{jt}(X_t) = \frac{\left( \sum_{j' \in J_r} \exp \left( \frac{v_{j'}(X_t)}{\lambda_r} \right) \right)^{\lambda_r - 1} \exp \left( \frac{v_j(X_t)}{\lambda_r} \right)}{G_t}$$

$$G_t = \sum_r \left( \sum_{j \in J_r} \exp \left( \frac{v_j(X_t)}{\lambda_r} \right) \right)^{\lambda_r}$$

Probability of choosing  $j$  in nest  $J_r$ :

$$p_{jt}(X_t) = \frac{\left( \sum_{j' \in J_r} \exp \left( \frac{v_{j'}(X_t)}{\lambda_r} \right) \right)^{\lambda_r - 1} \exp \left( \frac{v_j(X_t)}{\lambda_r} \right)}{G_t}$$

Probability of choosing nest  $r$ :

$$p_{rt}(X_t) = \frac{\left( \sum_{j' \in J_r} \exp \left( \frac{v_{j'}(X_t)}{\lambda_r} \right) \right)^{\lambda_r}}{G_t}$$

From nest probability:

$$(G_t p_{rt}(X_t))^{1/\lambda_r} = \sum_{j' \in J_r} \exp\left(\frac{v_{j'}(X_t)}{\lambda_r}\right)$$

From nest probability:

$$(G_t p_{rt}(X_t))^{1/\lambda_r} = \sum_{j' \in J_r} \exp\left(\frac{v_{j'}(X_t)}{\lambda_r}\right)$$

Substitute into choice probability:

$$p_{jt}(X_t) = \frac{(G_t p_{rt}(X_t))^{\frac{\lambda_r-1}{\lambda_r}} \exp\left(\frac{v_j(X_t)}{\lambda_r}\right)}{G_t}$$

From nest probability:

$$(G_t p_{rt}(X_t))^{1/\lambda_r} = \sum_{j' \in J_r} \exp\left(\frac{v_{j'}(X_t)}{\lambda_r}\right)$$

Substitute into choice probability:

$$p_{jt}(X_t) = \frac{(G_t p_{rt}(X_t))^{\frac{\lambda_r-1}{\lambda_r}} \exp\left(\frac{v_j(X_t)}{\lambda_r}\right)}{G_t}$$

Simplify:

$$p_{jt}(X_t) = G_t^{\frac{-1}{\lambda_r}} p_{rt}(X_t)^{\frac{\lambda_r-1}{\lambda_r}} \exp\left(\frac{v_j(X_t)}{\lambda_r}\right)$$

Take logs and rearrange:

$$(1/\lambda_r) \log(G_t) = -\log(p_{jt}(X_t)) + ((\lambda_r - 1)/\lambda_r) \log(p_{rt}(X_t)) + (1/\lambda_r) v_j(X_t)$$

Take logs and rearrange:

$$(1/\lambda_r) \log(G_t) = -\log(p_{jt}(X_t)) + ((\lambda_r - 1)/\lambda_r) \log(p_{rt}(X_t)) + (1/\lambda_r) v_j(X_t)$$

Multiply through by  $\lambda_r$ :

$$\log(G_t) = -\lambda_r \log(p_{jt}(X_t)) + (\lambda_r - 1) \log(p_{rt}(X_t)) + v_j(X_t)$$

Take logs and rearrange:

$$(1/\lambda_r) \log(G_t) = -\log(p_{jt}(X_t)) + ((\lambda_r - 1)/\lambda_r) \log(p_{rt}(X_t)) + (1/\lambda_r) v_j(X_t)$$

Multiply through by  $\lambda_r$ :

$$\log(G_t) = -\lambda_r \log(p_{jt}(X_t)) + (\lambda_r - 1) \log(p_{rt}(X_t)) + v_j(X_t)$$

Since  $V_t(X_t) = \log(G_t) + c$ , we're done



Take logs and rearrange:

$$(1/\lambda_r) \log(G_t) = -\log(p_{jt}(X_t)) + ((\lambda_r - 1)/\lambda_r) \log(p_{rt}(X_t)) + (1/\lambda_r) v_j(X_t)$$

Multiply through by  $\lambda_r$ :

$$\log(G_t) = -\lambda_r \log(p_{jt}(X_t)) + (\lambda_r - 1) \log(p_{rt}(X_t)) + v_j(X_t)$$

Since  $V_t(X_t) = \log(G_t) + c$ , we're done

Note: when  $\lambda_r = 1$ , this reduces to multinomial logit

When  $\epsilon \sim \text{GEV}$ , analytic expressions for  $V_t(X_t)$  generally difficult

When  $\epsilon \sim \text{GEV}$ , analytic expressions for  $V_t(X_t)$  generally difficult

But: given particular  $X_t$ , can solve for  $V_t(X_t)$  numerically

When  $\epsilon \sim \text{GEV}$ , analytic expressions for  $V_t(X_t)$  generally difficult

But: given particular  $X_t$ , can solve for  $V_t(X_t)$  numerically

Recall:  $p_{jt}(X_t) = \frac{\partial \log(G_t)}{\partial v_j(X_t)}$

When  $\epsilon \sim \text{GEV}$ , analytic expressions for  $V_t(X_t)$  generally difficult

But: given particular  $X_t$ , can solve for  $V_t(X_t)$  numerically

Recall:  $p_{jt}(X_t) = \frac{\partial \log(G_t)}{\partial v_j(X_t)}$

Stack for all choices:

$$\begin{bmatrix} p_{1t}(X_t) \\ \vdots \\ p_{Jt}(X_t) \end{bmatrix} = \begin{bmatrix} \frac{\partial \log(G_t)}{\partial v_{1t}(X_t)} \\ \vdots \\ \frac{\partial \log(G_t)}{\partial v_{Jt}(X_t)} \end{bmatrix}$$

When  $\epsilon \sim \text{GEV}$ , analytic expressions for  $V_t(X_t)$  generally difficult

But: given particular  $X_t$ , can solve for  $V_t(X_t)$  numerically

Recall:  $p_{jt}(X_t) = \frac{\partial \log(G_t)}{\partial v_j(X_t)}$

Stack for all choices:

$$\begin{bmatrix} p_{1t}(X_t) \\ \vdots \\ p_{Jt}(X_t) \end{bmatrix} = \begin{bmatrix} \frac{\partial \log(G_t)}{\partial v_{1t}(X_t)} \\ \vdots \\ \frac{\partial \log(G_t)}{\partial v_{Jt}(X_t)} \end{bmatrix}$$

One equation redundant  $\Rightarrow J - 1$  system solving for  $J - 1$  differences in  $v$ 's