

The infinite horizon value function:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} V(X'_i) dF_j(X'_i|X_i)$$

The infinite horizon value function:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} V(X'_i) dF_j(X'_i|X_i)$$

Expanding the continuation value:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} E_{\epsilon'} \left(\max_k v_k(X'_i) + \epsilon'_{ik} \right) dF_j(X'_i|X_i)$$

For T1EV ϵ 's, the expectation of the maximum has a closed form:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} \log \left(\sum_{k=1}^J \exp[v_k(X'_i)] \right) dF_j(X'_i|X_i) + \beta c$$

For T1EV ϵ 's, the expectation of the maximum has a closed form:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} \log \left(\sum_{k=1}^J \exp[v_k(X'_i)] \right) dF_j(X'_i|X_i) + \beta c$$

Key insight: The v 's appear on both sides of the equation

This creates a system that requires solving for a fixed point

For T1EV ϵ 's, the expectation of the maximum has a closed form:

$$v_j(X_i) = u_j(X_i) + \beta \int_{X'_i} \log \left(\sum_{k=1}^J \exp[v_k(X'_i)] \right) dF_j(X'_i|X_i) + \beta c$$

Key insight: The v 's appear on both sides of the equation

This creates a system that requires solving for a fixed point

(This works because it is a contraction mapping)

Let \mathcal{X} denote the number of states X can take on

Stack the conditional value functions for each possible state and choice:

$$\begin{bmatrix} v_1(X_1) \\ v_1(X_2) \\ \vdots \\ v_1(X_{\mathcal{X}}) \\ \vdots \\ v_J(X_{\mathcal{X}}) \end{bmatrix} = \begin{bmatrix} u_1(X_1) + \beta \int_{X'} \log \left(\sum_{k=1}^J \exp[v_k(X')] \right) dF_1(X'|X_1) + \beta c \\ u_1(X_2) + \beta \int_{X'} \log \left(\sum_{k=1}^J \exp[v_k(X')] \right) dF_1(X'|X_2) + \beta c \\ \vdots \\ u_1(X_{\mathcal{X}}) + \beta \int_{X'} \log \left(\sum_{k=1}^J \exp[v_k(X')] \right) dF_1(X'|X_{\mathcal{X}}) + \beta c \\ \vdots \\ u_J(X_{\mathcal{X}}) + \beta \int_{X'} \log \left(\sum_{k=1}^J \exp[v_k(X')] \right) dF_J(X'|X_{\mathcal{X}}) + \beta c \end{bmatrix}$$

Let \mathcal{X} denote the number of states X can take on

Stack the conditional value functions for each possible state and choice:

This system of equations can be written compactly as:

$$\mathbf{v} = \mathbf{u} + \beta \mathbf{T}(\mathbf{v})$$

where $\mathbf{T}(\cdot)$ represents the expectation operator

Solution algorithm:

- Plug in values for the parameters and take a guess at the v 's

Solution algorithm:

- Plug in values for the parameters and take a guess at the v 's
- Substitute in for the v 's on the right hand side which gives us a new set of v 's

Solution algorithm:

- Plug in values for the parameters and take a guess at the v 's
- Substitute in for the v 's on the right hand side which gives us a new set of v 's
- Repeat until convergence

Solution algorithm:

- Plug in values for the parameters and take a guess at the v 's
- Substitute in for the v 's on the right hand side which gives us a new set of v 's
- Repeat until convergence

Formally:

Iterate

$$\mathbf{v}^{(n+1)} = \mathbf{u} + \beta \mathbf{T}(\mathbf{v}^{(n)})$$

until

$$\|\mathbf{v}^{(n+1)} - \mathbf{v}^{(n)}\| < \delta$$

The complete estimation algorithm has a similar nested structure as finite horizon case:

Outer loop (parameter estimation):

- Guess parameter values $\theta^{(0)}$ (utility coefficients, β , etc.)

The complete estimation algorithm has a similar nested structure as finite horizon case:

Outer loop (parameter estimation):

- Guess parameter values $\theta^{(0)}$ (utility coefficients, β , etc.)

Inner loop (value function solution):

- For given $\theta^{(m)}$, solve fixed point: $\mathbf{v}^{(n+1)} = \mathbf{u}(\theta^{(m)}) + \beta \mathbf{T}(\mathbf{v}^{(n)})$
- Continue until $\|\mathbf{v}^{(n+1)} - \mathbf{v}^{(n)}\| < \delta$

The complete estimation algorithm has a similar nested structure as finite horizon case:

Outer loop (parameter estimation):

- Guess parameter values $\theta^{(0)}$ (utility coefficients, β , etc.)

Inner loop (value function solution):

- For given $\theta^{(m)}$, solve fixed point: $\mathbf{v}^{(n+1)} = \mathbf{u}(\theta^{(m)}) + \beta \mathbf{T}(\mathbf{v}^{(n)})$
- Continue until $\|\mathbf{v}^{(n+1)} - \mathbf{v}^{(n)}\| < \delta$

Back to outer loop:

- Calculate choice probabilities using solved $\mathbf{v}(\theta^{(m)})$
- Evaluate likelihood $\mathcal{L}(\theta^{(m)})$
- Update parameters to $\theta^{(m+1)}$ using nonlinear optimizer (e.g. L-BFGS)
- Repeat until likelihood convergence