

Confidence intervals around counterfactuals

## Confidence intervals around counterfactuals

- The counterfactuals produce, e.g., a new predicted distribution of  $Y$

## Confidence intervals around counterfactuals

- The counterfactuals produce, e.g., a new predicted distribution of  $Y$
- But we don't know what the confidence intervals are around the prediction

## Confidence intervals around counterfactuals

- The counterfactuals produce, e.g., a new predicted distribution of  $Y$
- But we don't know what the confidence intervals are around the prediction
- Our estimates inherently come with sampling variation

## Confidence intervals around counterfactuals

- The counterfactuals produce, e.g., a new predicted distribution of  $Y$
- But we don't know what the confidence intervals are around the prediction
- Our estimates inherently come with sampling variation
- We should incorporate that variation into our counterfactual

## Confidence intervals around counterfactuals

- The counterfactuals produce, e.g., a new predicted distribution of  $Y$
- But we don't know what the confidence intervals are around the prediction
- Our estimates inherently come with sampling variation
- We should incorporate that variation into our counterfactual
- How do we do this? Bootstrapping

Bootstrapping to get CIs around counterfactuals

Bootstrapping to get CIs around counterfactuals

Recall:  $-H^{-1}$  is the variance matrix of our estimates, where  $H$  is the Hessian



Bootstrapping to get CIs around counterfactuals

Recall:  $-H^{-1}$  is the variance matrix of our estimates, where  $H$  is the Hessian

Assume that  $\hat{\theta} \sim MVN(\theta, -H^{-1})$

Bootstrapping to get CIs around counterfactuals

Recall:  $-H^{-1}$  is the variance matrix of our estimates, where  $H$  is the Hessian

Assume that  $\hat{\theta} \sim MVN(\theta, -H^{-1})$

- but remember that we want to minimize  $-\ell$  so we just use  $H^{-1}$

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals
- Then we look at the 2.5th and 97.5th percentiles of cfl statistics

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals
- Then we look at the 2.5th and 97.5th percentiles of cfl statistics
- This will represent a 95% confidence interval of that statistic

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals
- Then we look at the 2.5th and 97.5th percentiles of cfl statistics
- This will represent a 95% confidence interval of that statistic
- This algorithm is known as the **parametric bootstrap**

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals
- Then we look at the 2.5th and 97.5th percentiles of cfl statistics
- This will represent a 95% confidence interval of that statistic
- This algorithm is known as the **parametric bootstrap**
- Contrast with **nonparametric bootstrap** which is when we randomly re-sample data

- We sample  $B$  times from  $MVN(\theta, -H^{-1})$  and compute our new counterfactuals
- Then we look at the 2.5th and 97.5th percentiles of cfl statistics
- This will represent a 95% confidence interval of that statistic
- This algorithm is known as the **parametric bootstrap**
- Contrast with **nonparametric bootstrap** which is when we randomly re-sample data
- P.B. can be intensive if it is costly to conduct counterfactuals  $\rightarrow$  rarely used