

# The DESI PRObabilistic Value-Added Bright Galaxy Survey (PROVA-BGS) Mock Challenge

CHANGHOON HAHN,<sup>1, 2, \*</sup> GYUBIN KWON, MALGORZATA SIUDEK, RITA TOJEIRO, AND GQP WG

<sup>1</sup>*Lawrence Berkeley National Laboratory, 1 Cyclotron Rd, Berkeley CA 94720, USA*

<sup>2</sup>*Berkeley Center for Cosmological Physics, University of California, Berkeley CA 94720, USA*

## ABSTRACT

In this project, we present the methodology for inferring physical properties of galaxies from joint SED fitting of the DESI optical photometry and spectroscopy. We construct realistic forward modeled DESI data from star formation and chemical enrichment histories of galaxies in the L-GALAXIES semi-analytic model and the ILLUSTRIS TNG hydrodynamic simulations. Then, using these mock observations, we demonstrate that the stellar mass and SFR posteriors from our SED fitting are consistent with constraints from other SED fitting methods in the literature and, more importantly, the input stellar masses and SFRs from the simulations. The SED fitting we present and validate in this project will be used to construct the PRObabilistic Value-Added Bright Galaxy Survey (PROVABGS) from DESI observations.

*Keywords:* keyword1 – keyword2 – keyword3

## 1. INTRODUCTION

The Dark Energy Spectroscopic Instrument will conduct the largest spectroscopic galaxy survey to date covering  $\sim 14,000 \text{ deg}^2$  (?). Over the next five years,

**What is DESI?** Provide an overview of DESI specifics, numbers, and science goals, which will mostly be cosmology (BAO, RSD, etc). DESI will be great Beyond its impact on cosmology, DESI will also be transformative for galaxy science.

provide value-added galaxy catalogs (VAGCs), which will be transformative for galaxy science. In the past, VAGCs such as the MPA-JHU<sup>1</sup> provided galaxy properties from emission line analyses of SDSS spectra ?.

Similarly, the NYU-VAGC (?),

These catalogs have been indispensable for establishing the global statistical view of galaxy properties (see ?, for a review). **previous successes with value-added catalogs**


**what will PROVABGS be good for?** extending previous statistical analyses on SDSS to a larger sample and to higher redshift. Mention some obvious ones: stellar mass function, luminosity function, star-formation sequence. more detailed galaxy-halo connection mention some cosmological applica-

\* hahn.changhoon@gmail.com

<sup>1</sup> <https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/>

TODO

TODO  
TODO



**Figure 1.** DESI will conduct the largest spectroscopic survey to date covering  $\sim 14,000 \text{ deg}^2$ . During dark time, DESI will measure  $>20$  million spectra of luminous red galaxies, emission line galaxies, and quasars out to  $z > 3$ . During bright time, DESI will measure the spectra of  $\sim 10$  million galaxies out to  $z \sim 0.4$  with the Bright Galaxy Survey (BGS). *Left:* With its  $\sim 14,000 \text{ deg}^2$  footprint (black), DESI will cover  $\sim 2\times$  the SDSS footprint (blue;  $\sim 7000 \text{ deg}^2$ ) and  $\sim 45\times$  the GAMA footprint (orange;  $\sim 300 \text{ deg}^2$ ). *Right:* Over this footprint, the BGS will provide spectra for a magnitude limited sample of  $\sim 10$  million galaxies down to  $r < 20$ , two orders of magnitude deeper than the SDSS main galaxy sample and 0.2 mag deeper than GAMA.


tions: multi-tracer analyses. Furthermore, since BGS will measure galaxy spectra down to  $r \sim 20$ , we will have faint dwarf galaxies at low redshift. provabgs introduces a new frontier of low signal-to-noise statistically powerful sample that will require

**Why do we need a mock challenge?** We want to test and cement our methodology specifically for our GQP analysis before SV data comes out. As part of the survey preparation, we have all the tools to accurately forward model observations. **details on some of the specific tools and what we're able to simulate: realistic spectroscopy. realistic photometry. realistic spectro-photometry** All of this gives us a rare opportunity to test our methodology on bespoke simulations.

A mock challenge is also great for testing new methodology. BGS is a bright time survey and will push the boundaries of low SNR spectra. But if we can find a way to infer robust galaxy properties the statistical payout is awesome. **Something also about LRGs** We're also trying to robustly fit spectra and photometry simultaneously. This has been done before (**citations**) but not extensively tested on simulations.

## 2. SIMULATIONS

DESI, with its robotically-actuated and fibre-fed spectrographs, will collect 5000 spectra simultaneously. The spectra cover the wavelength range 3600 to 9800 Å, with a spectral resolution of  $R = \lambda/\Delta\lambda$  between 2000 and 5500. During its five-year operation, starting in 2020, it will measure over 30 million spectra over  $14,000 \text{ deg}^2$  of the sky (**cite**). In addition, these DESI targets also have optical and infrared imaging data from the DESI Legacy Imaging Surveys (hereafter Legacy Surveys



**Figure 2.** *Left:* We forward model DESI  $g$ ,  $r$ , and  $z$  photometry (red) for our simulated galaxies (Section 2.1) by convolving their SEDs (dotted) with their broadband filters (dashed) and then applying an empirical noise model based on BGS objects in LS (Section 2.3). *Right:* The  $g - r$  and  $r - z$  color distribution of the forward modeled LGAL photometry is in good agreement with the color distribution of LS BGS objects (black contours).

(Dey et al. 2019). The Legacy Surveys are a combination of three public projects (Dark Energy Camera Legacy Survey, Beijing-Arizona Sky Survey, and Mayall  $z$ -band Legacy Survey) that jointly imaged the  $\sim 14,000 \text{ deg}^2$  DESI footprint in three optical bands ( $g$ ,  $r$ , and  $z$ ). Furthermore, DR8 of the Legacy Survey also includes photometry in the WISE  $W1$ ,  $W2$ ,  $W3$ , and  $W4$  infrared bands. The infrared photometry is from all imaging through year 4 of NEOWISE-Reactivation force-photometered in the unWISE maps at the locations of Legacy Surveys optical sources (cite). Below we describe how we simulate realistic DESI-like galaxy spectra and photometry from state-of-the-art simulations.

TODO

### 2.1. *L-Galaxies*

brief overview of L-GALAXIES (hereafter LGAL; Henriques et al. 2015)

### 2.2. *Spectral Energy Distributions*

For each simulated galaxy, LGAL provides the star formation histories (SFHs) and chemical enrichment histories (ZH) for its bulge and disk components separately, in approximately log-spaced lookback time bins. We treat each lookback time bin,  $i$ , as a single stellar population (SSP) of age  $t_i$ . Then, we derive the luminosities of the bulge and disk components by summing up the luminosities of all of their SSPs:

$$L^{\text{comp.}}(\lambda) = \sum_i \text{SFH}_i^{\text{comp.}} \Delta t_i L_{\text{SSP}}(\lambda; t_i, Z_i^{\text{comp.}}). \quad (1)$$

$\text{SFH}_i^{\text{comp.}}$  and  $Z_i^{\text{comp.}}$  are the star formation rate and metallicity of the bulge or disk component in lookback time bin  $i$ .  $\Delta t_i$  is the width of the bin.  $L_{\text{SSP}}$  corresponds to the luminosity of the SSP, which we calculate using the Flexible Stellar Population Synthesis (FSPS Conroy et al. 2009; Conroy & Gunn 2010) model. For FSPS, we use the MIST isochrones (Paxton et al. 2011, 2013, 2015; Choi et al. 2016; Dotter 2016), the MILES spectral library (?), and the Chabrier (2003) initial mass function (IMF).

Next, we apply velocity dispersions to  $L^{\text{comp}}(\lambda)$ . For the disk, we apply a fixed  $50\text{km/s}$  velocity dispersion; for the bulge, we derive its velocity dispersion using the [Zahid et al. \(2016\)](#) empirical relation that depends on the total bulge mass. Then, we apply dust attenuation to stellar emission in the disk component ( $L^{\text{disk}}$ ) based on the cold gas content and orientation of the disk. We derive the attenuation curve using a mixed-screen model with the [Mathis \(1983\)](#) dust extinction curve. Stellar emission from stars younger than 30Myr are further attenuated with a uniform dust screen and a wavelength dependent optical depth. @rita further details and citations for the mixed-screen model No dust attenuation is applied to the bulge component. @rita how come?

TODO  
TODO

Finally, we combine the attenuated disk component and the bulge component to construct the total luminosity of the simulated galaxy and then convert this rest-frame luminosity to observed-frame SED flux using its redshift,  $z$ .

$$f_{\text{SED}}(\lambda) = \frac{A(\lambda)L^{\text{disk}}(\lambda) + L^{\text{bulge}}(\lambda)}{4\pi d_L(z)^2(1+z)}. \quad (2)$$

$A(\lambda)$  is the dust attenuation for the disk component described above and  $d_L(z)$  is the luminosity distance. In the left panel of Figure 2, we present an example of the SED flux constructed for an arbitrary LGALgalaxy (black).

### 2.3. Forward Modeling DESI Photometry


In this section, we describe how we construct realistic LS-like photometry from the SEDs of simulated galaxies described in the last section. First, we convolve the SEDs with the broadband filters of the Legacy Survey to generate broadband photometric fluxes:

$$f_X = \int f_{\text{SED}}(\lambda) R_X(\lambda) d\lambda \quad (3)$$

where  $f_{\text{SED}}$  is the galaxy SED (Eq. 2) and  $R_X$  is the transmission for filter  $X$ . We generate photometry for the  $g$ ,  $r$ , and  $z$  optical bands. Next, we apply realistic measurement uncertainties to the derived photometry using a simple empirical . We match each simulated galaxy to a BGS object from LS DR9 with the nearest  $r$ -band magnitude and  $g - r$  and  $r - z$  color. The photometric uncertainties ( $\sigma_X$ ) and  $r$ -band fiber flux ( $f_r^{\text{fiber}}$ ) of the BGS object are then assigned to the simulated galaxy. We sample a Gaussian distribution with standard deviation  $\sigma_X$  and apply it to construct realistic LS-like photometry:

$$\hat{f}_X = f_X + n_X \quad \text{where } n_X \sim \mathcal{N}(0, \sigma_X). \quad (4)$$

Finally, we impose the target selection criteria of BGS ([Ruiz-Macias et al. 2021](#), Hahn *et al.* in prep.). In the left panel of Figure 2, we overplot the forward modeled photometry (red) ontop of the SED flux (black) for an arbitrary LGALgalaxy. For reference, we also plot  $R_X$  for the  $g$ ,  $r$ , and  $z$  bands of the Legacy Survey in blue, orange, and green respectively. On the right panel, we compare the  $g - r$  versus  $r - z$  color distribution for the forward modeled LGALgalaxies (red) to the color distribution of BGS objects in LS (black contour). The forward modeled photometry show good agreement with LS BGS objects in color space.



**Figure 3.** We construct simulated DESI spectra (solid) for LGAL simulated galaxies by applying a fiber aperture correction to the SED (dashed) and a realistic DESI noise model. We apply a fiber aperture correction by scaling down the full SED (dotted) by the  $r$ -band fiber fraction derived from the Legacy Surveys imaging. The noise model accounts for the DESI spectrograph response and an atmosphere model that accounts for the bright time observing conditions of BGS. Our forward model produces DESI-like spectra for all three arms of the DESI spectrographs:  $b$ ,  $r$ , and  $z$  (blue, orange, and green, respectively). For more details, we refer readers to Section 2.4.

#### 2.4. Forward Modeling DESI Spectra

In this section, we describe how we construct realistic DESI-like spectroscopy from the SEDs of simulated galaxies. Our forward model involves modeling the fiber aperture effect and applying a noise model that accurately reproduces the bright time observations of BGS.

DESI uses fiber-fed spectrographs with fibers that have angular radii of  $1''$ . Only the light from a galaxy within this fiber aperture is collected by the instrument. LS provides measurements of photometric fiber flux within a  $1''$  radius aperture ( $f_X^{\text{fiber}}$ ), which estimates the flux that passed through to the fibers. When we assign photometric uncertainties to our simulated galaxies based on  $r$ ,  $g-r$ , and  $r-z$  in Section 2.3, we also assign  $r$ -band fiber flux. We model the SED flux that passes through the fiber by scaling the SED flux by the  $r$  band fiber fraction, the ratio of the  $r$ -band fiber flux over the total  $r$  band flux:

$$f^{\text{spec}}(\lambda) = \left( \frac{f_r^{\text{fiber}}}{f_r} \right) f_{\text{SED}}(\lambda). \quad (5)$$

This fiber aperture correction assumes that there is no significant color dependence. We also assume that there are no significant biases in the fiber flux measurements in LS due to miscentering of objects.

**Do we want to say more about this assumption? Also cite Marta's paper investigating aperture effect.**

In addition to using it for the aperture correction, we also use  $f_r^{\text{fiber}}$  to derive “measured”  $\hat{f}_r^{\text{fiber}}$ :

TODO

$$\hat{f}_r^{\text{fiber}} = f_r^{\text{fiber}} + n_r^{\text{fiber}} \quad \text{where } n_r^{\text{fiber}} \sim \mathcal{N}\left(0, \frac{f_r^{\text{fiber}}}{f_r} \sigma_r\right). \quad (6)$$

We later use  $\hat{f}_r^{\text{fiber}}$  to set the prior on the nuisance parameter of our SED modeling (Section 3).

Next, we apply a noise model that simulates the DESI instrument response and bright time observing conditions of BGS. We use the spectral simulation pipeline of **cite survey simulations**

TODO

[paper and list details](#)<sup>2</sup>. More specifically, we use nominal dark time observing conditions with 180s exposure time. These conditions accurately reproduce the spectral noise and redshift success rates of observed BGS exposures in DESI survey validation data (Hahn *et al.* in prep.). In Figure 2.3, we present the forward modeled BGS spectrum of a LGAL galaxy (solid). For reference, we include the full SED (dotted) and fiber fraction scaled SED (dashed) of the galaxy.

### 3. JOINT SED MODELING OF PHOTOMETRY AND SPECTRA

#### 3.1. *Stellar Population Synthesis Modeling*

PROVABGS will provide inferred galaxy properties derived from joint SED modeling of DESI photometry and spectra. For the SED modeling, we use a state-of-the-art stellar population synthesis (SPS) model that uses a non-parametric SFH with a star-burst, a non-parametric ZH that varies with time, and a flexible dust attenuation prescription.

The form of the SFH is one of the most important factors in the accuracy of an SPS model. In general, the form of the SFH requires balancing between being flexible enough to describe the wide range of SFHs in observations while not being too flexible that it can describe any SFH at the expense of constraining power. If the model SFH is not flexible enough to describe actual SFHs of galaxies, then unbiased galaxy properties cannot be inferred using the SPS model. For instance, most SPS models (*e.g.* CIGALE, [Serra et al. 2011](#); BAGPIPES, [Carnall et al. 2017](#)) use parametric SFH such as the exponentially declining  $\tau$ -model. Such functional forms, however, produce biased estimates of galaxy properties (*e.g.*  $M_*$  and SFR) when used to fit mock observations of simulated galaxies ([Simha et al. 2014](#); [?](#); [Carnall et al. 2018](#)). On the other hand, many non-parametric forms of the SFH are overly flexible and allow unphysical SFHs ([Leja et al. 2019](#)), which unnecessarily increases parameter degeneracies and discards constraining power.

In our SPS model, we use a non-parametric SFH with two components: one based on non-negative matrix factorization (NMF) basis functions and a starburst component. For the first component, SFH is a linear combination of NMF SFH bases:


$$\text{SFH}^{\text{NMF}}(t, t_{\text{age}}) = \sum_{i=1}^4 \beta_i \frac{s_i^{\text{SFH}}(t)}{\int_0^{t_{\text{age}}} s_i^{\text{SFH}}(t) dt}. \quad (7)$$

$\{s_i^{\text{SFH}}\}$  are the NMF basis functions and  $\{\beta_i\}$  are the coefficients. The integral in the denominator normalizes the NMF basis functions to unity. We constrain  $\sum_i \beta_i = 1$ , so the total SFH of the component over the age of the galaxy ( $t_{\text{age}}$ ) is normalized to unity.  $\{s_i^{\text{SFH}}\}$  are from Tojeiro *et al.* (in prep.) and derived from the IllustrisTNG cosmological hydrodynamic simulation ([Nelson et al. 2018](#); [Pillepich et al. 2018](#); [Springel et al. 2018](#)). The SFHs of simulated galaxies IllustrisTNG are compiled, rebinned, and smoothed [more details here](#). Afterwards, we perform non-negative matrix factorization ([Lee & Seung 1999](#); [Cichocki & Phan 2009](#); [Févotte & Idier 2011](#)) on the smooth SFHs to derive  $\{s_i^{\text{SFH}}\}$ . We find that 4 components is sufficient to accurately reconstruct the SFHs from IllustrisTNG.

TODO

<sup>2</sup> <https://specs.readthedocs.io>





**Figure 4.** Non-negative matrix factorization basis functions for the SFH (left) and ZH (right). These basis functions are derived from the SFHs and ZHs of simulated galaxies in the IllustrisTNG cosmological hydrodynamic simulations.

Assuming that the SFHs of IllustrisTNG galaxies resemble the SFHs of actual observed galaxies, our NMF form provides a compact and flexible representation of the SFHs.

The NMF basis functions are derived from smooth SFHs, which means that it does not include any stochasticity. However, observations and high resolution zoom-in hydrodynamical simulations both find significant stochasticity in galaxy SFHs (Sparre et al. 2017; Caplar & Tacchella 2019; Hahn et al. 2019; Iyer et al. 2020). To include this stochasticity in our SPS model, we include a starburst component that consists of a SSP in the SFH. For the total SFH, we use

$$\text{SFH}(t, t_{\text{age}}) = (1 - f_{\text{burst}}) \text{SFH}^{\text{NMF}}(t, t_{\text{age}}) + f_{\text{burst}} \delta_{\text{D}}(t - t_{\text{burst}}). \quad (8)$$

$f_{\text{burst}}$  is the fraction of total stellar mass formed during the starburst;  $t_{\text{burst}}$  is the time at which the starburst occurs;  $\delta_{\text{D}}$  is the Dirac delta function. In total we have 6 free parameters in our SFH: 4 NMF basis coefficients ( $\beta_i$ ),  $f_{\text{burst}}$ , and  $t_{\text{burst}}$ .

Another key part of an SPS model is the ZH, or chemical enrichment history. Current SPS models mostly assume a ZH that does not vary over time (Carnall et al. 2017; Leja et al. 2019). Since galaxies in hydrodynamical simulations and observations do not have constant metallicities throughout their history, this assumption can significantly bias the inferred galaxy properties. Instead, for our ZH, we take a similar approach to the SFH and use NMF basis functions:

$$\text{ZH}(t) = \sum_{i=1}^2 \gamma_i s_i^{\text{ZH}}(t). \quad (9)$$

$\{s_i^{\text{ZH}}(t)\}$  are the ZH NMF basis functions and  $\{\gamma_i\}$  are the coefficients.  $\{s_i^{\text{ZH}}(t)\}$  are fit using the ZHs of simulated galaxies from IllustrisTNG. We use two NMF components, so our ZH prescription has 2 free parameters.

Next, we use the SFH and ZH above to model the unattenuated rest-frame luminosity as a linear combination of multiple SSPs, evaluated at logarithmically-spaced lookback time bins. We use a fixed

log-binning with the bin edges starting with  $(0, 10^{6.05}\text{yr})$ ,  $(10^{6.05}, 10^{6.15}\text{yr})$ , and continuing on with bins of width 0.1 dex. The binning is truncated at the age of the model galaxy. For a  $z = 0$  galaxy, we use 43  $t_{\text{lookback}}$  bins. We use log-spaced  $t_{\text{lookback}}$  bins because it better reproduces galaxy luminosities evaluated with much higher resolution  $t_{\text{lookback}}$  binning than linearly-spacing, for the same number of bins. At every  $t_{\text{lookback}}$  bin  $i$ , we evaluate the luminosity of a SSP with  $\text{ZH}(t_i)$ , where  $t_i$  is the center of  $t_{\text{lookback}}$  bin, and total stellar mass calculated by resampling the SFH in Eq. 8. We use FSPSto evaluate the SSP luminosities and use the MIST isochrones, MILES spectral library, and the Chabrier (2003) IMF (same as in Section 2.2). Since we use MIST isochrones, we impose a minimum and maximum limit to ZH based on its coverage:  $4.49 \times 10^{-5}$  and  $4.49 \times 10^{-2}$ , respectively. Our stellar metallicity range is significantly broader than previous studies, *e.g.* Carnall et al. (2017); Leja et al. (2017); Tacchella et al. (2021).

Before we combine the SSP luminosities, we apply dust attenuation. We use a two component Charlot & Fall (2000) dust attenuation model with birth cloud (BC) and diffuse-dust (ISM) components. The BC component represents the extra dust attenuation of young stars that are embedded in molecular clouds and HII regions. For SSPs younger than  $t_i < 100\text{Myr}$ , we apply the following BC dust attenuation:

$$L_i(\lambda) = L_i^{\text{unatten.}}(\lambda) \exp \left[ -\tau_{\text{BC}} \left( \frac{\lambda}{5500\text{\AA}} \right)^{-0.7} \right]. \quad (10)$$

$\tau_{\text{BC}}$  is the BC optical depth that determines the strength of the BC attenuation. Afterwards, *all* SSPs are attenuated by the diffuse dust using the Kriek & Conroy (2013) attenuation curve parameterization:

$$L_i(\lambda) = L_i^{\text{unatten.}}(\lambda) \exp \left[ -\tau_{\text{ISM}} \left( \frac{\lambda}{5500\text{\AA}} \right)^{n_{\text{dust}}} (k_{\text{Cal}}(\lambda) + D(\lambda)) \right]. \quad (11)$$

$\tau_{\text{ISM}}$  is the diffuse dust optical depth.  $n_{\text{dust}}$  is the ? dust index, which determines the slope of the attenuation curve.  $k_{\text{Cal}}(\lambda)$  is the ? attenuation curve and  $D(\lambda)$  is the UV dust bump, parameterized using a Lorentzian-like Drude profile:

$$D(\lambda) = \frac{E_b(\lambda \Delta\lambda)^2}{(\lambda^2 - \lambda_0^2)^2 + (\lambda \Delta\lambda)^2} \quad (12)$$

where  $\lambda_0 = 2175\text{\AA}$ ,  $\Delta\lambda = 350\text{\AA}$ , and  $E_b = ?$  are the central wavelength, full width at half maximum, and strength of the bump, respectively. Once dust attenuation is applied to the SSPs, we sum them up to get the rest-frame luminosity of the galaxy. In total, our SPS model has 12 free parameters:  $M_*$ , 4 SFH basis coefficients,  $f_{\text{burst}}$ ,  $t_{\text{burst}}$ , 2 ZH basis coefficients,  $\tau_{\text{BC}}$ ,  $\tau_{\text{ISM}}$ , and  $n_{\text{dust}}$ .

In practice, evaluating each SSP using FSPS requires X seconds. For each model evaluation, we evaluate  $\sim 43$  SSPs in each of the log-spaced  $t_{\text{lookback}}$  bins; this takes X seconds. Though this is not a prohibitive computational cost on its own, sampling a high dimensional parameter space for inference requires  $> 100,000$  evaluations — *i.e.*  $> 100$  CPU hours *per galaxy*. For the  $> 10$  million BGS galaxies, this would require *a billion* CPU hours. Instead, we use an emulator for the model luminosity, which uses a Principal Component Analysis (PCA) neural network (NN) following the approach of Alsing et al. (2019).

TODO

TODO  
TODOTODO  
TODO



**Table 1.** Parameters of the PROVABGS SPS model used for joint SED modeling of DESI photometry and spectroscopy.

name	description	prior
$\log M_*$	log galaxy stellar mass	uniform over [8, 13]
$\beta_1, \beta_2, \beta_3, \beta_4$	NMF basis coefficients for SFH	Dirichlet prior
$f_{\text{burst}}$	fraction of total stellar mass formed in starburst event	uniform over [0, 1]
$t_{\text{burst}}$	time of starburst event	uniform over [10Myr, 13.2Gyr]
$\gamma_1, \gamma_2$	NMF basis coefficients for ZH	uniform over []
$\tau_{\text{BC}}$	Birth cloud optical depth	uniform over []
$\tau_{\text{ISM}}$	diffuse-dust optical depth	uniform over []
$n_{\text{dust}}$	? dust index	uniform over []

Our emulator consists of a NN and PCA basis functions. The NN provides a flexible and accurate mapping between the SPS model parameters and PCA coefficients — *i.e.* the NN predicts PCA coefficients for a given set of SPS parameters. Then the linear combination of the predicted coefficients and PCA basis functions give us the emulated model luminosity. The PCA basis functions and NN are trained using 1,000,000 pairs of SPS parameters and model luminosity,  $(\theta, L(\lambda; \theta))$ . Throughout the wavelength range relevant for BGS,  $3000 < \lambda < 9800\text{\AA}$ , we achieve  $< 1\%$  accurate with the emulator. For details on the training, validation, and performance of our PCA NN emulator, we refer readers to Kwon *et al.* (in prep.).

From the rest-frame luminosity, we obtain the observed-frame, redshifted, flux in the same way as Eq. 2. In our case, redshift is not a free parameter since we will have spectroscopic redshifts for every DESI BGS galaxy. To model DESI photometry, we convolve the model flux with the LS broadband filters as in Eq. 3. To model DESI spectra, we first apply Gaussian velocity dispersion. In this work, we keep velocity dispersion fixed at 0km/s but in practice velocity dispersion can be set as a free parameter. Then the broadened flux is resampled into the wavelength binning of the observed DESI spectra, which has spectral resolution  $R = 2000 - 5000$  over  $3600 < \lambda < 9800\text{\AA}$ . Finally, the SPS model photometry and spectrum can be directly compared to observations. TODO

### 3.2. Parameter Inference

Using the SPS model above, we perform Bayesian parameter inference to derive posterior probability distributions of the SPS parameters from photometry and spectroscopy. From Bayes rule, we write down the posterior as

$$p(\theta | \mathbf{X}) \propto p(\theta) p(\mathbf{X} | \theta) \quad (13)$$

where  $\mathbf{X}$  is the photometry or spectrum and  $\theta$  is the set of SPS parameters.  $p(\mathbf{X} | \theta)$  is the likelihood, which we calculate separately for the photometry

$$\mathcal{L}^{\text{photo}} \propto \exp \left[ -\frac{1}{2} \left( \frac{X^{\text{photo}} - m^{\text{photo}}(\theta)}{\sigma^{\text{photo}}} \right)^2 \right] \quad (14)$$

and for the spectrum

$$\mathcal{L}^{\text{spec}} \propto \exp \left[ -\frac{1}{2} \left( \frac{X^{\text{spec}} - m^{\text{spec}}(\theta)}{\sigma^{\text{spec}}} \right)^2 \right]. \quad (15)$$

$m^{\text{photo}}$  and  $m^{\text{spec}}$  represent SPS model for photometry and spectroscopy.  $\sigma^{\text{photo}}$  and  $\sigma^{\text{spec}}$  represent the uncertainties on the measured photometry and spectrum. We consider the photometry independent from the spectrum so we combine the likelihoods when jointly model the spectrophotometry:

$$\log \mathcal{L} \approx \log \mathcal{L}^{\text{photo}} + \log \mathcal{L}^{\text{spec}}. \quad (16)$$

$p(\theta)$  is the prior on the SPS parameters. For most of our parameters, we use uninformative uniform priors with conservatively chosen ranges that are listed in Table ?? . However, for the priors of  $\{\beta_1, \beta_2, \beta_3, \beta_4\}$ , the NMF coefficients for the SFH, we use a Dirichlet distribution. With a Dirichlet distribution,  $\beta_i$  are within  $0 < \beta_i < 1$  and satisfy the constraint  $\sum_i \beta_i = 1$ . This maintains the normalization of the SFH in Eq. 7.

Now that we can evaluate the posterior using the likelihood and prior, we derive the posterior distributions using Markov Chain Monte Carlo (MCMC) sampling. We use the [Karamanis & Beutler \(2020\)](#) ensemble slice sampling MCMC algorithm with the ZEUS Python package. **highlight advantages of ensemble slice MCMC** TODO

When we sample the posterior, we do not directly sample our 12 dimensional SPS parameter space. This is because we impose a Dirichlet prior on the SFH NMF coefficients. Dirichlet distributions are difficult to directly and efficiently sample so instead we use the [Betancourt \(2012\)](#) sampling method, which allows us to transform an  $N$  dimensional Dirichlet distribution into an easier to sample  $N - 1$  dimensional space. Hence, we sample the posterior in the transformed 11 dimensional space.

James's work in convergence here. talk about how in principle speculator can easily be used with HMC because you can get derivatives with backpropagation. provide detailed profiling of SED fitting and projections for full 10million galaxy BGS sample.

In Figure 3.2 we present the posterior as well as the best-fit photometry and spectroscopy using speculator + emcee.

## 4. RESULTS

Figure 6


$$\text{SFR}_{100\text{Myr}} = \frac{\int_{t_{\text{age}} - 100\text{Myr}}^{t_{\text{age}}} \text{SFH}(t) dt}{100 \text{ Myr}} \quad (17)$$

In order to quantify the precision and accuracy of the inferred physical properties for our simulated galaxy population, we begin by assuming that the discrepancy between the inferred and true parameters for each galaxy  $\Delta_{\theta,i}$


$$\theta_i^{\text{inf}} = \theta_i^{\text{true}} + \Delta_{\theta,i} \quad (18)$$

where  $\Delta_{\theta,i}$  is sampled from a Gaussian distribution

$$\Delta_{\theta,i} \sim \mathcal{N}(\mu_{\Delta_\theta}, \sigma_{\Delta_\theta}). \quad (19)$$



**Figure 5.** *Top:* Posterior probability distribution of our 12 SPS model parameters derived from joint SED modeling of the mock DESI photometry and spectra. The contours mark the 68 and 95% percentiles of the distribution. *Bottom:* Comparison of our SPS model evaluated at the best-fit parameters (orange) with the mock observations (black). On the left panel, we compare the  $g$ ,  $r$ ,  $z$  band magnitudes; on the right, we compare the spectroscopy.



**Figure 6.** Comparison between the true,  $\theta_{\text{true}}$ , and inferred galaxy properties,  $\hat{\theta}$ . The from the left to right columns, we compare  $\log M_*$ ,  $\log \text{SFR}_{1\text{Gyr}}$ ,  $\log Z_{\text{MW}}$ , and  $\tau_{\text{ISM}}$ . The inferred galaxy properties are derived from SED modeling of spectra (top row), photometry (middle row), and joint spectra + photometry (bottom row).

This Gaussian distribution is described by population hyperparameters  $\mu_{\Delta_\theta}$  and  $\sigma_{\Delta_\theta}$ , the mean and standard deviation, which quantify the accuracy and precision of the inferred physical properties for the population.

Given the photometry and spectrum of our galaxies,  $\{\mathbf{D}_i\}$ , we can get the posteriors for these population hyperparameters  $\eta_\Delta = \{\mu_{\Delta_\theta}, \sigma_{\Delta_\theta}\}$  using a hierarchical Bayesian framework (?):

$$p(\eta_\Delta | \{\mathbf{D}_i\}) = \frac{p(\eta_\Delta) p(\{\mathbf{D}_i\} | \theta_\Delta)}{p(\{\mathbf{D}_i\})} \quad (20)$$

$$= \frac{p(\eta_\Delta)}{p(\{\mathbf{D}_i\})} \int p(\{\mathbf{D}_i\} | \{\theta_i\}) p(\{\theta_i\} | \eta_\Delta) d\{\theta_i\}. \quad (21)$$

Naively the posteriors for each of the galaxies are not correlated, so we can factorize the expression above

$$p(\eta_\Delta | \{\mathbf{D}_i\}) = \frac{p(\eta_\Delta)}{p(\{\mathbf{D}_i\})} \prod_{i=1}^N \int p(\mathbf{D}_i | \theta_i) p(\theta_i | \eta_\Delta) d\theta_i \quad (22)$$

$$= \frac{p(\eta_\Delta)}{p(\{\mathbf{D}_i\})} \prod_{i=1}^N \int \frac{p(\theta_i | \mathbf{D}_i) p(\mathbf{D}_i)}{p(\theta_i)} p(\theta_i | \eta_\Delta) d\theta_i \quad (23)$$

$$= p(\eta_\Delta) \prod_{i=1}^N \int \frac{p(\theta_i | \mathbf{D}_i) p(\theta_i | \eta_\Delta)}{p(\theta_i)} d\theta_i. \quad (24)$$

$p(\theta_i | \mathbf{D}_i)$  is the posterior for galaxy  $i$ . Hence, the integral can be which means the integral can be estimated using the MCMC sample from the posterior

$$p(\eta_\Delta | \{\mathbf{D}_i\}) = p(\eta_\Delta) \prod_{i=1}^N \frac{1}{S_i} \sum_{j=1}^{S_i} \frac{p(\theta_{i,j} | \eta_\Delta)}{p(\theta_{i,j})}. \quad (25)$$

$S_i$  is the number of MCMC samples and  $\theta_{i,j}$  is the  $j^{\text{th}}$  sample of galaxy  $i$ . We present the maximum a posteriori (MAP) estimates of  $\eta_\Delta$  for  $\log M_*$  and  $\log \text{SFR}$  in Figure 7.

$\eta_\Delta$  as a function of SNR/mag/colour.

discussion:

- sfh basis improves SFH accuracy?
- advantage of mcmc over MAP
- non-Gaussian posteriors
- science applicatoins
- comparison to other methods
- different observing condition doesn't impact our results?
- we use super simple dust mmodel. **CH:** more sophistiicated dust models doesn't matter?
- **CH:** we try fitting with a different IMF


## 5. SUMMARY

DESI is great.

Also advertise science papers with specific focus on mock challenge science papers.

## ACKNOWLEDGEMENTS

It's a pleasure to thank Joel Leja, Peter Melchior ... This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under contract No. DE-AC02-05CH11231. This project used resources of the National Energy Research Scientific



**Figure 7.** The discrepancies between the inferred and input/“true”  $M_*$ s (left) and SFRs (right) for our LGAL galaxies. In blue, we infer  $M_*$ s and SFRs using only photometry; in orange, we infer  $M_*$ s and SFRs by jointly fitting both photometry and spectroscopy. *Jointly fitting spectroscopy and photometry improves constraints on galaxy properties.*

Computing Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

## APPENDIX


### A. SED MODELING UNCERTAINTIES FROM STELLAR EVOLUTION

### B. COMPARISON TO OTHER METHODS

## REFERENCES

- Alsing J., et al., 2019, [arXiv:1911.11778 \[astro-ph\]](#)  
 Betancourt M. J., 2012, [arXiv:1010.3436 \[physics\]](#)  
 10.1063/1.3703631, pp 157–164  
 Caplar N., Tacchella S., 2019, [arXiv:1901.07556 \[astro-ph\]](#)  
 Carnall A. C., McLure R. J., Dunlop J. S., Davé R., 2017, [arXiv:1712.04452 \[astro-ph\]](#)  
 Carnall A. C., Leja J., Johnson B. D., McLure R. J., Dunlop J. S., Conroy C., 2018, [arXiv:1811.03635 \[astro-ph\]](#)  
 Chabrier G., 2003, [Publications of the Astronomical Society of the Pacific](#), 115, 763  
 Charlot S., Fall S. M., 2000, [The Astrophysical Journal](#), 539, 718





**Figure 8.** do we want to test bias against other properties? (e.g. obs condition?)

**Table 2.** Table of  $\theta_{\text{inf}} - \theta_{\text{true}}$  and uncertainties for different CIGALE setups

set	CIGALE A	CIGALE B	CIGALE C	CIGALE D
$\Delta M_{\text{tot}}$	0.07	0.09	0.27	0.05
$M_{\text{err}}$	0.13	0.20	0.16	0.16
$\Delta \text{Age}$	1.78	1.59	1.97	1.92
$\text{Age}_{\text{err}}$	2.36	2.63	2.47	2.47
$\Delta Z$	0.0037	0.0026	0.0027	0.0027
$Z_{\text{err}}$	0.0091	0.0085	0.0089	0.0089

Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *The Astrophysical Journal*, 823, 102

Cichocki A., Phan A.-H., 2009, *IEICE Transactions on Fundamentals of Electronics Communications and Computer Sciences*, 92, 708

Conroy C., Gunn J. E., 2010, *The Astrophysical Journal*, 712, 833


Conroy C., Gunn J. E., White M., 2009, *The Astrophysical Journal*, 699, 486

Dey A., et al., 2019, *AJ*, 157, 168


Dotter A., 2016, *The Astrophysical Journal Supplement Series*, 222, 8

Févotte C., Idier J., 2011, arXiv:1010.1763 [cs]


Hahn C., Tinker J. L., Wetzel A., 2019, arXiv:1910.01644 [astro-ph]



**Figure 9.** *Top:* Spectral energy distributions generated using MIST (black), Padova (blue), Parsec (orange), and BaSTI (green) isochrone libraries for an arbitrary set of SPS parameters. For our fiducial SED model, we use the MIST isochrone. *Bottom:* Ratios of the SEDs generated using different isochrone libraries over our fiducial MIST SED. The ratios reveal  $> 10\%$  discrepancies at rest-frame UV wavelengths and significant offsets at higher wavelengths.



**Figure 10.**



**Figure 11.** The properties inferred from CIGALE photometry fit as a function of true properties. Configuration CIGALE A, B, C, and D on one plot.

- Henriques B. M. B., White S. D. M., Thomas P. A., Angulo R., Guo Q., Lemson G., Springel V., Overzier R., 2015, [Monthly Notices of the Royal Astronomical Society](#), 451, 2663
- Iyer K. G., et al., 2020, [Monthly Notices of the Royal Astronomical Society](#), 498, 430
- Karamanis M., Beutler F., 2020, arXiv e-prints, p. arXiv:2002.06212
- Kriek M., Conroy C., 2013, [The Astrophysical Journal Letters](#), 775, L16
- Lee D. D., Seung H. S., 1999, [Nature](#), 401, 788
- Leja J., Johnson B. D., Conroy C., van Dokkum P. G., Byler N., 2017, [The Astrophysical Journal](#), 837, 170
- Leja J., Carnall A. C., Johnson B. D., Conroy C., Speagle J. S., 2019, [ApJ](#), 876, 3
- Mathis J. S., 1983, [The Astrophysical Journal](#), 267, 119
- Nelson D., et al., 2018, [Monthly Notices of the Royal Astronomical Society](#), 475, 624
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, [The Astrophysical Journal Supplement Series](#), 192, 3
- Paxton B., et al., 2013, [The Astrophysical Journal Supplement Series](#), 208, 4
- Paxton B., et al., 2015, [The Astrophysical Journal Supplement Series](#), 220, 15
- Pillepich A., et al., 2018, [Monthly Notices of the Royal Astronomical Society](#), 473, 4077
- Ruiz-Macias O., et al., 2021, [Monthly Notices of the Royal Astronomical Society](#), 502, 4328
- Serra P., Amblard A., Temi P., Burgarella D., Giovannoli E., Buat V., Noll S., Im S., 2011, [The Astrophysical Journal](#), 740, 22
- Simha V., Weinberg D. H., Conroy C., Dave R., Fardal M., Katz N., Oppenheimer B. D., 2014, arXiv e-prints, p. arXiv:1404.0402
- Sparre M., Hayward C. C., Feldmann R., Faucher-Giguère C.-A., Muratov A. L., Kereš D., Hopkins P. F., 2017, [Monthly Notices of the Royal Astronomical Society](#), 466, 88
- Springel V., et al., 2018, [Monthly Notices of the Royal Astronomical Society](#), 475, 676
- Tacchella S., et al., 2021, arXiv e-prints, 2102, arXiv:2102.12494
- Zahid H. J., Geller M. J., Fabricant D. G., Hwang H. S., 2016, [The Astrophysical Journal](#), 832, 203