

PROVABGS Probabilistic Stellar Mass Function of the BGS One-Percent Survey

CHANGHOON HAHN^{1,*}

¹*Department of Astrophysical Sciences, Princeton University, Peyton Hall, Princeton NJ 08544, USA*

(Dated: DRAFT -- 6d00025 -- 2021-01-14 -- NOT READY FOR DISTRIBUTION)

ABSTRACT

We present the probabilistic stellar mass function (pSMF) of galaxies in the DESI Bright Galaxy Survey (BGS), observed as part of the One-Percent Survey. We derive the pSMF from posteriors of stellar mass, M_* , for all the BGS galaxies using statistically rigorous hierarchical inference. The M_* posteriors are inferred from DESI photometry and spectroscopy using the [Hahn et al. \(2022\)](#) PROVABGS SED modeling framework. We

Keywords: keyword1 – keyword2 – keyword3

1. INTRODUCTION

With large galaxy surveys, such as the Sloan Digital Sky Survey (SDSS; [York et al. 2000](#)), Galaxy and Mass Assembly survey (GAMA; [Driver et al. 2011](#)), and PRISM Multi-object Survey (PRIMUS; [Coil et al. 2011](#)), the galaxy population can now largely be characterized with a small number of scaling relations and population statistics (see [Blanton & Moustakas 2009](#), for a review). The stellar mass function (SMF), for instance, precisely characterizes the overall stellar mass, M_* , distribution of galaxies and its evolution across cosmic history ([Moustakas et al. 2013](#); [Muzzin et al. 2013](#); [Leja et al. 2019](#); [Driver et al. 2022](#)).

The relationship between the stellar masses and star formation rates of galaxies reveal a bimodality in the galaxy population with star-forming galaxies lying on a tightly correlated “star formation sequence” (SFS; [Noeske et al. 2007](#); [Daddi et al. 2007](#); [Salim et al. 2007](#); [Speagle et al. 2014](#); [Hahn et al. 2019](#)). From $z \sim 2$, there is an overall decline in star formation rates of galaxies in the SFS. This is accompanied by an increase in the fraction of quiescent galaxies caused by the quenching of star formation in some galaxies (????). Additional scaling relations among galaxy properties such as the mass-metallicity relation (?) or **add more scaling relations here** have also been firmly established. They connect **some physical insight** More precise and accurate measurements of the statistical distributions of the properties for galaxy populations at different cosmic epochs have the potential to shed further light on galaxy formation and evolution.

For one, they have the potential to reveal new trends among galaxies undetected by previous observations and open new discovery space. They can also be used to test galaxy formation models

* changhoon.hahn@princeton.edu

TODO
TODO

spanning empirical models (*e.g.* UNIVERSEMACHINE; ?), semi-analytic models (*e.g.* ?Henriques et al. 2015; Somerville et al. 2015), and hydrodynamical simulations (see Somerville & Davé 2015, for a review). Empirical models, for example, have been used to measure the timescale of star formation quenching (???) or the dust content of galaxies (?).

Furthermore, observations have already been used to infer parameters that dictate the physical processes in semi-analytic models (*e.g.* Henriques et al. 2009; Lu et al. 2014; Henriques et al. 2015). Although full parameter exploration is currently computationally prohibitively for hydrodynamical simulations, they have been extensively compared to observations: *e.g.* Genel et al. (2014); ?; ?; ?; ?. Soon machine learning techniques for accelerating and emulating simulations will enable us to go beyond such comparisons and broadly explore parameter space and galaxy formation models (*e.g.* ?). While many different approaches are available for expanding our understanding of galaxies, they all require statistically powerful galaxy samples with well controlled systematics and well understood selection functions.

One survey that will provide galaxy samples with unprecedented statistical power is the Dark Energy Spectroscopic Instrument (DESI; ???). Over its 5 year operation, DESI will observe galaxy spectra using the 4-meter Mayll telescope at Kitt Peak National Observatory with a focal plane filled with 5000 robotically-actuated fibers that direct the light to ten optical spectrographs. It will observe ~ 40 million galaxy spectra over $360 < \lambda < 980$ nm with spectral resolution of $2000 < \lambda/\Delta\lambda < 5500$ over $\sim 14,000$ deg², a third of the sky. In addition, DESI galaxies will also have photometry from the Legacy Imaging Surveys Data Release 9 (LS; ?). LS is a combination of three public projects (Dark Energy Camera Legacy Survey, Beijing-Arizona Sky Survey, and Mayall *z*-band Legacy Survey) that jointly imaged the DESI footprint in three optical bands (*g*, *r*, and *z*). DESI began observing its main survey in May 14, 2021.

As part of the survey, DESI is conducting the Bright Galaxy Survey (BGS; Hahn et al. 2022). BGS spans the same 14,000deg² footprint and will include low redshift $z < 0.6$ galaxies that can be observed during bright time, when the night sky is $\sim 2.5\times$ brighter than nominal dark conditions, BGS will provide two galaxy samples: the BGS Bright sample, a $r < 19.5$ magnitude-limited sample of ~ 10 million galaxies, and the BGS Faint sample, a fainter $19.5 < r < 20.175$ sample of ~ 5 million galaxies selected using a surface brightness and color. The selection and completeness of the BGS samples are characterized in detail in ?. Compared to the seminal SDSS main galaxy survey, BGS will provide a galaxy sample two magnitudes deeper, over twice the sky, and double the median redshift $z \sim 0.2$. It will observe a broader range of galaxies than previous surveys and provide an opportunity to measure galaxy population statistics with unprecedented precision.

BGS will also be accompanied by a value-added catalog: the Probabilistic Value-Added BGS (PROVABGS; Hahn et al. 2022; Kwon et al. 2022). For every BGS galaxy, PROVABGS will provide physical properties such as stellar mass (M_*), average star formation rate ($\overline{\text{SFR}}$), stellar metallicity (Z_*), stellar age (t_{age}), and dust content. These galaxy properties will be from state-of-the-art Spectral Energy Distribution (SED) modeling of both DESI photometry and spectroscopy in a full Bayesian inference framework. The SED model is designed to minimize model misspecification by using a highly flexible non-parameteric star formation and metallicity histories as well as a flexible

dust attenuation model. Furthermore, the properties will be inferred using a fully Bayesian inference framework and provide statistically rigorous estimates of uncertainties and degeneracies among the parameters. Ultimately, PROVABGS will provide consistently measured galaxy properties that will enable analyses to fully take advantage of the statistical power of BGS with new techniques and approaches.

A key application for PROVABGS will be measuring population statistics using statistically rigorous methodology that correctly propagates the uncertainties in galaxy property measurements. Current population statistics are by and large derived from simply binning best-fit point estimates of galaxy properties. [Malz & Hogg \(2020\)](#) demonstrated, in the context of inferring redshift distributions from individual photometric redshift measurements, that using point estimates is statistically incorrect and can lead to biased redshift distributions. Similarly, the standard approach can also lead to bias population statistics.

Instead, we can estimate population statistics from combining individual PROVABGS posteriors of galaxy properties using population inference in a hierarchical Bayesian framework (*e.g.* [Hogg et al. 2010](#); [Foreman-Mackey et al. 2014](#); [Baronchelli et al. 2020](#)). This approach summary of why its better TODO In this work, we present the first of such population statistic measurement for BGS: the probabilistic stellar mass function (pSMF).

In particular, we present the pSMF of BGS galaxies observed during the DESI One-Percent Survey, a survey validation program conducted before the main survey operations. We also present the statistical methodology for the population inference as well as the our methods for accounting for observational incompleteness. We begin in Section 2 with an overview of the BGS galaxies observed during the DESI One-Percent Survey. Then, in Section 3, we briefly summarize the PROVABGS SED modeling framework used to infer the physical properties of the BGS galaxies. Afterwards, we present the pSMF inferred from the BGS observations in Section 4. We summarize and discuss our results in Section ?. Throughout the We use XXXXXX cosmology.

2. THE DESI BRIGHT GALAXY SURVEY: ONE-PERCENT SURVEY

DESI began its five years of operations in May 14, 2021. Before its start, DESI conducted the Survey Validation (SV) campaign to verify that the survey will meets its scientific and performance requirements. The SV campaign was divided into two main programs: the first, SV1, characterized the survey’s performance for different observing conditions and was used to optimize sample selection. The second, the One-Percent Survey (or SV3), observed a dataset that can be used for representative clustering measurements and deliver a ‘truth’ sample with high completeness over an area at least 1% of the expected main survey footprint. We refer readers to ? for details on the DESI SV programs. In this work, we focus on BGS galaxies observed during the One-Percent Survey.

The One-Percent Survey observed on 38 nights from April 2021 to the end of May 2021. During this time DESI observed 288 bright time exposures that cover 214 BGS ‘tiles’, planned DESI pointings. The tiles were arranged so that a set of 11 overlapping tiles has their centers arranged around a 0.12 deg circle, forming a ‘rosette’ completeness pattern. In total, the One-Percent Survey observed 20 rosettes covering 180 deg² spanning the northern galactic cap (see Figure 1 in [Hahn et al. 2022](#)).

All BGS spectra observed during the One-Percent Survey are reduced using the ‘Fuji’ version of the DESI spectroscopic data reduction pipeline (?). First, spectra are extracted from the spectrograph CCDs using the *Spectro-Perfectionsim* algorithm of ?. Then, fiber-to-fiber variations are corrected by flat-fielding and a sky model, empirically derived from sky fibers, is subtracted from each spectrum. Afterwards, the fluxes in the spectra are calibrated using stellar model fits to standard stars. The final processed spectra is then derived by co-adding the calibrated spectra across exposures of the same tile. In total, DESI observed spectra of 155,022 BGS Bright and 109,418 BGS Faint targets during the One-Percent Survey.

For each spectrum, redshift is measured using REDROCK¹ (?), a redshift fitting algorithm that uses χ^2 minimization computed from a linear combination of Principal Component Analysis (PCA) basis spectral templates in three template classes (“stellar”, “galaxy”, and “quasar”). REDROCK also provides measures of redshift uncertainty, ZERR and redshift confidence, $\Delta\chi^2$, which corresponds to the difference between the χ^2 values of the best-fit model and the next best-fit model. We restrict our sample to galaxy targets with reliable redshift measurements. We only keep targets with spectra classified as galaxy spectra by REDROCK, no REDROCK warning flags, $\Delta\chi^2 > 40$, and REDROCK redshift uncertainty ZERR $< 0.0005(1+z)$. We also exclude any targets observed using malfunctioning fiber positioners. Lastly, we impose a redshift range of $0 < z < 0.6$. After these cuts, our One-Percent Survey BGS sample includes 143,074 BGS Bright galaxies and 96,771 BGS Faint galaxies.

3. PROVABGS SED MODELING

For each BGS galaxy, we derive its M_* and other physical properties, $\overline{\text{SFR}}$, Z_{MW} , and $t_{\text{age,MW}}$ from DESI photometry and spectroscopy using the PROVABGS SED modeling framework (Hahn et al. 2022). PROVABGS models galaxy SEDs using stellar population synthesis with non-parametric star-formation history (SFH) with a starburst, a non-parametric metallicity history (ZH) that evolves with time, and a flexible dust attenuation prescription. The non-parametric SFH and ZH prescriptions are derived from SFHs and ZHs of simulated galaxies in the Illustris hydrodynamic simulation (Vogelsberger et al. 2014; Genel et al. 2014; Nelson et al. 2015) and provide compact and flexibly representations of SFHs and ZHs. For the stellar population synthesis, PROVABGS uses the Flexible Stellar Population Synthesis (FSPS; Conroy et al. 2009, 2010) model with MIST isochrones (Paxton et al. 2011, 2013, 2015; Choi et al. 2016; Dotter 2016), Chabrier (2003) initial mass function (IMF), and a combination of MILES (Sánchez-Blázquez et al. 2006) and BaSeL (Lejeune et al. 1997, 1998; Westera et al. 2002) spectral libraries.

Furthermore, PROVABGS provides a Bayesian inference framework that infers full posterior probability distributions of the SED model parameter: $p(\theta | \mathbf{X}^{\text{photo}}, \mathbf{X}^{\text{spec}})$, where $\mathbf{X}^{\text{photo}}$ represents the photometry and \mathbf{X}^{spec} represents the spectroscopy. In total, θ has 13 parameters: M_* , 6 parameters specifying the SFH ($\beta_1, \beta_2, \beta_3, \beta_4, f_{\text{burst}}, t_{\text{burst}}$), 2 parameters specifying ZH (γ_1, γ_2), 3 parameters specifying dust attenuation ($\tau_{\text{BC}}, \tau_{\text{ISM}}, n_{\text{dust}}$), and a nuisance parameter for the fiber aperture effect. Posteriors accurately estimate uncertainties and degeneracies among galaxy properties. Furthermore, as we later demonstrate, they are essential for statistically rigorous hierarchical population inference.

¹ <https://redrock.readthedocs.io>

In practice, accurately estimating a 13 dimensional posterior requires a large number ($\gtrsim 100,000$) SED model evaluations, which requires prohibitive computational resources — ~ 10 CPU hours per galaxy. To address this challenge, PROVABGS samples the posterior using the Karamanis & Beutler (2020) ensemble slice Markov Chain Monte Carlo (MCMC) sampling with the ZEUS Python package². PROVABGS further accelerates the inference by using neural emulators for the SED models. The emulators are accurate to subpercent level and $>100\times$ faster than the original SED model based on FSPS (Kwon et al. 2022). With ZEUS and neural emulation, deriving a posterior takes ~ 5 min per galaxy with PROVABGS. Moreover, Hahn et al. (2022) demonstrated PROVABGS can accurately infer M_* overall the full expected M_* range of BGS, using forward modeled synthetic DESI BGS observations.

In Figure 1, we demonstrate the PROVABGS SED modeling framework for a randomly selected BGS Bright galaxy with $z = 0.2242$ (target ID: 39627757520424630). In the top panels, we present the posteriors of galaxy properties, M_* , $\overline{\text{SFR}}$, Z_{MW} , and $t_{\text{age,MW}}$, inferred from DESI photometry and spectroscopy. We mark the 12, 40, 68, and 86 percentiles of posterior with the contours. The posteriors illustrate that we can precisely measure the properties of BGS galaxies from DESI photometry and spectroscopy. Furthermore, with the full posterior, we accurately estimate the uncertainties on the galaxy properties and the degeneracies among them (*e.g.* M_* and $\overline{\text{SFR}}$). In the bottom panels, we compare the PROVABGS SED model prediction using the best-fit parameter values (black) to DESI observations (blue). The left panel compares the optical g , r , and z band photometry while the right panel compares the spectra. The comparison shows good agreement between the best-fit model and the observations.

We derive a PROVABGS posterior (*e.g.* Figure 1) for every galaxy in the DESI One-Percent Survey. In Figure 2, we present the best-fit M_* measurements as a function of z for the BGS galaxies in DESI One-Percent Survey. We mark the galaxies in the BGS Bright sample in blue and the ones in the BGS Faint sample in orange.

4. RESULTS

From the posteriors of galaxy properties inferred using PROVABGS (Section 3), we derive the marginalized posteriors: $p(M_* | \mathbf{X}_i)$, the marginalized 1D posterior of M_* from observed spectrophotometry \mathbf{X}_i of galaxy i . Using these posteriors, we can estimate the probabilistic SMF (pSMF) of BGS galaxies using population inference in a hierarchical Bayesian framework (*e.g.* Hogg et al. 2010; Foreman-Mackey et al. 2014; Baronchelli et al. 2020). In other words, we can infer $p(\phi | \{\mathbf{X}_i\})$, the probability distribution of ϕ given the full DESI BGS observations, $\{\mathbf{X}_i\}$. ϕ is the set of population hyperparameters that describe the pSMF, $\Phi(M_*; \phi)$. We again emphasize that this approach is statistically rigorous and correctly propagates the uncertainties in our M_* measurements to the pSMF.

² <https://zeus-mcmc.readthedocs.io/>




Figure 1. *Top panels:* Posteriors of galaxy properties, M_* , $\overline{\text{SFR}}$, Z_{MW} , and $t_{\text{age,MW}}$, for a randomly selected BGS Bright galaxy with $z = 0.2242$ (target ID: 39627757520424630) inferred using the PROVABGS SED modeling framework from DESI photometry and spectroscopy. The contours mark the 12, 40, 68, and 86 percentiles of the posterior. With the PROVABGS posteriors, we accurately estimate the galaxy properties, their uncertainties, and any degeneracies among them. *Bottom panels:* Comparison of the best-fit PROVABGS SED model prediction (black) to observations (blue). We compare the g , r , and z band photometry in the left panel and spectra in the right panel. We infer the posterior of galaxy properties for every BGS galaxies in the DESI One-Percent Survey.

In this work, we estimate the pSMF using a Gaussian Mixture Model (GMM; Press et al. 1992; McLachlan & Peel 2000), which provides a highly flexible description of the M_* distribution:

$$\Phi(M_*; \phi) = \sum_{j=1}^k \mathcal{N}(M_*; \phi_j). \quad (1)$$




Figure 2. M_* as a function of z of BGS Bright (blue) and Faint (orange) galaxies in the DESI One-Percent Survey. For M_* , we use the best-fit values derived using PROVABGS. BGS Bright is a magnitude-limited sample to $r < 19.5$ while BGS Faint includes fainter galaxies $19.5 < r < 20.175$ selected using r_{fb} and color (?). In total, we infer the posteriors of 143,017 BGS Bright and 95,499 BGS Faint galaxies in the DESI One-Percent Survey spanning $0 < z < 0.6$.

k represents the number of Gaussian components. ϕ_j represent the mean and standard deviation of the j^{th} Gaussian component of the GMM. Previous works have used parametric functions (*e.g.* Schechter function) to describe the pSMF (Leja et al. 2019). We opt for GMMs in order to produce a non-parametric measurement of the pSMF. In subsequent work, Speranza et al. (in prep.), we will present the BGS pSMF measured using a parametric model with continuous redshift evolution.

To infer $p(\phi | \{\mathbf{X}_i\})$, we follow the same approach described in Hahn et al. (2022):

$$p(\phi | \{\mathbf{X}_i\}) = \frac{p(\phi) p(\{\mathbf{X}_i\} | \phi)}{p(\{\mathbf{X}_i\})} \quad (2)$$

$$= \frac{p(\phi)}{p(\{\mathbf{X}_i\})} \int p(\{\mathbf{X}_i\} | \{\theta_i\}) p(\{\theta_i\} | \phi) d\{\theta_i\}. \quad (3)$$

$$= \frac{p(\phi)}{p(\{\mathbf{X}_i\})} \prod_{i=1}^N \int p(\mathbf{X}_i | \theta_i) p(\theta_i | \phi) d\theta_i \quad (4)$$

$$= \frac{p(\phi)}{p(\{\mathbf{X}_i\})} \prod_{i=1}^N \int \frac{p(\theta_i | \mathbf{X}_i) p(\mathbf{X}_i)}{p(\theta_i)} p(\theta_i | \phi) d\theta_i \quad (5)$$

$$= p(\phi) \prod_{i=1}^N \int \frac{p(\theta_i | \mathbf{X}_i) p(\theta_i | \phi)}{p(\theta_i)} d\theta_i. \quad (6)$$

We can estimate the integral using S_i Monte Carlo samples from the individual posteriors $p(\theta_i | \mathbf{X}_i)$:

$$\approx p(\phi) \prod_{i=1}^N \frac{1}{S_i} \sum_{j=1}^{S_i} \frac{p(\theta_{i,j} | \phi)}{p(\theta_{i,j})}. \quad (7)$$

Since the sample of BGS galaxies is not volume-limited and complete as a function of M_* , we must account for the selection effect and incompleteness when estimating the pSMF. To account for the selection effects of the BGS samples, we include weights derived from z_i^{\max} , the maximum redshift that galaxy i could have and still be included in the BGS samples. We derive z_i^{\max} for every galaxy using by redshifting the SED predicted by the best-fit parameters. We then derive V_i^{\max} , the comoving volume out to z_i^{\max} , and include a factor of $1/V_i^{\max}$ in the galaxy weight w_i .

Next, we include correction weights for spectroscopic incompleteness driven by fiber assignment and redshift failures. Incompleteness from fiber assignment is due to the fact that DESI is not able to assign fibers to all galaxies included in the BGS target selection. Furthermore, due to the clustering of galaxies there is significant variation in the assignment probability. Meanwhile, incompleteness from redshift failure is caused by the fact that we do not successfully measure the redshift for every spectra and the redshift failure rate depends significantly on the surface brightnesses of the galaxies and the signal-to-noise ratio of the spectra. We describe how we derive the incompleteness correction weights for fiber assignment and redshift failures, $w_{i,\text{FA}}$ and $w_{i,\text{ZF}}$, in Appendix A. Each BGS galaxy is assigned a weight of $w_i = (w_{i,\text{FA}} \times w_{i,\text{ZF}})/V_i^{\max}$.

We modify Eq. 2 to include galaxy weights, w_i :

$$p(\phi | \{\mathbf{X}_i\}) \approx \frac{p(\phi)}{\prod_{i=1}^N p(\mathbf{X}_i)^{w_i}} \prod_{i=1}^N \left(\int p(\mathbf{X}_i | \theta_i) p(\theta_i | \phi) d\theta_i \right)^{w_i} \quad (8)$$

$$\approx \frac{p(\phi)}{\prod_{i=1}^N p(\mathbf{X}_i)^{w_i}} \prod_{i=1}^N \left(\sum_{j=1}^{S_i} \frac{p(\theta_{i,j} | \phi)}{p(\theta_{i,j})} \right)^{w_i} \quad (9)$$

$$\approx \frac{p(\phi)}{\prod_{i=1}^N p(\mathbf{X}_i)^{w_i}} \prod_{i=1}^N \left(\sum_{j=1}^{S_i} \frac{q_\phi(\theta_{i,j})}{p(\theta_{i,j})} \right)^{w_i}. \quad (10)$$

In practice, we do not derive the full posterior $p(\phi | \{\mathbf{X}_i\})$. Instead we derive the maximum a posteriori (MAP) hyperparameter ϕ_{MAP} that maximizes $p(\phi | \{\mathbf{X}_i\})$ or $\log p(\phi | \{\mathbf{X}_i\})$. We expand,

$$\log p(\phi | \{\mathbf{X}_i\}) \approx \log p(\phi) + \sum_{i=1}^N w_i \log \left(\sum_{j=1}^{S_i} \frac{q_\phi(\theta_{i,j})}{p(\theta_{i,j})} \right). \quad (11)$$

Since the first two terms are constant, we derive ϕ_{MAP} by maximizing

$$\max_{\phi} \sum_{i=1}^N w_i \log \left(\sum_{j=1}^{S_i} \frac{q_\phi(\theta_{i,j})}{p(\theta_{i,j})} \right). \quad (12)$$

using the ADAM optimizer (Kingma & Ba 2017). We derive ϕ_{MAP} for BGS galaxies in redshift bins of width $\Delta z = 0.04$ starting from $z = 0.01$ in order to examine the redshift evolution of the SMF within BGS.




Figure 3. The probabilistic SMF (pSMF) of BGS galaxies in the One-Percent Survey at $0.01 < z < 0.05$ (black line). We represent uncertainties on the pSMF, estimated using a standard jackknife technique (Appendix B), in the shaded regions. The solid line represents the pSMF above the completeness limit $M_* > M_{\text{lim}} = 10^{8.975} M_\odot$ (Appendix C). In the left panel, we present the pSMFs of BGS Bright (blue) and Faint (orange) galaxies. In the right panel, we include SMF measurements from previous spectroscopic surveys for comparison: SDSS (Moustakas et al. 2013; Bernardi et al. 2017) and GAMA (Driver et al. 2022). Overall, the pSMF of BGS are in good agreement with SMF measurements from previous surveys.




Figure 4. The BGS pSMF over the redshift range $0.01 < z < 0.17$ in bins of $\Delta z = 0.04$. The shaded regions represent the uncertainties on the pSMF, estimated using a standard jackknife technique. The solid lines represent the pSMF above the completeness limit $M_* > M_{\text{lim}}$ while the dashed lines represent the pSMF below the limit. There is no significant redshift evolution of the pSMF given the statistical uncertainties. The main BGS survey will observe $> 100\times$ more galaxies than the One-Percent Survey.

4.1. The Probabilistic Stellar Mass Function

We present the probabilistic SMF (pSMF) of $0.01 < z < 0.05$ BGS galaxies in the One-Percent Survey in Figure 3 (black line). The shaded regions represent the uncertainties of the pSMF from sample variance, which we derive using a standard jackknife technique (Appendix B) and are conservative estimates (Norberg et al. 2009). In the left panel, we also present the pSMFs of the BGS Bright (blue) and Faint (orange) galaxies. BGS Bright galaxies are selected using a $r > 19.5$ magnitude limit. As a result, the BGS Bright sample is M_* complete above $M_{\text{lim}} > 10^{8.975} M_\odot$. We derive M_{lim} in Appendix C and mark the pSMF above the completeness limit in solid and below the limit in dashed. Meanwhile, the BGS Faint sample is selected using a surface brightness and color selection. It includes fainter galaxies, $19.5 < r < 20.175$, with overall lower M_* than the BGS Bright sample.

We also include the SMF estimated using the standard approach (black dotted). This SMF is derived using the best-fit M_* point estimates with the same galaxy weights, w_i . At intermediate M_* range, $10^9 < M_* < 10^{11} M_\odot$, we find good agreement with the pSMF. However, the standard approach significantly underestimates the SMF outside this M_* range. These discrepancies are due to the fact that point estimates of M_* ignore the uncertainties which contribute significantly at the most and least massive ends of the SMF. The discrepancies are present in all other redshift bins and underscore the importance of correctly propagating the M_* uncertainties.

In the right panel, we compare the BGS pSMF to SMF measurements from previous spectroscopic surveys: SDSS (Moustakas et al. 2013; Bernardi et al. 2017) (black circle and square) and GAMA (Driver et al. 2022) (black triangle). We note that there is significant variance in SMF measurements in the literature, especially at the high M_* end. This is partly due to the different modeling methodologies used to derive M_* , which can contribute > 0.1 dex discrepancies (?). Furthermore, there are also discrepancies due to photometric corrections applied to SDSS photometry, assumptions on the stellar populations, and dust (Bernardi et al. 2017). In subsequent work, we will present a detailed comparison of BGS M_* measurements using different methods. Overall, we find good agreement with previous SMF measurements, especially in the intermediate M_* range where we precisely infer the pSMF.

In Figure 4, we present the redshift evolution of the pSMF over $0.01 < z < 0.17$ in redshift bins of $\Delta z = 0.04$. The shaded region represent the jackknife uncertainties for the pSMF. The solid line represents the pSMF above M_{lim} while the dashed lines represent the pSMF below the limit. We only include 4 redshift bins, since $M_{\text{lim}} > 10^{10.5} M_\odot$ for $z > 0.17$ (Table 2). The pSMFs in Figure 4 do not reveal a significant redshift dependence given their uncertainties. We note that the large uncertainties for the $0.01 < z < 0.05$ pSMF is driven by large-scale structure at $\text{RA} \sim 195$ deg, $\text{Dec} \sim 28$ deg, and $z \sim 0.244$. The main BGS survey will observe $> 100\times$ more BGS galaxies than the One-Percent Survey and enable pSMF measurements with unprecedented precision.

4.2. Star-Forming and Quiescent Galaxies in the BGS

In addition to the pSMF of the full galaxy population, we can also examine the pSMF of the star-forming and quiescent subpopulations using $\overline{\text{SFR}}$, average SFR over the last 1 Gyr, inferred using PROVABGS. In Figure 5, we present the distribution of M_* versus average specific SFR,




Figure 5. The M_* – sSFR distribution of BGS galaxies at $z < 0.2$. sSFR is derived using $\overline{\text{SFR}}$, average SFR over the last 1 Gyr, inferred using PROVABGS. The M_* – sSFR distribution is bimodal with star-forming galaxies lying on the star-forming sequence. We classify galaxies with $\text{sSFR} > 10^{-11.2} \text{ yr}^{-1}$ as star-forming galaxies and $\text{sSFR} < 10^{-11.2} \text{ yr}^{-1}$ as quiescent.




Figure 6. The pSMF of star-forming (left) and quiescent (right) BGS Bright galaxies over $0.01 < z < 0.17$ in bins of $\Delta z = 0.04$. Star-forming and quiescent galaxies are classified using an empirically determined $\text{sSFR} = 10^{-11.2} \text{ yr}^{-1}$ cut. We represent the uncertainties for the pSMF in the shaded regions and the pSMFs above/below the M_* completeness limits in solid/dashed lines. The pSMFs suggest a decline in massive, $M_* > 10^{11} M_\odot$ star-forming galaxies and an increase in the quiescent galaxy population at $M_* < 10^{11} M_\odot$ at lower redshifts.

$\overline{\text{sSFR}} = \overline{\text{SFR}}/M_*$, for BGS Bright (blue) and Faint (orange) galaxies at $z < 0.2$. The M_* – sSFR distribution of the BGS galaxies reveal a clear bimodality with star-forming galaxies lying on the SFS and quiescent galaxies lying $\gtrsim 1$ dex below the sequence. Figure 5 also confirms that BGS Faint




Figure 7. The quiescent fraction of BGS Bright galaxies over $0.01 < z < 0.17$ in bins of $\Delta z = 0.04$. We present the uncertainties in the shaded region and only include the quiescent fraction above the M_* completeness limit. The quiescent fractions increase with M_* at all redshifts. Furthermore, the quiescent fractions suggest an overall increase in the quiescent population with lower redshift.

galaxies have overall lower M_* than BGS Bright galaxies and are primarily star-forming galaxies. This is due to the fact that the $(z - W1) - 1.2(g - r) + 1.2$ color used to select BGS Faint galaxies is a proxy for $H\alpha$ and $H\beta$ emission lines.

To further examine the star-forming and quiescent galaxy populations, we classify BGS Bright galaxies as star-forming or quiescent using a $\overline{\text{sSFR}} = 10^{-11.2} \text{ yr}^{-1}$ cut. We determine this cut empirically based roughly on the sSFR of the “green valley” between the SFS and the quiescent mode. We opt for a $\overline{\text{sSFR}}$ cut rather than more sophisticated methods in the literature (*e.g.* Hahn et al. 2019; Donnari et al. 2019) for simplicity. In Figure 6, we present the pSMF of star-forming and quiescent BGS Bright galaxies at $0.01 < z < 0.17$ in bins of $\Delta z = 0.04$. The shaded regions represent the jackknife uncertainties for the pSMF. The solid lines represent the pSMFs above the completeness limit while the dashed lines represent the pSMFs below the limit. The pSMF of quiescent galaxies suggest an increase in the number of galaxies below $M_* < 10^{11} M_\odot$. Meanwhile, the pSMF of star-forming galaxies shows a possible decline at the massive end over $0.01 < z < 0.17$.

Next, we present the fraction of quiescent galaxies in BGS Bright as a function of M_* over $0.01 < z < 0.17$ in Figure 7. The quiescent fraction is derived by taking the ratio of the pSMFs of quiescent galaxies over all galaxies and measured for each $\Delta z = 0.04$ bin. The shaded region represent the uncertainties derived from propagating the jackknife uncertainties of the pSMFs. We focus on the quiescent fraction of BGS Bright galaxies above the M_* completeness limit: $M_* > M_{\text{lim}}$. At each redshift bin, the quiescent fraction increases with M_* to ~ 1 at $M_* \sim 10^{11.5} M_\odot$. The quiescent fraction also suggests an increase in the quiescent population with redshift; although a clear trend is obfuscated by significant statistical uncertainties. Upcoming observations from the DESI main

survey will increase the number of BGS galaxies by $>100\times$ and dramatically improve the precision of quiescent fraction measurements.

5. SUMMARY AND DISCUSSION

Over its five year operation, starting on May 2021, the DESI Bright Galaxy Survey (BGS) will observe the spectra of ~ 15 million galaxies out to $z < 0.6$ over $14,000 \text{ deg}^2$. BGS will produce two galaxy samples: a $r < 19.5$ magnitude-limited BGS Bright sample and a fainter $19.5 < r < 20.175$ surface brightness and color selected BGS Faint sample. Compared to the SDSS main galaxy survey, BGS galaxy samples will be two magnitudes deeper, over twice the sky, and double the median redshift $z \sim 0.2$. Furthermore, each galaxy in BGS will have measurements of its physical properties (*e.g.* M_* , SFR, Z_* , t_{age}) derived using state-of-the-art SED modeling of DESI spectrophotometry from PROVABGS.

The BGS samples will include diverse galaxy subpopulations that can be used to reveal new trends among galaxies that were previously undetectable and open new discovery space. They will also provide statistical power to measure scaling relations and population statistics to characterize the global galaxy population and test galaxy formation models with unprecedented precision.

In this work, we showcase the potential of BGS by presenting the probabilistic stellar mass function (pSMF) using only $\sim 250,000$ BGS galaxies observed during one of DESI's survey validation program. We present the hierarchical population inference framework that enable us to accurately propagate uncertainties on galaxy properties (M_*) to population statistics (pSMF). We also describe how we account for selection effect and incompleteness (Appendix A).

All of the pSMFs presented in this work are measured from BGS galaxies observed during the DESI One-Percent Survey. DESI has already completed nearly two years of observations. current status of BGS

subsequent works federico's paper as subsequent work with schetcher function fits detailed stellar mass comparison project.

Furthermore, in subsequent work we will extend the hierarchical population inference framework to the SFR- M_* distribution and present the probabilistic SFR- M_* distribution and quiescent fraction.

ACKNOWLEDGEMENTS

It's a pleasure to thank

APPENDIX

A. SPECTROSCOPIC COMPLETENESS

Spectroscopic galaxy surveys, such as BGS, do not successfully measure the redshift for all of the galaxies they target. As a result, this spectroscopic incompleteness must be accounted for when measuring galaxy population statistics such as the SMF. In this appendix, we present how we estimate the spectroscopic incompleteness for BGS and derive the weights we use to correct for its impact on the SMF.

TODO
TODO

For BGS, spectroscopic incompleteness is primarily driven by fiber assignment and redshift failures. DESI uses 10 fiber-fed spectrographs with 5000 fibers but targets more galaxies than available fibers. For instance, the BGS Bright and Faint samples have ~ 860 and 530 targets/deg², respectively. For the 8 deg² field-of-view of DESI, this roughly correspond to 11,000 targets, significantly more than the 5000 available fibers. DESI only measures the spectra of targets that are assigned fibers. In fact, of the 5000, a minimum of 400 ‘sky’ fibers are dedicated to measuring the sky background for accurate sky subtraction and an additional 100 fibers are assigned to standard stars for flux calibration ?.

Furthermore, each fiber is controlled by a robotic fiber positioner on the focal plane. These positioners can rotate on two arms and be positioned within a circular patrol region of radius 1.48 arcmin (????). Although the patrol regions of adjacent positioners slightly overlap, the geometry of the positioners cause higher incompleteness in regions with high target density (?). To mitigate the incompleteness from the fiber assignment, BGS will observe its footprint with four passes. With this strategy, BGS achieves $\sim 80\%$ fiber assignment completeness (?).

To estimate fiber assignment completeness, we run the fiber assignment algorithm (?) on BGS targets 128 separate times. For each BGS galaxy, i , we count the total number of times out of 128 that the galaxy is assigned a fiber: $N_{i,\text{FA}}$. Then to correct for the fiber assignment incompleteness, we assign correction weights

$$w_{i,\text{FA}} = \frac{128}{N_{i,\text{FA}}} \quad (\text{A1})$$

to each BGS galaxy. explain what this means




Figure 8. Redshift success rate of BGS Bright galaxies as a function of r_{fiber} and TSNR2. TSNR2 is a statistic that quantifies the signal-to-noise ratio of the observed spectrum. The color map represents the mean redshift success rate in each hexbin. We mark the TSNR2 bins (black dashed) that we use to separately fit the redshift success rate as a function of r_{fiber} using Eq. A3. In each TSNR2 bin, redshift success decreases as r_{fiber} increases.




Figure 9. Redshift success rates of BGS Bright galaxies as a function of r_{fiber} in 6 TSNR2 bins. The error bars represent the poisson uncertainties. In each panel, we include the best-fit analytic (Eq. A3) approximation of the redshift success rate (dashed) derived from χ^2 minimization. We use this analytic approximation to calculate the galaxy weights to correct for spectroscopic incompleteness caused by failures to accurately measure redshifts from observed spectra.

Although we measure a spectrum for each galaxy assigned a fiber, we do not accurately measure redshifts for every spectra. This redshift measurement failure significantly contributes to spectroscopic incompleteness. For BGS, redshift failure of an observed galaxy spectrum depends mainly on fiber magnitude and a statistic, TSNR2. Fiber magnitude is the predicted flux of the BGS object within a 1.5'' diameter fiber; we use r -band fiber magnitude, r_{fiber} . TSNR2 roughly corresponds to the signal-to-noise ratio of the spectrum and is the statistic used to calibrate the effective exposure times in DESI observations (CITE).

In Figure 8, we present the redshift, z , success rate of BGS Bright galaxies as a function of r_{fiber} and TSNR2. In each hexbin, the color map represents the mean z -success rate. We include all hexbins with more than 2 galaxies. Overall, the z -success rate depends significantly on r_{fiber} : galaxies with fainter r_{fiber} have lower z -success rates. However, the r_{fiber} dependence itself varies in bins of TSNR2. We mark the edges of the bins in black dashed: $\log \text{TSNR2} = 2.0, 2.5, 3.0, 3.5, 3.85$. Within each of the TSNR2 bins, the r_{fiber} dependence of the z -success rate does not vary significantly. In Figure 9, we present the z -success rate of BGS Bright galaxies as a function of r_{fiber} for each of the 6 TSNR2 bins. We mark the range of TSNR2 in the bottom left of each panel. The errorbars represent the Poisson uncertainties of the z -success rate.

TODO




Figure 10. The RA and Dec of the 12 jackknife fields of the BGS One-Percent Survey used to estimate the uncertainties on the SMF from sample variance. We mark each field with a distinct color.

To correct for the effect of redshift failures, we include an additional correction weight for each BGS galaxy:

$$w_{i,\text{ZF}} = \frac{1}{f_{z\text{-success}}(r_{\text{fiber},i}, \text{TSNR2}_i)}. \quad (\text{A2})$$

$f_{z\text{-success}}(r_{\text{fiber},i}, \text{TSNR2}_i)$ is the z -success rate as a function of r_{fiber} and TSNR2 of the galaxy. Galaxies with $f_{z\text{-success}} = 1$ (100% z -success) will have $w_{i,\text{ZF}} = 1.0$ while galaxies with $f_{z\text{-success}} = 0.1$ (10% z -success) will have $w_{i,\text{ZF}} = 10$. For $f_{z\text{-success}}(r_{\text{fiber},i}, \text{TSNR2}_i)$, we fit the following functional form for each TSNR2 bin:

$$f_{z\text{-success}}(r_{\text{fiber}}) = \frac{1}{2} \left(1 - \text{erf}(c_0(r_{\text{fiber}} - c_1)) \right). \quad (\text{A3})$$

In Figure 9, we present the best-fit $f_{z\text{-success}}(r_{\text{fiber}})$ for each of the TSNR2 bins in dashed. The best-fit coefficients, c_0, c_1 , are derived from χ^2 minimization. We repeat this procedure indepedently for BGS Bright galaxies as well as the BGS Faint galaxies with $(z - W1) - 1.2(g - r) + 1.2 \geq 0$, and BGS Faint galaxies with $(z - W1) - 1.2(g - r) + 1.2 < 0$. We list the best-fit values in bins of TSNR2 for each of the samples in Table 1.

B. UNCERTAINTIES ON THE SMF

We estimate the uncertainties of the SMF from sample variance using the standard jackknife technique. This involves splitting our BGS sample into subsamples and then estimating uncertainties using the subsample-to-subsample variations:

$$\sigma_{\Phi} = \left(\frac{N_{\text{jack}} - 1}{N_{\text{jack}}} \sum_{k=1}^{N_{\text{jack}}} (\Phi_k - \Phi)^2 \right). \quad (\text{B4})$$

N_{jack} is the number of jackknife subsamples and Φ_k represents the SMF estimated from the BGS galaxies excluding the jackknife subample k . In this work, we split the BGS sample into 12 jackknife fields based on the angular positions of galaxies. We present the jackknife fields in Figure 10 with distinct colors.

Table 1. Best-fit coefficients of the z -success rate as a function of r_{fiber} for different TSNR2 bins for BGS Bright and Faint samples.

TSNR2 range	c_0	c_1
BGS Bright		
$10^{1.5} - 10^2$	0.443	19.7
$10^2 - 10^{2.5}$	0.668	21.3
$10^{2.5} - 10^3$	0.888	22.1
$10^3 - 10^{3.5}$	0.822	22.4
$10^{3.5} - 10^{3.85}$	0.698	23.3
$10^{3.85} - 10^5$	0.465	21.8
BGS Faint		
$(z - W1) - 1.2(g - r) + 1.2 \geq 0$		
$10^{1.5} - 10^{2.5}$	1.67	21.1
$10^{2.5} - 10^3$	1.65	21.8
$10^3 - 10^{3.1}$	1.49	22.1
$10^{3.1} - 10^{3.2}$	1.32	22.3
$10^{3.2} - 10^{3.3}$	1.33	22.4
$10^{3.3} - 10^{3.5}$	0.907	23.1
$10^{3.5} - 10^{3.85}$	1.03	23.0
$10^{3.85} - 10^5$	0.924	21.6
BGS Faint		
$(z - W1) - 1.2(g - r) + 1.2 < 0$		
$10^{2.5} - 10^3$	1.48	20.9
$10^3 - 10^{3.1}$	2.40	21.2
$10^{3.1} - 10^{3.2}$	1.30	21.8
$10^{3.2} - 10^{3.3}$	1.27	22.0
$10^{3.3} - 10^{3.5}$	1.83	21.6
$10^{3.5} - 10^{3.85}$	0.798	22.9
$10^{3.85} - 10^5$	1.29	20.6

C. STELLAR MASS COMPLETENESS

In this appendix, we describe how we derive M_{lim} , the M_* limit above which our BGS Bright sample is complete. Although there are various methods for estimating M_{lim} in the literature, *e.g.* based on estimating the mass-to-light ratio ([Moustakas et al. 2013](#)), we adopt a simple approach that takes advantage of the fact that BGS Bright is a magnitude-limited sample.

To derive M_{lim} in redshift bins of width $\Delta z = 0.04$, we first split the galaxy sample into narrower bins of $\Delta z/2$. For each narrower redshift bin, $i\Delta z/2 < z < (i+1)\Delta z/2$, we take all the best-fit




Figure 11. The M_* distribution of BGS Bright galaxies with $0.01 < z < 0.03$ (blue) and the M_* distribution of same set of galaxies that would remain in the BGS Bright magnitude limit if they were redshifted to $z' = z + 0.02$: $r' < 19.5$. We set the stellar mass completeness limit, M_{lim} , for $0.01 < z < 0.05$ to the M_* where more than 10% of galaxies are excluded in the latter distribution.

Table 2. Stellar mass completeness limit, M_{lim} for redshift bins of width $\Delta z = 0.04$.

z range	$\log_{10} M_{\text{lim}}$
0.01 – 0.05	8.975
0.05 – 0.09	9.500
0.09 – 0.13	10.20
0.13 – 0.17	10.38
0.17 – 0.21	10.72

PROVABGS SEDs from all galaxies in the bin and artificially redshift it to $z' = z + \Delta z/2$:

$$f'_\lambda = f_\lambda \frac{d_L(z)^2}{d_L(z')^2}. \quad (\text{C5})$$

$d_L(z)$ represents the luminosity distance at redshift z . Afterward, we calculate the r -band magnitudes, r' , for f'_λ and impose the $r' < 19.5$ magnitude limit of the BGS Bright. We then compare the M_* distribution of all the galaxies in $i\Delta z/2 < z < (i+1)\Delta z/2$ to the galaxies in $i\Delta z/2 < z < (i+1)\Delta z/2$ with $r' < 19.5$. For instance, we present the M_* distributions of all BGS Bright galaxies in $0.01 < z < 0.03$ (blue) and the BGS Bright galaxies in $0.01 < z < 0.03$ with $r' < 19.5$ (orange) in Figure 11.

Since galaxies become fainter when they are placed at higher redshifts, *i.e.* $r' > r$, the $r' < 19.5$ sample has fewer low M_* galaxies. We determine the M_* at which, more than 10% of galaxies are excluded in the $r' < 19.5$ sample (black dashed) and set this limit as M_{lim} for the redshift bins: $0.01 < z < 0.05$. Our procedure for deriving M_{lim} takes advantage of the fact that galaxy samples at lower redshifts are complete down to lower M_* than at higher redshifts. We repeat this procedure




Figure 12. M_* and redshift relation of BGS Bright galaxies in the EDR (black) and the galaxies within the stellar mass completeness limit ($M_* < M_{\text{lim}}$; blue). M_{lim} is derived in redshift bins of width $\Delta z = 0.04$. The lowest redshift bin ($0.01 < z < 0.05$) is complete down to $M_* < 10^9 M_\odot$.

for all the $\Delta z = 0.04$ redshift bins that we use to measure the SMF. In Table 2, we list M_{lim} values for each of the redshift bins. Furthermore, we present the M_* and redshift relation of BGS Bright galaxies (black) and the stellar mass complete sample (blue) in Figure 12.

REFERENCES

- Baronchelli L., Nandra K., Buchner J., 2020, *Monthly Notices of the Royal Astronomical Society*, 498, 5284
- Bernardi M., Meert A., Sheth R. K., Fischer J. L., Huertas-Company M., Maraston C., Shankar F., Vikram V., 2017, *Monthly Notices of the Royal Astronomical Society*, 467, 2217
- Blanton M. R., Moustakas J., 2009, *Annual Review of Astronomy and Astrophysics*, 47, 159
- Chabrier G., 2003, *Publications of the Astronomical Society of the Pacific*, 115, 763
- Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *The Astrophysical Journal*, 823, 102
- Coil A. L., et al., 2011, *The Astrophysical Journal*, 741, 8
- Conroy C., Gunn J. E., White M., 2009, *The Astrophysical Journal*, 699, 486
- Conroy C., White M., Gunn J. E., 2010, *The Astrophysical Journal*, 708, 58
- Daddi E., et al., 2007, *The Astrophysical Journal*, 670, 156
- Donnari M., et al., 2019, *Monthly Notices of the Royal Astronomical Society*, 485, 4817
- Dotter A., 2016, *The Astrophysical Journal Supplement Series*, 222, 8
- Driver S. P., et al., 2011, *Monthly Notices of the Royal Astronomical Society*, 413, 971
- Driver S. P., et al., 2022, *Monthly Notices of the Royal Astronomical Society*, p. stac472
- Foreman-Mackey D., Hogg D. W., Morton T. D., 2014, *The Astrophysical Journal*, 795, 64
- Genel S., et al., 2014, *Monthly Notices of the Royal Astronomical Society*, 445, 175
- Hahn C., et al., 2019, *The Astrophysical Journal*, 872, 160

- Hahn C., et al., 2022, The DESI PRObabilistic Value-Added Bright Galaxy Survey (PROVABGS) Mock Challenge
- Henriques B. M. B., Thomas P. A., Oliver S., Roseboom I., 2009, [Monthly Notices of the Royal Astronomical Society](#), 396, 535
- Henriques B. M. B., White S. D. M., Thomas P. A., Angulo R., Guo Q., Lemson G., Springel V., Overzier R., 2015, [Monthly Notices of the Royal Astronomical Society](#), 451, 2663
- Hogg D. W., Myers A. D., Bovy J., 2010, [The Astrophysical Journal](#), 725, 2166
- Karamanis M., Beutler F., 2020, arXiv e-prints, p. arXiv:2002.06212
- Kingma D. P., Ba J., 2017, arXiv:1412.6980 [cs]
- Kwon K. J., Hahn C., Alsing J., 2022, Neural Stellar Population Synthesis Emulator for the DESI PROVABGS
- Leja J., Speagle J. S., Johnson B. D., Conroy C., van Dokkum P., Franx M., 2019, arXiv, p. arXiv:1910.04168
- Lejeune T., Cuisinier F., Buser R., 1997, [A & A Supplement series, Vol. 125, October II 1997, p.229-246.](#), 125, 229
- Lejeune T., Cuisinier F., Buser R., 1998, [Astronomy and Astrophysics Supplement, v.130, p.65-75](#), 130, 65
- Lu Y., et al., 2014, [The Astrophysical Journal](#), 795, 123
- Malz A. I., Hogg D. W., 2020, How to Obtain the Redshift Distribution from Probabilistic Redshift Estimates
- McLachlan G., Peel D., 2000, Finite Mixture Models. Wiley-Interscience
- Moustakas J., et al., 2013, [The Astrophysical Journal](#), 767, 50
- Muzzin A., et al., 2013, [The Astrophysical Journal](#), 777, 18
- Nelson D., et al., 2015, [Astronomy and Computing](#), 13, 12
- Noeske K. G., et al., 2007, [The Astrophysical Journal Letters](#), 660, L43
- Norberg P., Baugh C. M., Gaztañaga E., Croton D. J., 2009, [Monthly Notices of the Royal Astronomical Society](#), 396, 19
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, [The Astrophysical Journal Supplement Series](#), 192, 3
- Paxton B., et al., 2013, [The Astrophysical Journal Supplement Series](#), 208, 4
- Paxton B., et al., 2015, [The Astrophysical Journal Supplement Series](#), 220, 15
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1992, Numerical Recipes in C (2Nd Ed.): The Art of Scientific Computing. Cambridge University Press, New York, NY, USA
- Salim S., et al., 2007, [The Astrophysical Journal Supplement Series](#), 173, 267
- Sánchez-Blázquez P., et al., 2006, [Monthly Notices of the Royal Astronomical Society](#), 371, 703
- Somerville R. S., Davé R., 2015, [Annual Review of Astronomy and Astrophysics](#), 53, 51
- Somerville R. S., Popping G., Trager S. C., 2015, [Monthly Notices of the Royal Astronomical Society](#), 453, 4337
- Speagle J. S., Steinhardt C. L., Capak P. L., Silverman J. D., 2014, [The Astrophysical Journal Supplement Series](#), 214, 15
- Vogelsberger M., et al., 2014, [Monthly Notices of the Royal Astronomical Society](#), 444, 1518
- Westera P., Lejeune T., Buser R., Cuisinier F., Bruzual G., 2002, [Astronomy and Astrophysics](#), 381, 524
- York D. G., et al., 2000, [The Astronomical Journal](#), 120, 1579