

CAUSAL INFERENCE WITH ORDINAL OUTCOMES

DENSITY ESTIMATION BASED APPROACH

CHANHYUK PARK

WASHINGTON UNIVERSITY IN ST. LOUIS

MOTIVATION: THE DISCRETIZATION PROBLEM


- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
 - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
 - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it

MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
 - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
 - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it


MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
 - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
 - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it




MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
 - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
 - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it



MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
 - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
 - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it



THE GOAL

- *Identification*
- *Estimation*

THE GOAL

- *Identification*

Naive causal identification with these "index" may fail

⇒ **Normalized Latent Treatment Effect**

- *Estimation*

THE GOAL

- *Identification*

Naive causal identification with these "index" may fail

⇒ Normalized Latent Treatment Effect

- *Estimation*

Standard parametric regression is rigid

⇒ Flexible, density estimation based estimators

WHY NAIVE IDENTIFICATION FAILS

Preference over Political System (Y^*)



- The treatment effect we want: $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function g*
- *Arbitrary Numerical Index f*
- What we get: $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$


WHY NAIVE IDENTIFICATION FAILS

Preference over Political System (Y^*)




- The treatment effect we want: $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function g*
- *Arbitrary Numerical Index f*
- What we get: $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

WHY NAIVE IDENTIFICATION FAILS




- The treatment effect we want: $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function g*
- *Arbitrary Numerical Index f*
- What we get: $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

WHY NAIVE IDENTIFICATION FAILS



- The treatment effect we want: $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function g*
- *Arbitrary Numerical Index f*
- What we get: $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

WHY NAIVE IDENTIFICATION FAILS



- The treatment effect we want: $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function g*
- *Arbitrary Numerical Index f*
- What we get: $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 - ⇒ identify the Treatment Effect up to scale
 - ⇒ treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_j} = 1$
 - ② If there are multiple treatments, fixing $\tau_0 = 1$
 - ③ Pure probability scale

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 - ⇒ identify the Treatment Effect up to scale
 - ⇒ treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_i} = 1$
 - ② If there are multiple treatments, fixing $\tau_a = 1$
 - ③ Pure probability scale

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 - ⇒ identify the Treatment Effect up to scale
 - ⇒ treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_i} = 1$
 - ② If there are multiple treatments, fixing $\tau_a = 1$
 - ③ Pure probability scale

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 - ⇒ identify the Treatment Effect up to scale
 - ⇒ treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_i} = 1$
 - ② If there are multiple treatments, fixing $\tau_a = 1$
 - ③ Pure probability scale

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 - ⇒ identify the Treatment Effect up to scale
 - ⇒ treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_i} = 1$
 - ② If there are multiple treatments, fixing $\tau_a = 1$
 - ③ Pure probability scale

NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
 \implies identify the Treatment Effect up to scale
 \implies treatment effect τ^* cannot be distinguished from $C\tau^*$
- Normalization to cancel out C
 - ① Scale to fix $\sigma_{\varepsilon_i} = 1$
 $\implies \frac{\tau^*}{\sigma_{\varepsilon_i}}$ is identified
 - ② If there are multiple treatments, fixing $\tau_a = 1$
 - ③ Pure probability scale

HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models
 \implies Assume that ε_j follows specific distributions
- Distributional Assumptions can be violated
 - Pure misspecification
 - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models
 \implies Assume that ε_j follows specific distributions
- Distributional Assumptions can be violated
 - Pure misspecification
 - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models
 \implies Assume that ε_j follows specific distributions
- Distributional Assumptions can be violated
 - Pure misspecification
 - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models
 \implies Assume that ε_j follows specific distributions
- Distributional Assumptions can be violated
 - Pure misspecification
 - Unaccounted confounder
- Become *inconsistent*
- **Two Flexible Estimators** without rigid Distributional Assumptions

ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: **Kernel Density Estimation**
- Parametric Generative Model: **Normalizing Flows**
- Plug into MLE

$$\ell(\alpha, \beta, \tau) = \sum_{i=1}^n \sum_{j=0}^J \mathbf{1}_{Y_i=j} \log [\hat{F}_{\varepsilon_i}(D_i, \tau_i) \hat{F}_{\varepsilon_i}(D_i, \tau_i)]$$

ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: **Kernel Density Estimation**
 - Smooth each observation using a kernel (usually Gaussian)

$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right)$$

- Parametric Generative Model: **Normalizing Flows**
- Plug into MLE

$$\ell(\alpha, \beta, \tau) = \sum_{i=1}^n \sum_{j=0}^J 1_{Y_i=j} \log [\hat{F}_{\epsilon_i}(D_i, \tau_i) \hat{F}_{\epsilon_i}(D_i, \tau_i)]$$

ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: **Kernel Density Estimation**
 - Smooth each observation using a kernel (usually Gaussian)

$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right)$$

- Parametric Generative Model: **Normalizing Flows**
 - Use the change-of-variable formula

$$f_{\theta}(\varepsilon) = f_Z(T_{\theta}^{-1}(\varepsilon)) \left| \det \left(\frac{\partial T_{\theta}^{-1}(\varepsilon)}{\partial \varepsilon} \right) \right|,$$

- T_{θ} is a set of *invertible* transformation


- Plug into MLE

ALTERNATIVE: ESTIMATE THE DISTRIBUTION



- Nonparametric method: **Kernel Density Estimation**
- Parametric Generative Model: **Normalizing Flows**
- Plug into MLE

$$\ell(\alpha, \beta, \tau) = \sum_{i=1}^n \sum_{j=0}^J \mathbf{1}_{Y_i=j} \log [\hat{F}_{\varepsilon_i}(D_i, \tau_i) \hat{F}_{\varepsilon_i}(D_i, \tau_i)]$$

SIMULATION



SIMULATION – LOGNORMAL CASE



True treatment size: 0.46

REPLICATION – MATTINGLY ET AL. (2025)

- Effect of promotion videos on preference over political system
- *Strongly Prefer the US to Strongly Prefer China*

	ATE	NLTE			
	Original – OLS	Ordered Logit	Ordered Probit	KDE-based	NF-based
China	1.04*** (0.05)	0.73*** (0.04)	0.76*** (0.04)	0.36*** (0.07)	0.37*** (0.04)
USA	-0.43*** (0.04)	-0.35*** (0.04)	-0.38*** (0.04)	-0.18*** (0.05)	-0.17*** (0.03)
Competition	0.36*** (0.05)	0.21*** (0.04)	0.25*** (0.04)	0.13* (0.06)	0.11** (0.04)

*** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$

CONCLUSION

- *Identification*
- *Estimation*

CONCLUSION

- *Identification*

Naive use of ordinal "index" is not what we want

⇒ **Normalized Latent Treatment Effect**

- *Estimation*

CONCLUSION

- *Identification*

Naive use of ordinal "index" is not what we want

⇒ Normalized Latent Treatment Effect

- *Estimation*

Standard ordinal regressions risk inconsistency

⇒ KDE or NF based estimators