

# CAUSAL INFERENCE WITH ORDINAL OUTCOMES

---

DENSITY ESTIMATION BASED APPROACH

CHANHYUK PARK

WASHINGTON UNIVERSITY IN ST. LOUIS

## MOTIVATION: THE DISCRETIZATION PROBLEM


- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
  - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
  - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it

# MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
  - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
  - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it


# MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
  - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
  - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it




# MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
  - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
  - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it



# MOTIVATION: THE DISCRETIZATION PROBLEM

- Experiments to identify causal effects of treatments
- Outcomes are usually believed to be continuous in unidimensional space
  - Approval ratings (Canes-Wrone and De Marchi, 2002; Kriner and Schwartz, 2009)
  - Policy preferences (Scheve and Slaughter, 2001; Mayda and Rodrik, 2005; Wu, 2022)
- The Problem - How we measure it



# THE GOAL

- *Identification*
- *Estimation*

# THE GOAL

- *Identification*

Naive causal identification with these "index" may fail

⇒ **Normalized Latent Treatment Effect**

- *Estimation*



## THE GOAL

- *Identification*

Naive causal identification with these "index" may fail

⇒ Normalized Latent Treatment Effect

- *Estimation*

Standard parametric regression is rigid

⇒ Flexible, density estimation based estimators

# WHY NAIVE IDENTIFICATION FAILS

**Preference over Political System ( $Y^*$ )**



- The treatment effect we want:  $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function  $g$*
- *Arbitrary Numerical Index  $f$*
- What we get:  $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$


## WHY NAIVE IDENTIFICATION FAILS

**Preference over Political System ( $Y^*$ )**




- The treatment effect we want:  $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function  $g$*
- *Arbitrary Numerical Index  $f$*
- What we get:  $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

## WHY NAIVE IDENTIFICATION FAILS




- The treatment effect we want:  $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function  $g$*
- *Arbitrary Numerical Index  $f$*
- What we get:  $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

## WHY NAIVE IDENTIFICATION FAILS



- The treatment effect we want:  $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function  $g$*
- *Arbitrary Numerical Index  $f$*
- What we get:  $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

## WHY NAIVE IDENTIFICATION FAILS



- The treatment effect we want:  $\mathbb{E}[Y^*(1)] - \mathbb{E}[Y^*(0)] = \tau^*$
- *Unknown Reporting Function  $g$*
- *Arbitrary Numerical Index  $f$*
- What we get:  $\mathbb{E}[f(g(Y^*(1)))] - \mathbb{E}[f(g(Y^*(0)))] \neq \tau^*$

## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
  - ⇒ identify the Treatment Effect up to scale
  - ⇒ treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_j} = 1$
  - ② If there are multiple treatments, fixing  $\tau_0 = 1$
  - ③ Pure probability scale

## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
  - ⇒ identify the Treatment Effect up to scale
  - ⇒ treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_i} = 1$
  - ② If there are multiple treatments, fixing  $\tau_a = 1$
  - ③ Pure probability scale



## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
  - ⇒ identify the Treatment Effect up to scale
  - ⇒ treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_i} = 1$
  - ② If there are multiple treatments, fixing  $\tau_a = 1$
  - ③ Pure probability scale

## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
  - ⇒ identify the Treatment Effect up to scale
  - ⇒ treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_i} = 1$
  - ② If there are multiple treatments, fixing  $\tau_a = 1$
  - ③ Pure probability scale

## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers
  - ⇒ identify the Treatment Effect up to scale
  - ⇒ treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_i} = 1$
  - ② If there are multiple treatments, fixing  $\tau_a = 1$
  - ③ Pure probability scale

## NEW ESTIMANDS: NORMALIZED TREATMENT EFFECT

- Nonparametrically identified probability distribution of each answers  
     $\implies$  identify the Treatment Effect up to scale  
     $\implies$  treatment effect  $\tau^*$  cannot be distinguished from  $C\tau^*$
- Normalization to cancel out  $C$ 
  - ① Scale to fix  $\sigma_{\varepsilon_i} = 1$   
     $\implies \frac{\tau^*}{\sigma_{\varepsilon_i}}$  is identified
  - ② If there are multiple treatments, fixing  $\tau_a = 1$
  - ③ Pure probability scale

## HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models  
     $\implies$  Assume that  $\varepsilon_j$  follows specific distributions
- Distributional Assumptions can be violated
  - Pure misspecification
  - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

## HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models  
     $\implies$  Assume that  $\varepsilon_j$  follows specific distributions
- Distributional Assumptions can be violated
  - Pure misspecification
  - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

## HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models  
     $\implies$  Assume that  $\varepsilon_j$  follows specific distributions
- Distributional Assumptions can be violated
  - Pure misspecification
  - Unaccounted confounder
- Become *inconsistent*
- Two Flexible Estimators without rigid Distributional Assumptions

## HOW CAN WE ESTIMATE?

- Ordered Probit, Ordered Logit, and most IRT models  
     $\implies$  Assume that  $\varepsilon_j$  follows specific distributions
- Distributional Assumptions can be violated
  - Pure misspecification
  - Unaccounted confounder
- Become *inconsistent*
- **Two Flexible Estimators** without rigid Distributional Assumptions



## ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: Kernel Density Estimation
- Parametric Generative Model: Normalizing Flows

## ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: **Kernel Density Estimation**
  - Smooth each observation using a kernel (usually Gaussian)

$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right)$$

- Parametric Generative Model: **Normalizing Flows**

## ALTERNATIVE: ESTIMATE THE DISTRIBUTION

- Nonparametric method: **Kernel Density Estimation**
  - Smooth each observation using a kernel (usually Gaussian)


$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right)$$

- Parametric Generative Model: **Normalizing Flows**
  - Use the change-of-variable formula



$$f_{\theta}(\varepsilon) = f_Z(T_{\theta}^{-1}(\varepsilon)) \left| \det \left( \frac{\partial T_{\theta}^{-1}(\varepsilon)}{\partial \varepsilon} \right) \right|,$$

- $T_{\theta}$  is a set of *invertible* transformation

# SIMULATION



# SIMULATION – LOGNORMAL CASE



True treatment size: 0.46

## REPLICATION – MATTINGLY ET AL. (2025)

- Effect of promotion videos on preference over political system
- *Strongly Prefer the US to Strongly Prefer China*

	ATE	NLTE			
	Original – OLS	Ordered Logit	Ordered Probit	KDE-based	NF-based
China	1.04*** (0.05)	0.73*** (0.04)	0.76*** (0.04)	0.36*** (0.07)	0.37*** (0.04)
USA	–0.43*** (0.04)	–0.35*** (0.04)	–0.38*** (0.04)	–0.18*** (0.05)	–0.17*** (0.03)
Competition	0.36*** (0.05)	0.21*** (0.04)	0.25*** (0.04)	0.13* (0.06)	0.11** (0.04)

\*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; \* $p < 0.05$

## CONCLUSION

- *Identification*
- *Estimation*

# CONCLUSION

- *Identification*

Naive use of ordinal "index" is not what we want

⇒ **Normalized Latent Treatment Effect**

- *Estimation*



# CONCLUSION

- *Identification*

Naive use of ordinal "index" is not what we want

⇒ Normalized Latent Treatment Effect

- *Estimation*

Standard ordinal regressions risk inconsistency

⇒ KDE or NF based estimators