AMC12B 2024 Problem 19
Strategic Problem Analysis

Competition Math Framework v2.1

October 26, 2025

Problem Statement

Equilateral $\triangle ABC$ with side length 14 is rotated about its center by angle θ, where $0 < \theta < 60°$, to form $\triangle DEF$. The area of hexagon $ADBECF$ is $91\sqrt{3}$. What is $\tan\theta$?



(A) $\frac{3}{4}$ **(B)** $\frac{5\sqrt{3}}{11}$ **(C)** $\frac{4}{5}$ **(D)** $\frac{11}{13}$ **(E)** $\frac{7\sqrt{3}}{13}$

Note: The answer is not revealed here to allow students to work through the problem. See the Final Answer section at the end.

PROBLEM CONFIGURATION ANALYSIS

A. Problem Classification

- **Problem Type**: To Find (specific trigonometric value)
- **Competition Context**: AMC12B 2024, Problem 19
- **Difficulty Band**: Late-band (P19) - Requires integration of geometry and trigonometry
- **Mathematical Domain**: Geometry (equilateral triangles, rotation, hexagon), Trigonometry (angle relationships, identities)
- **Estimated Time**: 5-7 minutes
- **Point Value**: 6 points (standard AMC12 scoring)

B. Problem Structure Identification

- **Given Information**:
  - Equilateral triangle ABC with side length 14
  - Triangle rotated about center by angle θ where $0 < \theta < 60°$
  - Rotation creates triangle DEF
  - Hexagon $ADBECF$ formed by intersection has area $91\sqrt{3}$
  - Area of original triangle: $\frac{\sqrt{3}}{4} \cdot 14^2 = 49\sqrt{3}$
- **Constraints**:
  - $0 < \theta < 60°$ (rotation angle constraint)
  - Both triangles are equilateral with same side length
  - Both share the same center
  - Hexagon area is specified: $91\sqrt{3}$
- **Objective**: Find $\tan\theta$
- **Hidden Assumptions**:
  - The hexagon is formed by the overlapping region
  - Symmetry: the hexagon has 6 congruent triangular regions
  - The circumradius $R = \frac{14\sqrt{3}}{3}$ (for equilateral triangle)
  - Small triangular corners are cut off from the original triangle
- **Answer Choice Leverage**:

- Three choices contain $\sqrt{3}$: (B), (E)
- Two choices are rational: (A), (C), (D)
- All values are between 0.75 and 1.1, consistent with $0 < \theta < 60°$
- Can test answer choices if stuck

- **Proof Requirements**: N/A (computational problem)

C. Strategic Orientation

- **Hypothesis**: Use area relationships and trigonometry to find θ
- **Conclusion**: Need exact value of $\tan\theta$
- **Notation**:
  - O = center (circumcenter) of triangles
  - $R = \frac{14\sqrt{3}}{3}$ = circumradius
  - $[\cdot]$ = area notation
  - $[ABC] = 49\sqrt{3}$ (original triangle area)
  - $[ADBECF] = 91\sqrt{3}$ (hexagon area)
- **Intuitive Approach**:
  1. Relate hexagon area to triangle area and rotation angle
  2. Use symmetry: hexagon can be divided into 6 smaller triangular regions
  3. Set up trigonometric equations using area formulas
  4. Solve for $\sin\theta$ and $\cos\theta$, then find $\tan\theta$
- **Cross-Domain Potential**:
  - Geometry $\rightarrow$ Trigonometry (angle relationships)
  - Area methods $\rightarrow$ Algebraic equations
  - Coordinate geometry possible but likely more complex

STRATEGIC FRAMEWORK APPLICATION

A. Psychological Strategies

- **Mental Toughness**: This is a P19 problem combining geometry and trigonometry. Multiple solution paths exist - if one approach stalls, pivot to another.
- **Creativity Cultivation**: Visualize the rotation. The hexagon is formed by 6 triangular "petals" around the center. Look for symmetry to simplify calculations.

- **Iterative Refinement**: Start with area relationships, then use trigonometric identities to isolate θ.

B. Investigation Strategies

- **Startup Orientation**: Draw a careful diagram showing both triangles and the hexagon. Mark the center O.
- **Get Your Hands Dirty**:
  - Calculate circumradius: $R = \frac{14}{\sqrt{3}} = \frac{14\sqrt{3}}{3}$
  - Compute triangle area: $[ABC] = 49\sqrt{3}$
  - Observe: Hexagon area $91\sqrt{3} > [ABC] = 49\sqrt{3}$, which seems wrong!
  - Actually, we need to understand the hexagon better...
- **Penultimate Step**: Once we have $\sin\theta$ and $\cos\theta$, compute $\tan\theta = \frac{\sin\theta}{\cos\theta}$
- **Make It Easier**: Use symmetry - the hexagon has 3-fold rotational symmetry, so compute area of one sector
- **Conjecture via Small Cases**: Consider limiting cases:
  - If $\theta = 0$, triangles coincide, hexagon area = triangle area
  - If $\theta = 60°$, triangles coincide again
  - Maximum hexagon area occurs somewhere in between
- **Visualization**: The hexagon consists of 6 identical triangular regions emanating from center O

C. Late-Band Strategic Playbook

- **Wishful Thinking**: "I wish I could express the hexagon area in terms of θ..." $\Rightarrow$ Use triangular decomposition from center
- **Pattern Recognition**: Hexagon formed by rotation suggests using central angles and areas of triangular sectors
- **Answer Choice Testing**: If algebraic approach stalls, test answer choices by working backwards

D. Argument Construction Strategy

- **Direct Approach**: Set up area equation involving θ, solve using trigonometric identities
- **Decomposition**: Break hexagon into 6 triangular pieces, each with area expressible in terms of θ
- **Symmetry Exploitation**: Use 3-fold rotational symmetry to simplify calculations

E. Cross-Domain Translation

- **Geometric to Trigonometric**: Translate area relationships into trigonometric equations
- **Visual to Algebraic**: Use diagram to identify key angles and distances, then algebraize

TACTICAL METHODOLOGY SELECTION

A. Core Mathematical Tactics

- **Primary Tactics**:
  - **Symmetry Exploitation**: Use 3-fold rotational symmetry of configuration
  - **Area Decomposition**: Break complex hexagon into manageable triangular pieces
  - **Trigonometric Identities**: Use sum formulas, double angle formulas
- **Domain-Specific Tactics**:
  - **Equilateral Triangle Properties**:
    * Area: $A = \frac{\sqrt{3}}{4}s^2$
    * Circumradius: $R = \frac{s}{\sqrt{3}} = \frac{s\sqrt{3}}{3}$
    * Central angles: $120°$ between vertices
  - **Area of Triangle from Vertex and Angle**:
    * $[\triangle OPQ] = \frac{1}{2} \cdot OP \cdot OQ \cdot \sin(\angle POQ)$
    * For $OP = OQ = R$: $[\triangle OPQ] = \frac{1}{2}R^2 \sin(\angle POQ)$
  - **Trigonometric Identities**:
    * $\sin(120° - \theta) = \sin 120° \cos\theta - \cos 120° \sin\theta$
    * $= \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta$
    * $\sin^2\theta + \cos^2\theta = 1$
- **Crossover Tactics**:

- Geometric decomposition + Trigonometric calculation
- Visual symmetry + Algebraic manipulation

CONVERSION TECHNIQUES APPLICATION

A. High-Probability Techniques

- **Primary Technique**: **Area Decomposition with Trigonometry**
- **Recognition Cues**:
  - "Rotated about center" $\Rightarrow$ Use central angles
  - "Equilateral triangle" $\Rightarrow$ Use symmetry and special angles
  - "Area of hexagon" $\Rightarrow$ Decompose into triangular regions
  - "Find $\tan\theta$" $\Rightarrow$ Need to find both $\sin\theta$ and $\cos\theta$
- **Application**:
  1. Identify center O as circumcenter
  2. Calculate circumradius $R = \frac{14\sqrt{3}}{3}$
  3. Decompose hexagon into 6 triangular regions from O
  4. Express areas using $[\triangle] = \frac{1}{2}R^2 \sin(\text{angle})$
  5. Set up equation: hexagon area = sum of triangular areas
  6. Solve for $\sin\theta$ and $\cos\theta$
  7. Compute $\tan\theta$

B. Alternative Techniques

- **Coordinate Geometry Approach**:
  - Place triangle with center at origin
  - Set up coordinates for vertices
  - Apply rotation matrix
  - Calculate hexagon area using shoelace formula
  - More computational but systematic
- **Direct Trigonometric Setup**:
  - Use relationship: $[ADBECF] = 3([FOC] + [COE])$ by symmetry
  - Express each triangular area in terms of θ
  - Solve resulting trigonometric equation

- **Answer Choice Testing**:
  - For each answer, compute $\sin\theta$ and $\cos\theta$
  - Verify hexagon area equals $91\sqrt{3}$
  - Time-consuming but guaranteed to work

KEY INSIGHT

The Critical Insights:

1. **Hexagon Decomposition**: The hexagon $ADBECF$ can be viewed as consisting of 6 identical triangular "sectors" emanating from the center O. By symmetry, we can compute the area of two adjacent sectors and multiply by 3.

2. **Area Relationship**:

$$\begin{aligned}[ADBECF] &= 3 \times ([\triangle OFC] + [\triangle OCE]) \\ &= 3 \times \left(\frac{1}{2}R^2 \sin\theta + \frac{1}{2}R^2 \sin(120^\circ - \theta)\right)\end{aligned}$$

   where $R = \frac{14\sqrt{3}}{3}$ is the circumradius.

3. **Simplification Strategy**: Using $R^2 = \frac{196 \cdot 3}{9} = \frac{196}{3}$, we get:

$$[ADBECF] = \frac{3}{2} \cdot \frac{196}{3} \cdot (\sin\theta + \sin(120^\circ - \theta)) = 98(\sin\theta + \sin(120^\circ - \theta))$$

4. **Key Equation**: Setting this equal to $91\sqrt{3}$:

$$\sin\theta + \sin(120^\circ - \theta) = \frac{91\sqrt{3}}{98}$$

5. **Angle Addition Formula**: Expanding the sine:

$$\begin{aligned}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta &= \frac{91\sqrt{3}}{98} \\ \frac{3}{2}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta &= \frac{91\sqrt{3}}{98}\end{aligned}$$

   Multiplying by $\frac{2}{\sqrt{3}}$:

$$\sqrt{3}\sin\theta + \cos\theta = \frac{13}{7}$$

6. **Finding Both Components**: Use $\sin^2\theta + \cos^2\theta = 1$ with $\cos\theta = \frac{13}{7} - \sqrt{3}\sin\theta$ to get a quadratic in $\sin\theta$.

VERIFICATION STRATEGY DESIGN

A. Verification Level Selection

- **Chosen Level**: Level 5 (Thorough Verification) - 1-2 minutes
- **Justification**:
  - Problem 19 is late-band with complex trigonometry
  - Multiple algebraic steps prone to error
  - Need to verify both $\sin\theta$ and $\cos\theta$ satisfy constraints
  - Should verify $\tan\theta$ matches an answer choice
  - Can verify by computing hexagon area with found angle
- **Time Budget**: 1-2 minutes for thorough verification

B. Verification Checklist

- **Solution Validity**:
  - Check: $0 < \sin\theta < \sin 60^\circ = \frac{\sqrt{3}}{2}$
  - Verify: $\sin\theta = \frac{5\sqrt{3}}{14} \approx 0.619 < 0.866$ ✓
  - Check: $\cos\theta > 0$ for $0 < \theta < 60^\circ$
  - Verify: $\cos\theta = \frac{11}{14} > 0$ ✓
  - Check: $\sin^2\theta + \cos^2\theta = 1$
  - Verify: $\left(\frac{5\sqrt{3}}{14}\right)^2 + \left(\frac{11}{14}\right)^2 = \frac{75}{196} + \frac{121}{196} = \frac{196}{196} = 1$ ✓
- **Equation Verification**:
  - Verify: $\sqrt{3}\sin\theta + \cos\theta = \sqrt{3} \cdot \frac{5\sqrt{3}}{14} + \frac{11}{14} = \frac{15}{14} + \frac{11}{14} = \frac{26}{14} = \frac{13}{7}$ ✓
- **Answer Format**:
  - Compute: $\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{5\sqrt{3}/14}{11/14} = \frac{5\sqrt{3}}{11}$
  - This matches choice (B) ✓
- **Alternative Verification**:
  - Work backwards: compute hexagon area using θ with $\tan\theta = \frac{5\sqrt{3}}{11}$
  - Should get $91\sqrt{3}$

C. Domain-Specific Verification

- **Geometry Verification**:
  - Circumradius formula: $R = \frac{14\sqrt{3}}{3}$ ✓
  - Triangle area: $\frac{\sqrt{3}}{4} \cdot 14^2 = 49\sqrt{3}$ ✓
  - Hexagon should have area between $[ABC]$ and something larger
- **Trigonometry Verification**:
  - All trig identities applied correctly
  - Quadratic solution gives two roots; verify correct one chosen
  - The other root $\sin\theta = \frac{4\sqrt{3}}{7} > \frac{\sqrt{3}}{2}$ violates $\theta < 60°$

SOLUTION ROADMAP

A. Step-by-Step Solution

Step 1: Setup and Notation
Let O be the center (circumcenter) of both equilateral triangles. The circumradius is:

$$R = \frac{\text{side}}{\sqrt{3}} = \frac{14}{\sqrt{3}} = \frac{14\sqrt{3}}{3}$$

The area of equilateral triangle ABC is:

$$[ABC] = \frac{\sqrt{3}}{4} \cdot 14^2 = 49\sqrt{3}$$

Checkpoint: We have $R = \frac{14\sqrt{3}}{3}$ and original triangle area.

Step 2: Understanding the Hexagon
When triangle ABC is rotated by angle θ about O to form triangle DEF, the vertices rotate as follows:

- $A \to D$
- $B \to E$
- $C \to F$

The hexagon $ADBECF$ is the region covered by the overlapping configuration. By symmetry, this hexagon can be decomposed into 6 triangular sectors emanating from O.
Consider the vertices in order around O. In the original triangle, the angle between consecutive vertices (e.g., $\angle AOB$) is $120°$. After rotation by θ:

- $\angle AOD = \theta$
- $\angle DOB = 120^\circ - \theta$
- By symmetry, similar angles repeat around the hexagon

Checkpoint: The hexagon area can be expressed as a sum of triangular sectors.

Step 3: Area Decomposition
By symmetry (3-fold rotational), the hexagon consists of 3 pairs of triangles. Consider one pair: $\triangle OFC$ and $\triangle OCE$.

$$[ADBECF] = 3 \times ([\triangle OFC] + [\triangle OCE])$$

For a triangle with two sides R and included angle α:

$$[\triangle] = \frac{1}{2}R^2 \sin\alpha$$

We have:

- $[\triangle OFC] = \frac{1}{2}R^2 \sin(\angle FOC) = \frac{1}{2}R^2 \sin\theta$
- $[\triangle OCE] = \frac{1}{2}R^2 \sin(\angle COE) = \frac{1}{2}R^2 \sin(120^\circ - \theta)$

Therefore:

$$\begin{aligned}[ADBECF] &= 3 \times \frac{1}{2}R^2(\sin\theta + \sin(120^\circ - \theta)) \\ &= \frac{3R^2}{2}(\sin\theta + \sin(120^\circ - \theta))\end{aligned}$$

Checkpoint: We have an expression for hexagon area in terms of θ.

Step 4: Numerical Substitution
Calculate R^2:

$$R^2 = \left(\frac{14\sqrt{3}}{3}\right)^2 = \frac{196 \cdot 3}{9} = \frac{588}{9} = \frac{196}{3}$$

Substitute:

$$\begin{aligned}91\sqrt{3} &= \frac{3}{2} \cdot \frac{196}{3} \cdot (\sin\theta + \sin(120^\circ - \theta)) \\ 91\sqrt{3} &= \frac{196}{2} \cdot (\sin\theta + \sin(120^\circ - \theta)) \\ 91\sqrt{3} &= 98 \cdot (\sin\theta + \sin(120^\circ - \theta))\end{aligned}$$

Dividing both sides by 98:

$$\sin\theta + \sin(120^\circ - \theta) = \frac{91\sqrt{3}}{98}$$

Checkpoint: We have a trigonometric equation to solve.

Step 5: Expanding the Sine
Use the angle subtraction formula:

$$\begin{aligned}\sin(120^\circ - \theta) &= \sin 120^\circ \cos\theta - \cos 120^\circ \sin\theta \\ &= \frac{\sqrt{3}}{2}\cos\theta - \left(-\frac{1}{2}\right)\sin\theta \\ &= \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta\end{aligned}$$

Substituting:

$$\begin{aligned}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta &= \frac{91\sqrt{3}}{98} \\ \frac{3}{2}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta &= \frac{91\sqrt{3}}{98}\end{aligned}$$

Multiply both sides by $\frac{2}{\sqrt{3}}$:

$$\begin{aligned}\sqrt{3}\sin\theta + \cos\theta &= \frac{2}{\sqrt{3}} \cdot \frac{91\sqrt{3}}{98} \\ &= \frac{2 \cdot 91}{98} \\ &= \frac{182}{98} \\ &= \frac{13}{7}\end{aligned}$$

Checkpoint: We have $\sqrt{3}\sin\theta + \cos\theta = \frac{13}{7}$.

Step 6: Solving for $\sin\theta$ and $\cos\theta$
From $\sqrt{3}\sin\theta + \cos\theta = \frac{13}{7}$, we get:

$$\cos\theta = \frac{13}{7} - \sqrt{3}\sin\theta$$

Using $\sin^2\theta + \cos^2\theta = 1$:

$$\begin{aligned}\sin^2\theta + \left(\frac{13}{7} - \sqrt{3}\sin\theta\right)^2 &= 1 \\ \sin^2\theta + \frac{169}{49} - \frac{26\sqrt{3}}{7}\sin\theta + 3\sin^2\theta &= 1 \\ 4\sin^2\theta - \frac{26\sqrt{3}}{7}\sin\theta + \frac{169}{49} &= 1 \\ 4\sin^2\theta - \frac{26\sqrt{3}}{7}\sin\theta + \frac{169-49}{49} &= 0 \\ 4\sin^2\theta - \frac{26\sqrt{3}}{7}\sin\theta + \frac{120}{49} &= 0\end{aligned}$$

Multiply through by 49 to clear denominators:

$$196\sin^2\theta - 182\sqrt{3}\sin\theta + 120 = 0$$

Divide by 4 for simplification:

$$49\sin^2\theta - \frac{91\sqrt{3}}{2}\sin\theta + 30 = 0$$

Multiply by 2:

$$98\sin^2\theta - 91\sqrt{3}\sin\theta + 60 = 0$$

Using the quadratic formula with $a = 98$, $b = -91\sqrt{3}$, $c = 60$:

$$\begin{aligned}\sin\theta &= \frac{91\sqrt{3} \pm \sqrt{(91\sqrt{3})^2 - 4(98)(60)}}{2(98)} \\ &= \frac{91\sqrt{3} \pm \sqrt{8281 \cdot 3 - 23520}}{196} \\ &= \frac{91\sqrt{3} \pm \sqrt{24843 - 23520}}{196} \\ &= \frac{91\sqrt{3} \pm \sqrt{1323}}{196}\end{aligned}$$

The key simplification: $1323 = 441 \cdot 3 = 21^2 \cdot 3$, so $\sqrt{1323} = 21\sqrt{3}$.
This perfect square factor makes the discriminant simplify nicely! Therefore:

$$\begin{aligned}\sin\theta &= \frac{91\sqrt{3} \pm 21\sqrt{3}}{196} \\ &= \frac{(91 \pm 21)\sqrt{3}}{196}\end{aligned}$$

This gives us two solutions:

- $\sin\theta = \frac{112\sqrt{3}}{196} = \frac{4\sqrt{3}}{7} = \frac{8\sqrt{3}}{14}$
- $\sin\theta = \frac{70\sqrt{3}}{196} = \frac{5\sqrt{3}}{14}$

Checking the constraint $0 < \theta < 60°$:
For $\theta < 60°$, we need $\sin\theta < \sin 60° = \frac{\sqrt{3}}{2} = \frac{7\sqrt{3}}{14}$.

- First root: $\frac{8\sqrt{3}}{14} > \frac{7\sqrt{3}}{14}$ **(INVALID - violates $\theta < 60°$)**
- Second root: $\frac{5\sqrt{3}}{14} < \frac{7\sqrt{3}}{14}$ ✓**(VALID)**

Therefore, $\sin\theta = \frac{5\sqrt{3}}{14}$.
Now find $\cos\theta$:

$$\cos\theta = \frac{13}{7} - \sqrt{3}\cdot\frac{5\sqrt{3}}{14} = \frac{13}{7} - \frac{15}{14} = \frac{26-15}{14} = \frac{11}{14}$$

Check: $\sin^2\theta + \cos^2\theta = \frac{75}{196} + \frac{121}{196} = \frac{196}{196} = 1$ ✓
Check constraint: $\sin 60^\circ = \frac{\sqrt{3}}{2} = \frac{7\sqrt{3}}{14}$, and $\frac{5\sqrt{3}}{14} < \frac{7\sqrt{3}}{14}$ ✓
The other solution $\sin\theta = \frac{4\sqrt{3}}{7} = \frac{8\sqrt{3}}{14} > \frac{7\sqrt{3}}{14}$ violates $\theta < 60^\circ$.
Checkpoint: We have $\sin\theta = \frac{5\sqrt{3}}{14}$ and $\cos\theta = \frac{11}{14}$.

Step 7: Computing $\tan\theta$

$$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{5\sqrt{3}/14}{11/14} = \frac{5\sqrt{3}}{11}$$

Final Answer: $\tan\theta = \frac{5\sqrt{3}}{11}$, which is choice **(B)**.

B. Alternative Solution (Using Area Relationship)

The hexagon area can also be computed as:

$$[ADBECF] = [\triangle ABC] + [\triangle DEF] - 3[\text{small corner triangles}]$$

But since both triangles have the same area and the configuration is symmetric, we can use:

$$[ADBECF] = 2[\triangle ABC] - [\text{non-overlapping regions}]$$

This leads to the same system of equations as above.

C. Verification of Answer

With $\tan\theta = \frac{5\sqrt{3}}{11}$:

- $\sin\theta = \frac{5\sqrt{3}}{14}$ (using $\sin^2 + \cos^2 = 1$ and $\tan = \frac{\sin}{\cos}$)
- $\cos\theta = \frac{11}{14}$
- Check: $\sqrt{3}\sin\theta + \cos\theta = \sqrt{3}\cdot\frac{5\sqrt{3}}{14} + \frac{11}{14} = \frac{15+11}{14} = \frac{26}{14} = \frac{13}{7}$ ✓

WARNING: COMMON PITFALL

Common Pitfalls to Avoid:

1. **Incorrect hexagon interpretation**: The hexagon $ADBECF$ is NOT the intersection of the two triangles, but rather the union or a specific configuration.

Carefully understand the geometry from the diagram.

2. **Wrong angle identification**: The angle θ is the rotation angle from O, not an angle in the hexagon itself. Make sure to use $\angle FOC = \theta$ correctly.

3. **Circumradius calculation error**: For an equilateral triangle with side s, the circumradius is $R = \frac{s\sqrt{3}}{3}$, NOT $\frac{s}{2}$ (which is the inradius times 2).

4. **Trigonometric identity mistakes**:
   - $\sin(120^\circ - \theta) = \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta$ (be careful with signs!)
   - Don't forget $\cos 120^\circ = -\frac{1}{2}$ (negative!)

5. **Selecting wrong root**: The quadratic gives two solutions for $\sin\theta$. Must check which satisfies $0 < \theta < 60^\circ$, i.e., $0 < \sin\theta < \frac{\sqrt{3}}{2}$.

6. **Arithmetic errors**: With $\sqrt{3}$ appearing frequently, arithmetic mistakes are common. Double-check each step.

7. **Area formula confusion**: Remember $[\triangle] = \frac{1}{2}ab\sin C$ where C is the included angle between sides a and b.

8. **Symmetry misapplication**: The hexagon has 3-fold, not 6-fold symmetry in the area calculation. Factor of 3, not 6.

PRO TIP

Competition Tips:

1. **Draw a clear diagram**: This problem is much easier with a good visualization. Mark the center O, label all vertices, and show the rotation clearly.

2. **Use symmetry aggressively**: The 3-fold rotational symmetry means you only need to compute 2 triangular sectors, then multiply by 3. This saves time and reduces errors.

3. **Circumradius quick formula**: For equilateral triangle with side s: $R = \frac{s}{\sqrt{3}} = \frac{s\sqrt{3}}{3}$. Memorize this!

4. **Angle relationships**: In an equilateral triangle, vertices are separated by 120° central angles. After rotation by θ, the new angles are θ and $120^\circ - \theta$.

5. **Simplification strategy**: When you get $\frac{3}{2}\sin\theta + \frac{\sqrt{3}}{2}\cos\theta = \frac{91\sqrt{3}}{98}$, immediately factor out $\frac{1}{2}$ and multiply by $\frac{2}{\sqrt{3}}$ to get integer coefficients.

6. **Answer choice hint**: Three choices contain $\sqrt{3}$. For rotation of equilateral triangle, $\sqrt{3}$ in the answer suggests sin and cos values involve $\sqrt{3}$.

7. **Verification shortcut**: Once you find $\sin\theta$ and $\cos\theta$, verify $\sin^2 + \cos^2 = 1$ before computing $\tan\theta$. This catches errors early.

8. **Calculator usage**: If allowed, use calculator for:

   - Computing $R^2 = \frac{196}{3}$
   - Verifying $\sin^2 + \cos^2 = 1$
   - Checking numerical approximations

   But do symbolic manipulation by hand to avoid rounding errors.

9. **Time management**: This is a 5-7 minute problem. If stuck after 3 minutes, consider answer choice testing or moving on and returning later.

COMPETITION STRATEGY INTEGRATION

A. Time Management

- **Estimated Time**: 5-7 minutes total
  - Setup and diagram: 1 minute
  - Area decomposition: 1.5 minutes
  - Trigonometric manipulation: 2 minutes
  - Solving for $\sin\theta$ and $\cos\theta$: 1.5 minutes
  - Computing $\tan\theta$ and verification: 1 minute
- **Time Checkpoints**:
  - At 1.5 min: Should have setup complete with $R = \frac{14\sqrt{3}}{3}$
  - At 3 min: Should have equation $\sin\theta + \sin(120^\circ - \theta) = \frac{91\sqrt{3}}{98}$
  - At 5 min: Should have $\sqrt{3}\sin\theta + \cos\theta = \frac{13}{7}$
  - At 6.5 min: Should have $\tan\theta = \frac{5\sqrt{3}}{11}$
- **Priority Level**: **Medium-High** - This is P19, challenging but doable with systematic approach. Worth attempting if comfortable with geometry and trigonometry.

B. Risk Assessment

- **Confidence Level**: 85-90% after verification
  - Clear geometric setup
  - Systematic trigonometric manipulation
  - Answer verifiable through multiple checks

  - Matches answer choice

- **Common Error Sources**:
  - Circumradius calculation ($R = \frac{s\sqrt{3}}{3}$, not $\frac{s}{2}$)
  - Angle addition formula signs
  - Quadratic solution selection (which root is valid?)
  - Arithmetic with $\sqrt{3}$ terms

- **Abandonment Criteria**:
  - If setup is unclear after 2 minutes $\Rightarrow$ try answer choice testing
  - If trigonometry becomes very messy after 5 minutes $\Rightarrow$ flag and move on
  - If can't solve quadratic after 6 minutes $\Rightarrow$ make educated guess from answer choices

- **Flag for Review**: Yes if time permits - double-check the quadratic solution and angle constraint

C. Strategic Context

- **Problem Positioning**: P19 is genuinely challenging but follows standard late-band patterns: geometric setup + trigonometric calculation.

- **Score Impact**: Worth 6 points. Good ROI if you're comfortable with geometry and trigonometry.

- **Time Budget Implications**: 5-7 minutes is reasonable. Don't exceed 8 minutes.

- **Technique Practice**: This problem tests:
  - Geometric visualization and symmetry
  - Area decomposition techniques
  - Trigonometric identities and manipulation
  - Solving systems with sin and cos
  - Quadratic solving with symbolic coefficients

Final Answer

From the area relationship and trigonometric manipulation, we found:

$$\sin\theta = \frac{5\sqrt{3}}{14}$$
$$\cos\theta = \frac{11}{14}$$

Therefore:

$$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{5\sqrt{3}/14}{11/14} = \frac{5\sqrt{3}}{11}$$

$$\boxed{\textbf{(B)}\ \frac{5\sqrt{3}}{11}}$$

Confidence Assessment

- **Confidence Level**: 90%
  - Geometric setup is clear and uses standard techniques
  - Trigonometric manipulation is systematic
  - Solution satisfies all constraints: $0 < \theta < 60°$
  - Verified: $\sin^2\theta + \cos^2\theta = 1$ ✓
  - Verified: $\sqrt{3}\sin\theta + \cos\theta = \frac{13}{7}$ ✓
  - Answer matches choice (B)
- **Verification Applied**: Level 5 (Thorough Verification)
  - Checked angle constraint for $\theta < 60°$
  - Verified Pythagorean identity
  - Verified original trigonometric equation
  - Confirmed answer in answer choices
  - Rejected invalid quadratic root
- **Flag for Review**: No - solution is thorough and well-verified
- **Time Spent**: Approximately 6 minutes (within target range)
  - 1 min: Understanding geometry and calculating R
  - 1.5 min: Setting up area decomposition
  - 2 min: Trigonometric manipulation to linear equation
  - 1 min: Solving for $\sin\theta$ and $\cos\theta$
  - 30 sec: Computing $\tan\theta$ and final check

Learning Points

Key Takeaways from This Problem:

1. **Rotation and Symmetry**: When a regular figure is rotated about its center, exploit rotational symmetry to decompose complex regions into simpler pieces.

2. **Area Decomposition**: Express complex areas as sums of triangular sectors from the center. Use $[\triangle] = \frac{1}{2}r^2 \sin\alpha$ for sectors.

3. **Circumradius Formula**: For equilateral triangle with side s: $R = \frac{s\sqrt{3}}{3}$. This appears frequently in AMC problems.

4. **Angle Relationships**: After rotating by θ, new angles become θ and $120° - \theta$ due to the $120°$ spacing in equilateral triangles.

5. **Trigonometric Equation Solving**: The strategy of isolating one trig function and using $\sin^2 + \cos^2 = 1$ to get a quadratic is fundamental for late-band problems.

6. **Solution Validation**: Always check which quadratic root satisfies the original constraints. Here, $\theta < 60°$ means $\sin\theta < \frac{\sqrt{3}}{2}$.

7. **Multiple Solution Paths**: This problem can be solved via:

   - Area decomposition (used above)
   - Coordinate geometry with rotation matrices
   - Direct calculation of overlap regions

   Choose the path that plays to your strengths!

Related Problems to Practice:

- Other AMC12 problems involving rotation and area
- Trigonometric identities with $\sin(120° \pm \theta)$
- Equilateral triangle properties and circumradius
- Area decomposition techniques
- Solving systems: $a\sin\theta + b\cos\theta = c$

Formulas to Remember:
Equilateral Triangle (side s):

- Area: $A = \frac{\sqrt{3}}{4}s^2$
- Circumradius: $R = \frac{s\sqrt{3}}{3}$

- Inradius: $r = \frac{s\sqrt{3}}{6} = \frac{R}{2}$
- Central angles: $120°$ between vertices

Triangle Area from Sides and Angle:

$$[\triangle] = \frac{1}{2}ab\sin C$$

where C is the angle between sides a and b.

Key Trigonometric Identity:

$$\sin(120° - \theta) = \frac{\sqrt{3}}{2}\cos\theta + \frac{1}{2}\sin\theta$$