

Status of petrale sole (*Eopsetta jordani*) along the U.S. west coast in 2019

Chantel R. Wetzel¹

¹Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East, Seattle, Washington 98112

June 2019

DRAFT SAFE

Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by NOAA Fisheries. It does not represent and should not be construed to represent any agency determination or policy.

This report may be cited as:

Wetzel, C.R. 2019. Status of petrale sole (*Eopsetta jordani*) along the U.S. west coast in 2019. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Status of petrale sole (*Eopsetta jordani*) along the U.S. west coast in 2019

Contents

Executive Summary	i
Stock	i
Landings	i
Data and Assessment	iii
Updated Data	iii
Stock Biomass	iv
Recruitment	vii
Exploitation Status	ix
Ecosystem Considerations	xii
Reference Points	xii
Management Performance	xiii
Unresolved Problems and Major Uncertainties	xiv
Decision Table	xiv
Research and Data Needs	xvii
1 Introduction	1
1.1 Basic Information	1
1.2 Life History	2
1.3 Historical and Current Fishery Information	2
1.4 Summary of Management History and Performance	4
1.5 Fisheries off Canada and Alaska	5
2 Data	5
2.1 Fishery-Independent Data	6
2.1.1 NWFSC West Coast Groundfish Bottom Trawl Survey	6
2.1.2 AFSC/NWFSC West Coast Triennial Shelf Survey	8
2.2 Fishery-Dependent Data	9

2.2.1	Commercial Fishery Landings	9
2.2.2	Discards	10
2.2.3	Foreign Landings	11
2.2.4	Historical Commercial Catch-Per-Unit Effort/Logbooks	11
2.2.5	Fishery Length and Age Data	11
2.3	Biological Data	12
2.3.1	Natural Mortality	12
2.3.2	Maturation and Fecundity	13
2.3.3	Sex Ratio	14
2.3.4	Length-Weight Relationship	14
2.3.5	Growth (Length-at-Age)	15
2.3.6	Ageing Precision and Bias	15
2.3.7	Environmental and Ecosystem Data	15
3	Assessment Model	16
3.1	History of Modeling Approaches Used for This Stock	16
3.2	General Model Specifications and Assumptions	17
3.2.1	Changes Between the 2015 Update and Current Assessment Model .	17
3.2.2	Summary of Fleets and Areas	18
3.2.3	Priors	18
3.2.4	Data Weighting	18
3.2.5	Estimated and Fixed Parameters	18
3.2.6	Key Assumptions and Structural Choices	19
3.2.7	Bridging Analysis	19
3.2.8	Convergence	19
3.3	Base Model Results	19
3.3.1	Parameter Estimates	20
3.3.2	Fits to the Data	21
3.3.3	Population Trajectory	22
3.3.4	Sensitivity Analyses	23
3.3.5	Retrospective Analysis	24
3.3.6	Historical Analysis	24
3.3.7	Likelihood Profiles	24
3.3.8	Reference Points	24

4 Harvest Projections and Decision Tables	25
5 Regional Management Considerations	25
6 Research Needs	25
7 Acknowledgments	26
8 References	27
9 Tables	31
10 Figures	69
11 Appendix A. Detailed Fit to Length Composition Data	190
12 Appendix B. Detailed Fit to Age Composition Data	213
13 Appendix C. List of Auxiliary Files Available	230

Executive Summary

executive-summary

Stock

stock

This assessment reports the status of the petrale sole (*Eopsetta jordani*) off U.S. coast of California, Oregon, and Washington using data through 2018. While petrale sole are modeled as a single stock, the spatial aspects of the coast-wide population are addressed through geographic separation of data sources/fleets where possible. There is currently no genetic evidence suggesting distinct biological stocks of petrale sole off the U.S. coast. The limited tagging data available to describe adult movement suggests that petrale sole may have some homing ability for deep water spawning sites but also have the ability to move long distances between spawning sites, inter-spawning season, as well as seasonally.

Landings

landings

While records do not exist, the earliest catches of petrale sole are reported in 1876 in California and 1884 in Oregon. In this assessment, fishery removals have been divided among 4 fleets: 1) winter North trawl, 2) summer North trawl, 3) winter South trawl, and 4) summer South trawl. Landings for the North fleet are defined as fish landed in Washington and Oregon ports. Landings for the South fleet are defined as fish landed in California ports. Recent annual catches between 1981-2018 range between 755 and 3008 mt per year and the most recent year landing are shown in Table a. Petrale sole are caught nearly exclusively by trawl fleets; non-trawl gears contribute less than 3% of the catches. Based on the 2005 assessment, annual catch limits (ACLs) were reduced to 2499 mt for 2007-2008. Following the 2009 assessment ACLs were further reduced to a low of 976 mt for 2011 and have subsequently increased to a high value of 3,136 for 2017. From the inception of the fishery through the war years, the vast majority of catches occurred between March and October (the summer fishery), when the stock is dispersed over the continental shelf. The post-World War II period witnessed a steady decline in the amount and proportion of annual catches occurring during the summer months (March-October). Conversely, petrale sole catch during the winter season (November-February), when the fishery targets spawning aggregations, has exhibited a steadily increasing trend since the 1940s. From the mid-1980s through the early 2000s, catches during the winter months were roughly equivalent to or exceeded catches throughout the remainder of the year, whereas during the past 10 years the relative catches during the winter and summer have been more variable across years (a). petrale sole are a desirable market species and discarding has historically been low.

Table a: Landings (mt) for the past 10 years for petrale sole by source.

`tab:Exec_catch`

Year	Winter (N)	Summer (N)	Winter (S)	Summer (S)	Total Landings
2009	846.71	641.75	469.66	250.38	2208.49
2010	258.09	292.34	77.60	120.95	748.98
2011	221.60	423.11	39.59	77.70	762.00
2012	406.05	477.71	124.46	107.63	1115.85
2013	509.04	1007.26	130.10	278.35	1924.74
2014	852.90	860.31	273.40	354.19	2340.80




Figure a: 'Landings of by the Northern and Southern winter and summer fleets of the US west coast.' `fig:Exec_catch1`

Data and Assessment

data-and-assessment

This an update assessment for petrale sole, which was last assessed in 2013 and updated in 2015. The update assessment was conducted using the length- and age-structured modeling software Stock Synthesis (version 3.30.13). The coastwide population was modeled allowing separate growth and mortality parameters for each sex (a two-sex model) with the fishing year beginning on November 1 and ending on October 31. The fisheries are structured seasonally based on winter (November to February) and summer (March to October) fishing seasons due to the development and growth of the wintertime fishery, which began in the 1950s. In recent decades wintertime catches have often exceed summertime catches. The fisheries modeled as the North Winter and North Summer, where the north includes both Washington and Oregon, and South Winter and South Summer encompasses California fisheries.

The model includes catch, length- and age-frequency data from the trawl fleets as well as standardized winter fishery catch-per-unit-effort (CPUE) indices. Biological data are derived from both port and on-board observer sampling programs. The National Marine Fisheries Service (NMFS) AFSC/NWFSC West Coast Triennial Shelf Survey early (1980, 1983, 1986, 1989, 1992) and late period (1995, 1998, 2001, and 2004) and the NWFSC West Coast Groundfish Bottom Trawl Survey (2003-2018) relative biomass indices and biological sampling provide fishery independent information on relative trend and demographics of the petrale sole stock.

Updated Data

updated-data

The base stock assessment model structure is consistent with the 2013 assessment and the 2015 update, except as noted here. Modifications from the previous assessment model include:

1. Model fitting using latest version of Stock Synthesis (SS v.3.30.13).
2. Added commercial fishery catch data (2015-2018).
3. Updated historical composition data from the commercial fishery (length and age data) and new data (2015 - 2018) expanded to trip and catch based on current best practices.
4. Updated discard rate, average weight, and discard length composition data (2014-2017).
5. NWFSC West Coast Groundfish Bottom Trawl Survey index of abundance was calculated using VAST.
6. Updated NWFSC West Coast Groundfish Bottom Trawl Survey length and age data (2015-2018) with all years data expanded to the tow and strata.
7. AFSC/NWFSC West Coast Triennial Shelf Survey early and late index of abundance were calculated using VAST.

8. Model tuning to re-weight data.
9. Length-weight relationship parameters estimated outside of the stock assessment model from the NWFSC West Coast Groundfish Bottom Trawl Survey data up to 2018 and input as fixed values.
10. Update the natural mortality prior for female and male fish.

Stock Biomass

stock-biomass

Petrale sole were lightly exploited during the early 1900s, but by the 1950s the fishery was well developed and showing clear signs of depletion and declines in catches and biomass (Figures [a](#) and [b](#)). The rate of decline in spawning biomass accelerated through the 1930s-1970s reaching minimums generally around or below 10% of the unexploited levels during the 1980s through the early 2000s (Figure [c](#)). The petrale sole spawning stock biomass is estimated to have increased in recent year due to reduced catches during rebuilding and in response to above average recruitment in 2006, 2007, and 2008. The 2019 estimated spawning biomass relative to unfished equilibrium spawning biomass is above the target of 25% of unfished spawning biomass at 32.3% (\sim 95% asymptotic interval: \pm 21.9%-42.6%).

Table b: Recent trend in estimated spawning biomass (mt) and estimated relative spawning biomass (depletion).

Year	Spawning Biomass (mt)	~ 95% Confidence Interval	Estimated Relative Spawning Biomass	tab:SpawningDeplete_mod1	
				~ 95%	Confidence Interval
2010	3494	2845 - 4143	0.114	0.077	- 0.152
2011	4414	3606 - 5222	0.144	0.098	- 0.191
2012	5904	4858 - 6950	0.193	0.132	- 0.255
2013	7751	6410 - 9091	0.254	0.174	- 0.333
2014	9284	7682 - 10886	0.304	0.209	- 0.398
2015	10202	8441 - 11962	0.334	0.231	- 0.436
2016	10439	8593 - 12284	0.342	0.237	- 0.446
2017	10531	8606 - 12457	0.345	0.240	- 0.449
2018	10213	8186 - 12240	0.334	0.231	- 0.437
2019	9867	7682 - 12052	0.323	0.219	- 0.426




Figure b: Estimated time-series of spawning biomass trajectory (circles and line; median; light broken lines; 95% credibility intervals) for the base assessment model. fig:Spawnbio_all

Spawning depletion with ~95% asymptotic intervals




Figure c: Estimated time-series of relative spawning biomass (depletion) (circles and line: median; light broken lines: 95% credibility intervals) for the base assessment model. | [fig:RelDeplete_all](#)

Recruitment

recruitment

Annual recruitment was treated as stochastic, and estimated as annual deviations from log-mean recruitment where mean recruitment is the fitted Beverton-Holt stock recruitment curve. The time-series of estimated recruitments shows a relationship with the decline in spawning biomass, punctuated by larger recruitments (Figure d) in 2006, 2007, and 2008. The five largest estimated recruitment estimated with the model (in ascending order) occurred in 2006, 2007, 1998, 1966, and 2008. The four lowest recruitments estimated within the model (in ascending order) occurred in 1986, 1992, 1973, and 1987.

Table c: Recent estimated trend in recruitment and estimated recruitment deviations determined from the base model. The recruitment deviations for 2018 and 2019 were fixed at zero within the model.

Year	Estimated Recruitment	~ 95% Confidence Interval	Estimated Recruitment Devs.	~ 95% Confidence Interval
2010	11740	7281 - 18929	-0.083	-0.411 - 0.244
2011	13434	8205 - 21994	-0.014	-0.358 - 0.330
2012	19781	12473 - 31372	0.306	0.002 - 0.609
2013	12763	7331 - 22221	-0.183	-0.631 - 0.265
2014	13826	7883 - 24251	-0.130	-0.586 - 0.325
2015	13751	7464 - 25334	-0.149	-0.664 - 0.366
2016	13924	6927 - 27992	-0.160	-0.771 - 0.451
2017	15071	6851 - 33154	-0.104	-0.837 - 0.630
2018	17012	7437 - 38914	0.000	-0.784 - 0.784
2019	16935	7409 - 38710	0.000	-0.784 - 0.784

Age-0 recruits (1,000s) with ~95% asymptotic intervals




Figure d: Time-series of estimated petrale sole recruitments for the base model with 95% confidence or credibility intervals. [fig:Recruits_all](#)

Exploitation Status

`exploitation-status`

The relative spawning biomass of petrale sole was estimated to have dropped below the management target (25%) for the first time in 1976. The stock continued to decline and first fell below the minimum stock size threshold level of 12.5% in 1982. The relative spawning biomass remained around the threshold stock size until approximately 2010, with the stock reaching its lowest relative spawning biomass level in 1993 at 5.8%. In 2009 petrale sole was formally declared overfished. Fishing mortality rates sharply declined during the rebuilding period relative to previous year rates which exceeded the target (Figure e). After reduced harvests, the 2015 update stock assessment estimated the stock to have rebuilt to the management target (25%) in 2014. This update estimates that the relative spawning biomass exceed 25% in 2013 with harvest rates in the most recent years remaining just under of the target rate (Figure e).

Table d: Recent trend in spawning potential ratio 1-SPR and summary exploitation rate for age 3+ biomass for petrale sole.

Year	1-SPR	~ 95% Confidence Interval	Exploitation Rate	tab:SPR_Exploit_mod1 ~ 95% Confidence Interval
2009	0.822	0.757 - 0.888	0.271	0.223 - 0.318
2010	0.625	0.521 - 0.728	0.091	0.072 - 0.111
2011	0.552	0.446 - 0.658	0.062	0.049 - 0.075
2012	0.568	0.466 - 0.671	0.074	0.060 - 0.089
2013	0.632	0.535 - 0.730	0.111	0.090 - 0.131
2014	0.633	0.537 - 0.729	0.126	0.104 - 0.149
2015	0.644	0.549 - 0.738	0.138	0.113 - 0.163
2016	0.618	0.521 - 0.714	0.129	0.105 - 0.153
2017	0.654	0.561 - 0.747	0.155	0.126 - 0.185
2018	0.649	0.555 - 0.743	0.152	0.121 - 0.183




Figure e: Estimated relative spawning potential ratio 1-SPR for the base model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the SPR30% harvest rate. The last year in the time-series is 2018. | [fig:SPR_all](#)




Figure f: Phase plot of estimated 1-SPR(%) vs. relative spawning biomass (B/B_{target}) for the base case model. The red circle indicates 2018 estimated status and exploitation for petrale sole. |
fig:Phase_all

Ecosystem Considerations

ecosystem-considerations

Ecosystem factors have not been explicitly modeled in this assessment, but there are several aspects of the California current ecosystem that may impact petrale sole population dynamics and warrant further research. Castillo (1992) and Castillo et al. (1995) suggest that density-independent survival of early life stages is low and show that offshore Ekman transportation of eggs and larvae may be an important source of variation in year-class strength in the Columbia INPFC area. The effects of the Pacific Decadal Oscillation on California current temperature and productivity (Mantua et al. 1997) may also contribute to non-stationary recruitment dynamics for petrale sole. The prevalence of a strong late 1990s year-class for many West Coast groundfish species suggests that environmentally driven recruitment variation may be correlated among species with relatively diverse life history strategies. Although current research efforts along these lines are limited, a more explicit exploration of ecosystem processes may be possible in future petrale sole stock assessments if resources are available for such investigations.

Reference Points

reference-points

This stock assessment estimates that the spawning biomass of petrale sole is above the management target. Due to reduced landings and the large 2008 year-class, an increasing trend in spawning biomass was estimated in the base model. The estimated depletion in 2019 is 32.3% ($\sim 95\%$ asymptotic interval: $\pm 21.9\%-42.6\%$), corresponding to an spawning biomass of 9,867 mt ($\sim 95\%$ asymptotic interval: 7,682-12,052 mt). Unfished age 3+ biomass was estimated to be 49,439.6 mt in the base model. The target spawning biomass based on the biomass target ($SB_{25\%}$) is 7,638.7 mt, with an equilibrium catch of 2,830.3 mt. Equilibrium yield at the proxy F_{MSY} harvest rate corresponding to $SPR_{30\%}$ is 2,819.8 mt. Estimated MSY catch is at a 2,835.9 spawning biomass of 7,005.9 mt (22.9% relative spawning biomass).

Table e: Summary of reference points and management quantities for the base case.

Quantity	Estimate	tab:Ref_pts_mod1	
		~2.5%	~97.5%
		Confidence Interval	Confidence Interval
Unfished spawning biomass (mt)	30554.7	24634.6	36474.8
Unfished age 3+ biomass (mt)	49439.6	41597	57282.2
Unfished recruitment (R0, thousands)	18626.7	11147.4	26106
Spawning biomass(2019 mt)	9867.3	7682.4	12052.2
Relative spawning biomass (depletion) (2019)	0.323	0.219	0.426
Reference points based on SB_{25%}			
Proxy spawning biomass ($B_{25\%}$)	7638.7	6158.7	9118.7
SPR resulting in $B_{25\%}$ ($SPR_{B25\%}$)	0.286	0.258	0.313
Exploitation rate resulting in $B_{25\%}$	0.182	0.163	0.2
Yield with $SPR_{B25\%}$ at $B_{25\%}$ (mt)	2830.3	2624.2	3036.4
Reference points based on SPR proxy for MSY			
Spawning biomass	8096.3	6199.3	9993.3
$SPR_{30\%}$			
Exploitation rate corresponding to $SPR_{30\%}$	0.173	0.145	0.2
Yield with $SPR_{30\%}$ at SB_{SPR} (mt)	2819.8	2590.1	3049.5
Reference points based on estimated MSY values			
Spawning biomass at MSY (SB_{MSY})	7005.9	5242.2	8769.6
SPR_{MSY}	0.266	0.201	0.331
Exploitation rate at MSY	0.195	0.164	0.225
MSY (mt)	2835.9	2641.9	3029.9

Management Performance

management-performance

The 2009 stock assessment estimated petrale sole to be at 11.6% of unfished spawning stock biomass in 2010. Based on the 2009 stock assessment, the 2010 coast-wide ACL was reduced to 1,200 mt to reflect the overfished status of the stock and the 2011 coast-wide overfishing limit (OFL) and ACL were set at 1,021 mt and 976 mt, respectively (Table f).

Recent coast-wide annual landings have not exceeded the ACL. The 2009, 2011, and 2013 full assessments estimated that petrale sole have been below the management target since the 1960s and below the overfished threshold between the early 1980s and the early 2000s with fishing mortality rates in excess of the current F-target for flatfish of SPR30%. The 2015 update assessment estimated that the stock had recovered with the relative spawning biomass exceeding the management target.

Table f: Recent trend in total catch and landings (mt) relative to the management guidelines. Estimated total catch reflect the landings plus the model estimated discarded biomass based on discard rate data.

Year	OFL (mt; ABC prior to 2011)	ACL (mt; OY prior to 2011)	Total Landings (mt)	Estimated Total Catch (mt)
2009	2811	2433	2209	2329
2010	2751	1200	755	867
2011	1021	976	768	791
2012	1275	1160	1135	1159
2013	2711	2592	1936	1973
2014	2774	2652	2373	2398
2015	3073	2816	2686	2727
2016	3208	2910	2506	2543
2017	3208	3136	3008	3050
2018	3152	3013	2840	2882

Unresolved Problems and Major Uncertainties

unresolved-problems-and-major-uncertainties

Parameter uncertainty is explicitly captured in the asymptotic confidence intervals reported throughout this assessment for key parameters and management quantities. These intervals reflect the uncertainty in the model fit to the data sources included in the assessment, but do not include uncertainty associated with alternative model configurations, weighting of data sources (a combination of input sample sizes and relative weighting of likelihood components), or fixed parameters.

There are a number of major uncertainties regarding model parameters that have been explored via sensitivity analysis. The most notable explorations involved the sensitivity of model estimates to:

1. Value of female natural mortality.
2. Sex ratio between female and male petrale sole.
3. Changes in estimated based on alternative data weighting approaches.

Additionally, to date a reconstructed historical Washington catch history has not been included in the petrale sole stock assessment. Washington state is currently undergoing efforts to determine historical catches.

Decision Table

decision-table

Model uncertainty has been described by the estimated uncertainty within the base model and by the sensitivities to different model structure.

Table g: Projections of potential OFL (mt) and ABC (mt) and the estimated spawning biomass and relative depletion based on ABC removals. The 2019 and 2020 removals are set at the harvest limits currently set by management of XXX mt per year.

Year	OFL	ABC	Spawning Biomass (mt)	Relative Depletion
2019	3436	3301	9867	0.323
2020	3189	3065	9191	0.301
2021	2987	2870	8630	0.282
2022	2843	2731	8235	0.270
2023	2764	2655	8035	0.263
2024	2742	2633	8009	0.262
2025	2757	2648	8087	0.265
2026	2790	2680	8201	0.268
2027	2824	2713	8304	0.272
2028	2851	2738	8379	0.274
2029	2869	2756	8427	0.276
2030	2880	2766	8455	0.277

Table h: Decision table summary of 10-year projections beginning in 2021 for alternate states of nature based on an axis of uncertainty for the base model. The removals in 2019 and 2020 were set at the defined management specification of XXX mt for each year assuming full attainment. The range of natural mortality values corresponded to the 12.5 and 87.5th quantile from the uncertainty around final spawning biomass. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. The SPR30 catch stream is based on the equilibrium yield applying the SPR30 harvest rate.

		States of nature					
		Low State		Base		High State	
	Year	Catch	Spawning Biomass	Depletion	Spawning Biomass	Depletion	Spawning Biomass
ABC	2021						
	2022						
	2023						
	2024						
	2025						
	2026						
	2027						
	2028						
	2029						
	2030						
SPR target = 0.34	2021						
	2022						
	2023						
	2024						
	2025						
	2026						
	2027						
	2028						
	2029						
	2030						

Research and Data Needs

research-and-data-needs

Progress on a number of research topics and data issues would substantially improve the ability of this assessment to reliably and precisely model petrale sole population dynamics in the future:

1. In the past many assessments have derived historical catches independently. The states of California and Oregon have completed comprehensive historical catch reconstructions. At the time of this assessment, a comprehensive historical catch reconstruction is not available for Washington. Completion of a Washington catch reconstruction would provide the best possible estimated catch series that accounts for all the catch and better resolves historical catch uncertainty for flatfish as a group.
2. Due to limited data, new studies on the maturity relationships for petrale sole would be beneficial.
3. Where possible, historical otolith samples aged using a combination of surface and break-and-burn methods should be re-aged using the break-and-burn method. Early surface read otoliths should also be re-aged using the break-and-burn method. Historical otoliths aged with a standard method will allow the further evaluation of the potential impacts of consistent under ageing using surface methods, changes in selectivity during early periods of time without any composition information, and potential changes in growth.
4. Studies on stock structure and movement of petrale sole, particularly with regard to the winter-summer spawning migration of petrale sole and the likely trans-boundary movement of petrale sole between U.S. and Canadian waters seasonally.
5. The extent of spatial variability on productivity processes such as growth, recruitment, and maturity is currently unknown and would benefit from further research.

Table i: Base model results summary.

Quantity	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
OFL (mt)	2751	1021	1275	2711	2774	3073	3208	3208	3152	1
ACL (mt)	1200	976	1160	2592	2652	2816	2910	3136	3013	1
Landings (mt)	755	768	1135	1936	2373	2686	2506	3008	2840	
Total Est. Catch (mt)	867	791	1159	1973	2398	2727	2543	3050	2882	
1-SPR	0.625	0.552	0.568	0.632	0.633	0.644	0.618	0.654	0.649	
Exploitation rate	0.091	0.062	0.074	0.111	0.126	0.138	0.129	0.155	0.152	
Age 3+ biomass (mt)	9487.05	12726.80	15602.80	17806.70	18073.00	19748.50	19715.50	19637.00	18918.70	18265.30
Spawning Biomass	3494	4414	5904	7751	9284	10202	10439	10531	10213	9867
95% CI	2845 - 4143	3606 - 5222	4858 - 6950	6410 - 9091	7682 - 10886	8441 - 11962	8593 - 12284	8606 - 12457	8186 - 12240	7682 - 12052
Relative Depletion	0.114	0.144	0.193	0.254	0.304	0.334	0.342	0.345	0.334	0.323
95% CI	0.077 - 0.152	0.098 - 0.191	0.132 - 0.255	0.174 - 0.333	0.209 - 0.398	0.231 - 0.436	0.237 - 0.446	0.240 - 0.449	0.231 - 0.437	0.219 - 0.426
Recruits	11740	13434	19781	12763	13826	13751	13924	15071	17012	16935
95% CI	7281 - 18929	8205 - 21994	12473 - 31372	7331 - 22221	7883 - 24251	7464 - 25334	6927 - 27992	6851 - 33154	7437 - 38914	7409 - 38710




Figure g: Equilibrium yield curve for the base case model. Values are based on the 2018 fishery selectivity and with steepness fixed at 0.84. [fig:Yield_all](#)

1 Introduction

introduction

This updated assessment does not attempt to reiterate all background information for petrale sole presented in the 2013 assessment document. Instead, only a few key assumptions are restated, along with a detailed description of changes made during the course of the update. Those interested in a more complete description of petrale sole life-history and the details of previous assessments should refer to the 2013 assessment (Haltuch et al. 2013b).

1.1 Basic Information

basic-information

Petrale sole (*Eopsetta jordani*) is a right-eyed flounder in the family Pleuronectidae ranging from the western Gulf of Alaska to the Coronado Islands, northern Baja California (Kramer et al. 1995, Love et al. 2005) with a preference for soft substrates at depths ranging from 0-550 m (Love et al. 2005). Common names include brill, California sole, Jordan's flounder, cape sole, round nose sole, English sole, soglia, petorau, nameta, and tsubame garei (Smith 1937, Gates and Frey 1974, Eschmeyer and Herald 1983, Love 1996). In northern and central California petrale sole are dominant on the middle and outer continental shelf. PacFIN fishery logbook data show that adults are caught in depths from 18 to 1,280 m off the U.S. West Coast with a majority of the catches of petrale sole being taken between 70-220 m during March through October, and between 290-440 m during November through February.

Past assessments completed by Demory (1984), Turnock et al. (1993), and Sampson and Lee (1999) considered petrale sole in the Columbia and U.S.-Vancouver INPFC areas a single stock. Sampson and Lee (1999) assumed that petrale sole in the Eureka and Monterey INPFC areas represented two additional distinct stocks. The 2005 petrale sole assessment assumed two stocks, northern (U.S.-Vancouver and Columbia INPFC areas) and southern (Eureka, Monterey and Conception INPFC areas), to maintain continuity with previous assessments. Three stocks (West Coast Vancouver Island, Queen Charlotte Sound, and Heceta Strait) are considered for petrale sole in the waters off British Columbia, Canada (Starr and Fargo 2004). The 2009, 2011, 2013, and 2015 assessments integrate the previously separate north-south assessments to provide a coast-wide status evaluation. The decision to conduct a single-area assessment is based on strong evidence of a mixed stock from tagging studies, a lack of genetic studies on stock structure, and a lack of evidence for differences in growth between the 2005 northern and southern assessment areas and from examination of the fishery size-at-age data, as well as confounding differences in data collection between Washington, Oregon, and California. This 2019 update assessment provides a coast-wide status evaluation for petrale sole using data through 2018.

Fishing fleets are separated both geographically and seasonally to account for spatial and seasonal patterns in catch given the coast-wide assessment area. The petrale sole fisheries possess a distinct seasonality, with catches peaking during the winter months, so the fisheries are divided into winter (November-February) and summer (March-October) fisheries. Note

that the “fishing year” for this assessment (November 1 to October 31) differs from the standard calendar year. The U.S.-Canadian border is the northern boundary for the assessed stock, although the basis for this choice is due to political and current management needs rather than the population dynamics. Given the lack of clear information regarding the status of distinct biological populations, this assessment treats the U.S. Petrale sole resource from the Mexican border to the Canadian border as a single coast-wide stock.

1.2 Life History

life-history

Petrale sole spawn during the winter at several discrete deep water sites (270-460 m) off the U.S. West Coast, from November to April, with peak spawning taking place from December to February (Harry 1959, Best 1960, Gregory and Jow 1976, Castillo et al. 1993, Reilly et al. 1994, Love 1996). Females spawn once each year and fecundity varies with fish size, with one large female laying as many as 1.5 million eggs (Porter 1964). Petrale sole eggs are planktonic, ranging in size from 1.2 to 1.3 mm, and are found in deep water habitats at water temperatures of 4-10 degrees C and salinities of 25-30 ppt (Best 1960, Ketchen and Forrester 1966, Alderdice and Forrest 1971, Gregory and Jow 1976). The duration of the egg stage can range from approximately 6 to 14 days (Alderdice and Forrest 1971, Love 1996). The most favorable conditions for egg incubation and larval growth are 6-7 degrees C and 27.5-29.5 ppt (Ketchen and Forrester 1966, Alderdice and Forrest 1971, Castillo 1995).

Adult petrale sole achieve a maximum size of around 50 cm and 63 cm for males and females, respectively (Best 1963, Pedersen 1975). The maximum length reported for petrale sole is 70 cm (Eschmeyer and Herald 1983, Love et al. 2005) while the maximum observed break-and-burn age is 31 years (Haltuch et al. 2013b).

1.3 Historical and Current Fishery Information

historical-and-current-fishery-information

Petrale sole have been caught in the flatfish fishery off the U.S. Pacific coast since the late 19th century. The fishery first developed off of California where, prior to 1876, fishing in San Francisco Bay was by hand or set lines and beach seining (Scofield 1948). By 1880 two San Francisco based trawler companies were running a total of six boats, extending the fishing grounds beyond the Golden Gate Bridge northward to Point Reyes (Scofield 1948). Steam trawlers entered the fishery during 1888 and 1889, and four steam tugs based out of San Francisco were sufficient to flood market with flatfish (Scofield 1948). By 1915 San Francisco and Santa Cruz trawlers were operating at depths of about 45-100 m with catches averaging 10,000 lbs per tow or 3,000 lbs per hour (Scofield 1948). Flatfish comprised approximately 90% of the catch with 20-25% being discarded as unmarketable (Scofield 1948). During 1915 laws were enacted that prohibited dragging in California waters and making it illegal to possess a trawl net from Santa Barbara County southward (Scofield 1948). By 1934 twenty 56-72 foot diesel engine trawlers operated out of San Francisco fishing between about 55

and 185 m (Scofield 1948). From 1944-1947 the number of California trawlers fluctuated between 16 and 46 boats (Scofield 1948). Although the flatfish fishery in California was well developed by the 1950s and 1960s, catch statistics were not reported until 1970 (Heimann and Carlisle 1970). In this early California report petrale sole landings during 1916 to 1930 were not separated from the total flatfish landings.

The earliest trawl fishing off Oregon began during 1884-1885, and the fishery was solidly established by 1937, with the fishery increasing rapidly during WWII (Harry and Morgan, 1961). Initially trawlers stayed close to the fishing grounds adjacent to Newport and Astoria, operating at about 35-90 m between Stonewall Bank and Depoe Bay. Fishing operations gradually extended into deep water. For example, Newport-based trawlers were commonly fishing at about 185 m in 1949, at about 185-365 m by 1952, and at about 550 m by 1953.

Alverson and Chatwin (1957) describe the history of the petrale sole fishery off of Washington and British Columbia with fishing grounds ranging from Cape Flattery to Destruction Island. Petrale sole catches off of Washington were small until the late 1930s with the fishery extending to about 365 m following the development of deep water rockfish fisheries during the 1950s.

By the 1950s the petrale sole fishery was showing signs of depletion with reports suggesting that petrale sole abundance had declined by at least 50% from 1942 to 1947 (Harry 1956). Sampson and Lee (1999) reported that three fishery regulations were implemented during 1957-67: 1) a winter closure off Oregon, Washington and British Columbia, 2) a 3,000 lb per trip limit, and 3) no more than two trips per month during 1957. With the 1977 enactment of the Magnuson Fishery Conservation and Management Act (MFCMA) the large foreign-dominated fishery that had developed since the late 1960s was replaced by the domestic fishery that continues today. Petrale sole are harvested almost exclusively by bottom trawls in the U.S. West Coast groundfish fishery. Recent petrale sole catches exhibit marked seasonal variation, with substantial portions of the annual harvest taken from the spawning grounds during December and January. Evidence suggests that the winter fishery on the deep water spawning grounds developed sporadically during the 1950s and 1960s as fishers discovered new locations (e.g., Alverson and Chatwin (1957); Ketchen and Forrester (1966)). Both historical and current petrale sole fisheries have primarily relied upon trawl fleets. Fishery removals were divided among 4 fleets: 1) winter North trawl, 2) summer North trawl, 3) winter South trawl, and 4) summer South trawl. Landings for the North fleet are defined as fish landed in Washington and Oregon ports. Landings for the South fleet are defined as fish landed in California ports.

Historical landings reconstructions show peak catches from the summer fishery occurred during the 1940s and 1950s and subsequently declined, during which time the fleet moved to fishing in deeper waters during the winter. After the period of peak landings during the 1940s and 1950s, total landings were somewhat stable until about the late 1970s, and then generally declined until the mid-2000s. (Table 1, Figure 1). During 2009 the fishery was declared overfished and during 2010 management restrictions limited the catch to 755 mt (Table 1, Figure 1). Recent years overfishing limit (OFL), annual catch limit (ACL), landings, and estimated total dead are shown in Table 2.

1.4 Summary of Management History and Performance

summary-of-management-history-and-performance

Beginning in 1983 the Pacific Fishery Management Council (PFMC) established coast-wide annual catch limits (ACLs) for the annual harvests of petrale sole in the waters off the U.S. West Coast. The first assessment of West Coast petrale sole occurred in 1984 (Demory 1984). Based on the 1999 assessment a coast-wide ACL of 2,762 mt was specified and remained unchanged between 2001 and 2006.

The 2005 assessment of petrale sole stock assessment split the stock into two areas, the northern area that included U.S.-Vancouver and Columbia INPFC areas and the southern area that included the Eureka, Monterey and Conception INPFC areas (Lai et al. 2005). While petrale sole stock structure is not well understood, CPUE and geographical differences between states were used to support the use of two separate assessment areas. In 2005 petrale sole were estimated to be at 34 and 29% of unfished spawning stock biomass in the northern and southern areas, respectively. In spite of different models and data, the biomass trends were qualitatively similar in both areas, providing support for a coast-wide stock. This assessment estimated that petrale sole had historically been below the Pacific Council's minimum stock size threshold of 25% of unfished biomass from the mid-1970s until just prior to the completion of the assessment, with estimated harvest rates in excess of the target fishing mortality rate implemented for petrale sole at that time (F40%). However, the 2005 stock assessment determined that the stock was in the precautionary zone and was not overfished (i.e., the spawning stock biomass was not below 25% of the unfished spawning stock biomass). Based on the 2005 stock assessment results, ACLs were set at 3,025 mt and 2,919 mt for 2007 and 2008, respectively, with an ACT of 2,499 mt for both years.

In comparison to the 1999 assessment of petrale sole, the 2005 assessment represented a significant change in the perception of petrale sole stock status. The stock assessment conducted in 1999 (Washington-Oregon only) estimated the spawning stock biomass in 1998 at 39% of unfished stock biomass. Although the estimates of 1998 spawning-stock biomass were little changed between the 1999 and 2005 (Northern area) assessments, the estimated depletion in the 2005 assessment was much lower. The change in status between the 1999 and 2005 analyses was due to the introduction of a reconstructed catch history in 2005, which spanned the entire period of removals. The 1999 stock assessment used a catch history that started in 1977, after the bulk of the removals from the fishery had already taken place. Thus the 1999 stock assessment produced a more optimistic view of the petrale stock's level of depletion. The stock's estimated decline in status between the 2005 and 2009 assessments was driven primarily by a significant decline in the trawl-survey index over that period. The 2011 assessment concluded that the stock status continued to be below the target of 25% of unfished biomass.

The 2009 coast-wide stock assessment estimated that the petrale sole stock had declined from its 2005 high to 11.6% of the unfished spawning stock biomass (Haltuch and Hicks 2009). The petrale sole was declared overfished based on newly adopted management targets (e.g., target spawning biomass for flatfish stocks defined as 25% and overfished threshold of 12.5%

of unfished spawning stock biomass) resulting in a rebuilding plan and catch restrictions for petrale sole. The stock was declared rebuilt based on the results of the 2015 update stock assessment which estimated the coastwide biomass at 30.7% of unfished spawning stock output with ACLs of 3,136 and 3,013 in 2017 and 2018 respectively (Stawitz et al. 2015).

For additional information on changes in the petrale sole fishery please see the 2013 stock assessment (Haltuch et al. 2013b).

1.5 Fisheries off Canada and Alaska

[fisheries-off-canada-and-alaska](#)

The Canadian fishery developed rapidly during the late 1940s to mid-1950s following the discovery of petrale sole spawning aggregations off the West Coast of Vancouver Island (Anon 2001). Annual landings of petrale sole in British Columbia peaked at 4,800 mt in 1948 but declined significantly after the mid-1960s (Anon 2001). By the 1970s, analysis conducted by Pederson (1975) suggested that petrale sole abundance was low and abundance remained low into the 1990s. In the early 1990s vessel trip quotas were established to try to halt the decline in petrale sole abundance (Anon 2001). Winter quarter landings of petrale sole were limited to 44,000 lb per trip during 1985-91; to 10,000 lb per trip during 1991-95; and to 2,000 lb per trip in 1996. Biological data collected during 1980-1996 showed a prolonged decline in the proportion of young fish entering the population (Anon 2001). Therefore, no directed fishing for petrale sole has been permitted in Canada since 1996 due to a continuing decline in long term abundance (Fargo 1997, Anon 2001). As of 2005 petrale sole off of British Columbia were treated as three “stocks” and were still considered to be at low levels. The recent assessments for the Canadian stocks have been based on catch histories and limited biological data.

In Alaska petrale sole are not targeted in the Bering Sea/Aleutian Island fisheries and are managed as a minor species in the “Other Flatfish” stock complex.

2 Data

[data](#)

Data used in the petrale sole assessment are summarized in Figure 2. The data that were added or reprocessed for this assessment are:

1. Commercial catches (2015-2018 added);
2. Commercial length and age data (all years reprocessed, 2015-2018 added);
3. Observed discard rates, average weights, and lengths (2002-2017 reprocessed, 2014-2017 added);

4. AFSC/NWFSC West Coast Triennial Shelf Survey early and late indices of abundance and length composition data (1980-2004 reprocessed); and
5. NWFSC West Coast Groundfish Bottom Trawl Survey index of abundance, length and age composition data (2003-2018 reprocessed, 2015-2018 added).

A description of each data source is provided below.

2.1 Fishery-Independent Data

`fishery-independent-data`

2.1.1 NWFSC West Coast Groundfish Bottom Trawl Survey

`nwfsc-west-coast-groundfish-bottom-trawl-survey`

Three sources of information are produced by this survey: an index of relative abundance, length-frequency distributions, and age-frequency distributions. Only years in which the NWFSC West Coast Groundfish Bottom Trawl Survey included the continental shelf (55-183 m) are considered (2003-2018), since the highest percent of positive survey tows with petrale sole are found on the continental shelf.

The NWFSC West Coast Groundfish Bottom Trawl Survey is based on a random-grid design; covering the coastal waters from a depth of 55 m to 1,280 m (Bradburn et al. 2011). This design uses four industry chartered vessels per year, assigned to a roughly equal number of randomly selected grid cells and divided into two ‘passes’ of the coast that are executed from north to south. Two vessels fish during each pass, which are conducted from late May to early October each year. This design therefore incorporates both vessel-to-vessel differences in catchability as well as variance associated with selecting a relatively small number (~700) of possible cells from a very large set of possible cells spread from the Mexican to the Canadian border.

The NWFSC West Coast Groundfish Bottom Trawl Survey commonly encounters petrale sole along the U.S West Coast, except south of Point Conception (Figure 3). The catch-per-unit-effort estimated from the survey is roughly constant north of 38° (Figure 4). The survey does fish shallower than 54 m and no petrale sole were caught deeper than 550 m. Figure 5 shows that the postie tows catch rate by depth peaks between 100-200 meters and declines as depth increases.

The data from the NWFSC West Coast Groundfish Bottom Trawl Survey was analyzed using a spatio-temporal delta model implemented as an R package, VAST (Thorson and Barnett 2017, Thorson 2019), which is publicly available online (<https://github.com/James-Thorson/VAST>). Spatial and spatio-temporal variation is specifically included in both encounter probability and positive catch rates, a logit-link for encounter probability and a log-link for positive catch rates. Vessel-year effects were included for each unique combination of vessel and year in the data to account for the random selection of commercial vessels used during

sampling (Helser et al. 2004, Thorson and Ward 2013). Spatial variation was approximated using 250 knots, and the model used the bias-correction algorithm (Thorson and Kristensen 2016) in Template Model Builder (Kristensen et al. 2016). Further details regarding model structure are available in the user manual ([https://github.com/James-Thorson/VAST/blob/master/examples/VAST user manual.pdf](https://github.com/James-Thorson/VAST/blob/master/examples/VAST%20user%20manual.pdf)). The stratification is provided in Table 3.

The gamma distribution with random strata-year and vessel effects was chosen as the final model. The Q-Q plot does not show any departures from the assumed distribution (Figure 6). The Pearson residuals for the encounter and catch rates for gamma distribution model are shown in Figures 7 and 8.

The estimated index of abundance is shown in Table 4. For contrast, the 2015 model estimated, the 2019 design based, and the 2019 VAST indices are shown in Figure 9 and the spatial density by year estimated by VAST are shown in Figure 10. The index for the NWFSC West Coast Groundfish Bottom Trawl Survey shows an increase in the population between 2009 and 2014 and roughly stable through 2017, and decrease in the most recent year.

Length bins from 12 to 62 cm in 2 cm increments were used to summarize the length frequency of the survey catches in each year. Table 5 shows the number of lengths taken by the survey. The first bin includes all observations less than 14 cm and the last bin includes all fish larger than 62 cm. The length frequency distributions for the NWFSC West Coast Groundfish Bottom Trawl Survey from 2003-2018 generally show a strong cohort growing through 2005 and smaller fish entering the population beginning in 2007 rows with a large 2014 cohort entering the populations (Figure 11).

Age distributions included bins from age 1 to age 17, with the last bin including all fish of greater age. Table 6 shows the number of ages taken by the survey. The marginal NWFSC West Coast Groundfish Bottom Trawl Survey age-compositions, which allow for easier viewing of strong cohorts, show the strong 1998 cohort ageing from 2003 to 2007, with younger fish appearing between 2008-2014 (Figure 12). The exception to this is the female composition in 2005, where only one female fish was aged from the tow with the largest catch rate. The expansion of numbers to tow can greatly affect the marginal age distribution, but does not have as much effect on the conditional age-at-length data.

The input sample sizes for length and marginal age-composition data for all fishery-independent surveys were calculated based on the approach used in the 2013 full and 2015 update assessment as:

$$N = (0.138 * (\sum^N fish_y / \sum^N tows_y) + 1) * \sum^N tows_y$$

where fish is the number of petrale sole by year y and the total number of tows by year. The effective sample size of conditional-age-at-length data was set at the number of fish at each length by sex and by year. The conditional-age-at-length data were not expanded and were binned by according to length, age, sex, and year.

2.1.2 AFSC/NWFSC West Coast Triennial Shelf Survey

`afscnwfsc-west-coast-triennial-shelf-survey`

The AFSC/NWFSC West Coast Triennial Shelf Survey (referred to as the Triennial Survey for short) was first conducted by the AFSC in 1977 and spanned the time-frame from 1977-2004. The survey's design and sampling methods are most recently described in Weinberg et al. (2002). Its basic design was a series of equally-spaced transects from which searches for tows in a specific depth range were initiated. The survey design has changed slightly over the period of time. In general, all of the surveys were conducted in the mid-summer through early fall: the 1977 survey was conducted from early July through late September; the surveys from 1980 through 1989 ran from mid-July to late September; the 1992 survey spanned from mid-July through early October; the 1995 survey was conducted from early June to late August; the 1998 survey ran from early June through early August; and the 2001 and 2004 surveys were conducted in May-July.

Haul depths ranged from 91-457 m during the 1977 survey with no hauls shallower than 91 m. The surveys in 1980, 1983, and 1986 covered the West Coast south to 36.8° N latitude and a depth range of 55-366 m. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5° N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55-500 m and surveyed south to 34.5° N. In the final year of the Triennial Survey series, 2004, the NWFSC's Fishery Resource and Monitoring division (FRAM) conducted the survey and followed very similar protocols as the Alaska Fisheries Science Center (AFSC).

Due to changes in survey timing, the Triennial Survey data have been split into independent early (1980-1992) and late (1995-2004) survey time series. The splitting of this time series was investigated during the 2009 STAR panel due to the changes in survey timing and the expected change in petrale sole catchability because of the stock's seasonal onshore-offshore migrations (Cook et al. 2009). For these reasons, as well as because the split improved fits to the split time series and made small changes to the estimation of the selectivity curves, the 2009 STAR panel supported the split.

The Triennial Survey commonly encounters petrale sole along the U.S West Coast (Figure 13). The catch-per-unit-effort estimated from the survey is roughly constant across the surveyed latitudes (Figure 14). Additionally, petrale sole were captured across the survey depths between 55-500 m (Figure 15).

The data from the petrale sole was analyzed using a spatio-temporal delta model implemented as an R package, VAST (Thorson and Barnett 2017, Thorson 2019), described above in Section 2.1.1. Spatial variation was approximated using 250 knots, and the model used the bias-correction algorithm (Thorson and Kristensen 2016) in Template Model Builder (Kristensen et al. 2016). The index of abundance was estimated using VAST separately for the early and late periods of the survey. The stratifications are provided in Tables 7 and 8.

The gamma distribution with random strata-year and vessel effects was chosen as the final model for both the early and late time periods. The Q-Q plots do not show any departures

from the assumed distribution (Figures 17 and 20). The Pearson residuals for the encounter and catch rate for the early and late periods are shown in Figures 18, 19, 21, and 22.

The estimated index of abundance is shown in Table 4. For contrast, the 2015 model estimated, the 2019 design based, and the 2019 VAST indices are shown in Figure 23 and the estimated density of petrale sole is show in Figures 24 and 25. The index for the Triennial Survey across the early and late period shows an slight increase in the population between 1980 and 2001 with a spike in the final year of 2004.

Length bins from 12 to 62 cm in 2 cm increments were used to summarize the length frequency of the survey catches in each year. Table 9 shows the number of lengths taken by the survey. The first bin includes all observations less than 14 cm and the last bin includes all fish larger than 62 cm. The length frequency distributions for the Triennial Survey from 1980-2004 are shown in Figures 26 and 27.

There are no petrale sole age data from the Triennial Survey.

The input sample sizes for length data were calculated using the same approach for the NWFSC West Coast Groundfish Bottom Trawl Survey data described in Section 2.1.1.

2.2 Fishery-Dependent Data

`fishery-dependent-data`

2.2.1 Commercial Fishery Landings

`commercial-fishery-landings`

All landings for this update assessment were summarized by port of landing, where available, as well as for a northern fleet consisting of Washington and Oregon and a southern fleet consisting of California. Landings for Washington and Oregon are summed into a single northern fleet due to the fact that vessels commonly fish and land in each other's waters and ports.

The PacFIN database (1981-2018 for California and Washington; 1987-2018 for Oregon) extracted XXX ADD DATE XXX. Historical catches were not updated from the previous assessment in 2013. The 2013 assessment historical Washington catches were obtained from WDFW landings reconstruction for 1935, 1939 and 1949- 1969 (pers. comm. T. Tsou and G. Lippert) and the Pacific Marine Fisheries Commission (PMFC) Data Series for 1956-1980 (PFMC 1979). The 2013 assessment historical Oregon landings were obtained from reconstruction for 1932 to 1986 (Karnowski et al. 2014). The 2013 assessment historical California landings used catch reconstruction data extending from 1931-1980 (Ralston et al. 2010) and California Department of Fish and Game (CDFG) Fish Bulletins for 1916-1930 landings (Heimann and Carlisle 1970) as reconstructed by Lai et al. (2005). The California fishery began in 1876 but no landings data are available from 1876-1915. Therefore a linear interpolation between landings of 1 ton in 1876 and the landings recorded for 1916 are used to filling this period.

Landings for the fishing year, beginning on 1 November, are summarized by fleet in Table 1 and Figure 1. The landings of petrale sole by gear types other than groundfish-trawl have been inconsequential, averaging less than 2.5% of the coast-wide landings. The non-trawl landings are included in the trawl landings.

2.2.2 Discards

discards

Data on discards of petrale sole are available from two different data sources. The earliest source is referred to as the Pikitch data and comes from a study organized by Ellen Pikitch that collected trawl discards from 1985-1987 (Pikitch et al. 1988). The northern and southern boundaries of the study were 48°42' N latitude and 42°60' N latitude respectively, which is primarily within the Columbia INPFC area (Pikitch et al. 1988, Rogers and Pikitch 1992). Participation in the study was voluntary and included vessels using bottom, midwater, and shrimp trawl gears. Observers of normal fishing operations on commercial vessels collected the data, estimated the total weight of the catch by tow, and recorded the weight of species retained and discarded in the sample. Results of the Pikitch data were obtained from John Wallace (personal communication, NWFSC, NOAA) in the form of ratios of discard weight to retained weight of petrale sole and sex-specific length frequencies. The Pikitch discard estimates were applied to both the summer and winter northern fisheries and are shown in Table 10.

The second source is from the West Coast Groundfish Observer Program (WCGOP). This program is part of the NWFSC and has been recording discard observations since 2003. Table 10 shows the discard ratios (discarded/(discarded + retained)) of petrale sole from WCGOP. Since 2011, when the trawl rationalization program was implemented, observer coverage rates increased to nearly 100% for all the limited entry trawl vessels in the program and discard rates declined compared to pre-2011 rates. Discard rates were obtained for both the catch-share and the non-catch share sector for petrale sole. A single discard rate was calculated by weighting discard rates based on the commercial landings by each sector. Coefficient of variations were calculated for the non-catch shares sector and pre-catch share years by bootstrapping vessels within ports because the observer program randomly chooses vessels within ports to be observed. The discard rates from WCGOP are shown in Table 10.

Starting in 2015 a small number of vessels switched to electronic monitoring discards at sea (4, 7, and 8 vessels in 2015, 2016, and 2017 respectively) rather than a human observer and as of this update assessment only 3 years of data are available. Discarding rates at sea of petrale sole by these vessels were very low, near zero. This update assessment did not evaluate these data to estimate an electronic monitoring specific discard rate, but rather applied the discard ratio from the observed vessels in the WCGOP database. Future assessments should evaluate this assumption in greater detail.

Discard mean body weight data were obtained from the WCGOP data and used in this update assessment for each of the four fishing fleets. The mean body weight of discarded fish

from each fleet are shown in Figures 28 - 31. The summer fisheries, both north and south, had relatively large sample numbers which is reflected in a lower CV by year relative to the winter fisheries.

Discard length composition data were obtained from the WCGOP data and used in this update assessment to estimate retention curves for each of the four fishing fleets. The discard length data from each fleet are shown in Figures 32 and 33.

The data, historical and current, provided by the WCGOP are updated annually based on the most recent standards of QA/QC methods. Hence, these data can have minor changes over time. To ensure the data from the years since the last update assessment (2014-2018) were consistent with the earlier data, data from all years were replaced based to reflect the current standings of the WCGOP data.

2.2.3 Foreign Landings

`foreign-landings`

The impact of landings of petrale sole by foreign fishing fleets prior to the institution of the exclusive economic zone (EEZ) of the U.S. West Coast is currently not quantified and remains an area for research.

2.2.4 Historical Commercial Catch-Per-Unit Effort/Logbooks

`historical-commercial-catch-per-unit-effortlogbooks`

Commercial logbook data for petrale sole was first used to construct CPUE indices of abundance in the 1999 assessment for Oregon fleets from 1987-1997 (Sampson and Lee 1999). Since the first inclusion 1999, the commercial CPUE indices were extended and or updated based on management changes and new statistical methods through 2009. For additional information on the use of CPUE indices in the assessment of petrale sole please see the 2013 assessment (Haltuch et al. 2013b).

CPUE calculations for the Winter fishery on aggregations of petrale sole described in the 2013 assessment were retained for this assessment (Haltuch et al. 2013b) (Figures 35 and 36). Two CPUE indices from 1987-2009 with catchability modeled as a power function are used in this update assessment, one for the north and south winter fisheries.

2.2.5 Fishery Length and Age Data

`fishery-length-and-age-data`

The PacFIN BDS database contains data from Oregon Department of Fish and Wildlife (ODFW; 1966-present) and Washington Department of Fish and Wildlife (WDFW; 1955-present), but only 1986-present data from California Department of Fish and Game (CDFG).

The CDFG data set for the years 1948-1992 was extracted and provided from CALCOM by Brenda Erwin (CDFG) in 2011.

The historical Oregon data for petrale sole has changed substantially since 2015. The state has identified samples across years that were not collected according to the state's standardized sampling protocol that were included in PacFIN due to error. These samples likely represent samples that were collected for special projects. Oregon has removed some of these samples for petrale sole from PacFIN, but not all as of 2019. To remove the remaining non-standard samples, Oregon PacFIN data was filtered according to the "Sample Quality" based on directions from Ali Whitman at ODFW where samples with a sample quality code of 63 represent non-standardized samples. Filtering out these records removed all Oregon samples from 1966-1970 and 1972-1975 and a portion of samples from 1971.

Commercial length-frequency distributions based on the fishing year were developed for each state for which observations were available. For each fleet, the raw observations (compiled from the PacFIN and CalCOM databases) were expanded to the sample level, to allow for any fish that were not measured, then to the trip level to account for the relative size of the landing from which the sample was obtained. The expanded length observations were then expanded by the landings in each state for the combined Washington and Oregon fleet. Age frequencies were computed in the same manner, except that age observations for Washington and Oregon were not combined due to aging error considerations.

Length and age data collected from commercial landings for each fleet are summarized by the number of trips and fish sampled by year (Tables 11 and 12). Figures 32, 33, and 34 show plots of the commercial length and age composition data across time for each fishery fleet.

The calculation for input sample sizes for the commercial length and age data was done to be consistent with the 2015 update assessment. The input sample size for commercial lengths and ages were set equal to the number of trips by year for each fleet.

2.3 Biological Data

biological-data

2.3.1 Natural Mortality

natural-mortality

The instantaneous rate of natural mortality for a wild fish population is notoriously difficult to estimate. One accepted method is to examine the age distribution of an unexploited or lightly exploited stock. This method cannot readily be applied to petrale sole given the long history of exploitation off the U.S. West Coast. Ketchen and Forrester (1966) estimated that the natural mortality coefficients were $0.18\text{-}0.26 \text{ yr}^{-1}$ for males and $0.19\text{-}0.21 \text{ yr}^{-1}$ for females based on a catch curve analysis of 1943-1945 Washington trawl data from Swiftsure Bank, off the southwest corner of Vancouver Island. However, petrale sole catches were relatively high during mid-1940s through the 1950s. Starr and Fargo (2004) estimated the instantaneous

rate of natural mortality (M) using Hoenig's method (Hoenig 1983) estimating M values of 0.22 and 0.15 yr^{-1} were estimated given maximum ages of 20 and 30 years, respectively.

An archived set of commercial samples, collected from Northern California between the late 1950s and early 1980s, recently found that multiple samples were aged between 20-31 years old, suggesting a similar range of M values for U.S. West Coast petrale sole. U.S. stock assessments prior to 2009 and current British Columbia stock assessments assumed a value of $M = 0.2 \text{ yr}^{-1}$ for both sexes. The 2013 stock assessment used a meta-analysis value produced the following normal prior distributions for females (mean = 0.151, sd = 0.16) and males (0.206, sd = 0.218) based on early research by Owen Hamel (pers. comm.) with maximum age for females and males of 32 and 29 years, respectively.

Hamel (2015) refined and published a method for combining meta-analytic approaches relating the M rate to other life-history parameters such as longevity, size, growth rate, and reproductive effort to provide a prior on M . In that same issue of *ICES Journal of Marine Science*, Then et al. (2015) provided an updated data set of estimates of M and related life history parameters across a large number of fish species from which to develop an M estimator for fish species in general. They concluded by recommending M estimates be based on maximum age alone, based on an updated Hoenig non-linear least squares estimator $M = 4.899A_{\max}^{-0.916}$. The approach of basing M priors on maximum age alone was one that was already being used for West Coast rockfish assessments. However, in fitting the alternative model forms relating M to A_{\max} , Then et al. (2015) did not consistently apply their transformation. In particular, in real space, one would expect substantial heteroscedasticity in both the observation and process error associated with the observed relationship of M to A_{\max} . Therefore, it would be reasonable to fit all models under a log transformation. This was not done. Re-evaluating the data used in Then et al. (2015) by fitting the one-parameter A_{\max} model under a log-log transformation (such that the slope is forced to be -1 in the transformed space (Hamel 2015)), the point estimate for M is:

$$M = \frac{5.4}{A_{\max}}$$

The above is also the median of the prior. The prior is defined as a lognormal distribution with mean $\ln(5.4/A_{\max})$ and SE = 0.438.

The natural mortality prior was updated for this update assessment using the above approach. Maximum age was assumed to be 32 and 29 years for females and males, respectively, the same assumption applied in the 2013 assessment. Using the Hamel et al. approach above, the prior value for females in regular space is 0.169 and for males is 0.186.

2.3.2 Maturation and Fecundity

maturation-and-fecundity

Petrale sole maturity-at-length information is generally sparse in space and time, has not been collected in a systematic fashion across time, is of varying quality, and does not always agree

between studies. It is possible that maturity may have changed over time. However, it is not possible to assess this quantitatively owing to differences in when historical samples on which maturity ogives could be based were taken, and how maturity stage (visual vs. histological) was determined. The 2005 petrale sole assessment used the most recent study for the West Coast of the U.S. that was based on observations collected during 2002 from Oregon and Washington (Hannah et al. 2002). The 50% size-at-maturity was estimated at 33.1 cm with maturity asymptoting to 1.0 for larger fish (Figure 37).

To date, there has been limited information regarding fecundity at age or length of petrale sole. The 2013 stock assessment assumed that fecundity of female petrale sole was equal to biomass (Figure 38). Since the last full assessment, new research has been done examining the fecundity of petrale sole (Lefebvre et al. n.d.). The study concluded a difference in fecundity between California and Washington petrale sole where a 40 cm fish in California is more fecund compared to northern fish of the same size (Figure 39). However, northern fish of the largest size were more fecund relative to fish in California. The current petrale sole model is a single area coastwide model, which assumes fish along the U.S. have the same biology (e.g. natural mortality, growth, fecundity). The estimates of fecundity for petrale sole were considered new data and based on the guidelines for update stock assessments, these data were not included in the base model. However, a sensitivity to including these data was provided. The next full assessment should include the new data about fecundity at length.

2.3.3 Sex Ratio

sex-ratio

Past assessments of petrale sole have assumed a 50% sex ratio between females and males off the U.S West Coast. Similarly, Canadian data from the 2004 published stock assessment also suggests sex ratios of petrale sole in British Columbia are generally 50% males and 50% females (Starr and Fargo 2004). To be consistent with the full assessment this update assessment retains the equal sex ratio assumption. However, examining the NWFSC West Coast Groundfish Bottom Trawl Survey data the proportion of females in the population across the mid-range lengths is approximately 0.40 with the proportion increasing to 1 at the largest lengths due to dimorphic growth (Figure 40). The next full assessment should evaluate the sex ratio for petrale sole.

2.3.4 Length-Weight Relationship

length-weight-relationship

The length-weight relationship for petrale sole was estimated outside the model using all biological data available from the NWFSC West Coast Groundfish Bottom Trawl Survey data, where the female weight-at-length in grams was estimated at $2.08e-06L^{3.47}$ and males at $3.05e-06L^{3.36}$ where L is length in cm (Figures 41).

2.3.5 Growth (Length-at-Age)

growth-length-at-age

The length-at-age was estimated for male and female petrale sole. Figure 42 shows the lengths and ages as well as predicted von Bertalanffy fits to the data from the fishery and the NWFSC West Coast Groundfish Bottom Trawl Survey data. Females grow larger than males and sex-specific growth parameters were estimated at the following values:

XXX DOUBLE CHECK THESE VALUES WHEN AGES ARRIVE XXX

Females $L_\infty = 54$; $k = 0.16$

Males $L_\infty = 41$; $k = 0.25$

These values were used as starting parameter values within the base model prior to estimating each parameter for male and female petrale sole.

2.3.6 Ageing Precision and Bias

ageing-precision-and-bias

Historically, petrale sole otoliths have been read by multiple ageing labs using surface and break and burn methods. In order to conduct a comprehensive estimation of ageing bias and imprecision, the 2009 assessment compiled and analyzed all of the available double-read data from the state of Oregon, the Cooperative Aging Project (CAP), and the Washington Department of Fish and Wildlife (WDFW), as well information from a bomb radiocarbon age validation study for petrale sole off the U.S. West Coast (Haltuch and Hicks 2009, Haltuch et al. (2013a)).

The 2013 stock assessment applied read method and lab specific ageing error vectors (Haltuch et al. 2013b). The same approach to ageing error based on data source are age reading method applied in the 2013 assessment was applied in this update stock assessment. The ageing error vectors are shown in Tables 13 and 14. For a detailed description please see the 2013 stock assessment (Haltuch et al. 2013b).

2.3.7 Environmental and Ecosystem Data

environmental-and-ecosystem-data

This update assessment did not evaluate potential ecosystem data and methodologies for petrale sole.

3 Assessment Model

assessment-model

3.1 History of Modeling Approaches Used for This Stock

history-of-modeling-approaches-used-for-this-stock

Early stock assessments only assessed petrale sole in the combined U.S.-Vancouver and Columbia INPFC areas, i.e. petrale sole in these areas were treated as a unit stock, using time series of data that began during the 1970s (Demory 1984, Turnock et al. 1993). The first assessment used stock reduction analysis and the second assessment used the length-based Stock Synthesis model. The third petrale sole assessment utilized the hybrid length-and-age-based Stock Synthesis 1 model, using data from 1977-1998 (Sampson and Lee 1999). During the 1999 stock assessment an attempt was made to include separate area assessments for the Eureka and Monterey INPFC areas but acceptable models could not be configured due to a lack of data (Sampson and Lee 1999).

The 2005 petrale sole assessment was conducted as two separate stocks, the northern stock encompassing the U.S. Vancouver and Columbia INPFC areas and the southern stock including the Eureka, Monterey and Conception INPFC areas, using Stock Synthesis 2, a length-age structured model. Both the northern- and southern-area models specified the fishing year as beginning on November 1 and continuing through October 31 of the following year, with a November-February winter fishery and a March-October summer fishery. Landings prior to 1957 were assumed to have been taken during the summer season in years where monthly data were not available to split the catches seasonally. The complete catch history was reconstructed for petrale sole for the 2005 stock assessment, with the northern area model starting in 1910 and the southern area model in 1876. In 2005, the STAR panel noted that the petrale sole stock trends were similar in both northern and southern areas, in spite of the different modeling choices made for each area, and that a single coast-wide assessment should be considered. The 2009 and 2011 assessments treated petrale sole as a single coast-wide stock, with the fleets and landings structured by state (WA, OR, CA) area of catch. During the 2011 STAR panel concerns were raised regarding the difficulty of discriminating landings from Washington and Oregon waters, particularly in light of the Oregon historical landings reconstruction that includes a summary of data by port of landing but not by catch area, due to the fact that the Oregon and Washington vessels commonly fish in each other's waters and land in each other's ports. The availability of the historical comprehensive landings reconstruction for Oregon by port of landing lead the STAR panel to recommend combining the Washington and Oregon fleets within the coast-wide stock assessment using port of landing rather than catch area. Starting with the 2013 stock assessment, the coast-wide stock assessment now summarizes petrale sole landings by the port of landing and combines Washington and Oregon into a single fleet (Haltuch et al. 2013b). The 2015 this 2019 update assessment assumes the same approach as the 2013 stock assessment.

3.2 General Model Specifications and Assumptions

general-model-specifications-and-assumptions

Stock Synthesis version 3.30.03.13 was used to estimate the parameters in the model (Methot and Wetzel 2013). R4SS, version 1.33.2, along with R version 3.4.3 were used to investigate and plot model fits. A summary of the data sources used in the model (details discussed above) is shown in Figure 2.

3.2.1 Changes Between the 2015 Update and Current Assessment Model

changes-between-the-2015-update-and-current-assessment-model

As with the 2013 petrale sole stock assessment, the current model is implemented as a single-area model. The current update assessment has been upgraded to a new version of SS (3.30.13). A thorough description of the 2013 assessment model, which is used in this update assessment, is presented separately below; this section linking the two models is intended to clearly identify where substantive changes were made. These changes include:

1. Fitting using SS v.3.30.13.
2. Added commercial fishery catch data (2015-2018).
3. Added composition data from the commercial fishery (length and age data 2015-2018) and recalculated data expansions based upon the current methods.
4. Reprocessed all discard data sources and added discard rate, average weight, and length composition data (2014-2017).
5. Added 2015-2018 NWFSC West Coast Groundfish Bottom Trawl Survey data and calculated the index of abundance VAST.
6. Added NWFSC West Coast Groundfish Bottom Trawl Survey length and age data 2015-2018.
7. Triennial Survey early and late indices of abundance were calculated using VAST.
8. Model tuning to re-weight data.
9. Length-weight relationship parameters estimated outside of the stock assessment model from the NWFSC West Coast Groundfish Bottom Trawl Survey data up to 2018 and input as fixed values.
10. Update the natural mortality prior for female and male fish.

The general model set-up is described in Table 15.

3.2.2 Summary of Fleets and Areas

summary-of-fleets-and-areas

Fishery removals were divided among 4 fleets: 1) winter North trawl, 2) summer North trawl, 3) winter South trawl, and 4) summer South trawl. Landings for the North fleet are defined as fish landed in Washington and Oregon ports. Landings for the South fleet are defined as fish landed in California ports. Other removals are very small and are included in the trawl fishery removals. The data available for each fleet are described in Figure 2.

3.2.3 Priors

priors

Priors were applied only to parameters for steepness (h) and natural mortality (M). The steepness prior is based on the Myers (1999) meta-analysis of flatfish steepness and the natural mortality prior is based on a meta-analysis completed by Hamel (2015). The prior for steepness assumed a beta distribution with a mean equal to 0.80 (Figure 44).

The natural mortality prior was updated for this update assessment using the Hamel meta-analysis approach. Maximum age was assumed to be 32 and 29 years for females and males (Figure 43), respectively, the same assumption regarding maximum age as applied in the 2013 assessment.

3.2.4 Data Weighting

data-weighting

Length and conditional-age-at-length compositions from the NWFSC West Coast Groundfish Bottom Trawl Survey were fit along with length and marginal age compositions from the fishery and the Triennial Survey. Length data started with a input sample size determined from the equation listed in Sections 2.1.1 (survey data) and 2.2.5 (fishery data). It was assumed for conditional-age-at-length data that each age was a random sample within the length bin and the model started with a sample size equal to the number of fish in that length bin.

The update assessment model was weighted using the McAllister and Ianelli (1997) method (Harmonic Mean weighting), consistent with the 2015 update assessment. The McAllister and Ianelli data weight approach looks at the difference between individual observations and predictions. A sensitivity was performed examining the difference between alternative weighting approaches. The weights applied to each length and age data set for the base model are shown in Table 16.

3.2.5 Estimated and Fixed Parameters

estimated-and-fixed-parameters

There were 304 estimated parameters in the base model. These included one parameters for R_0 , natural mortality, steepness, growth, selectivity, retention, time blocking of the fleets and

the surveys, commercial CPUE catchability, recruitment deviations, and forecast recruitment deviations (Table 17).

Fixed parameters in the model were as follows. The standard deviation of recruitment deviates was fixed at 0.40. Maturity-at-length was fixed as described above in Section 2.3.2. Length-weight parameters were fixed at estimates using all length-weight observations (Figure 41).

3.2.6 Key Assumptions and Structural Choices

key-assumptions-and-structural-choices

All structural choices for stock assessment models are likely to be important under some circumstances. In this update assessment update these choices are generally made to be consistent with the previous assessment (Haltuch et al. 2013b). Major choices in the structuring of this stock assessment model include a coast-wide model with seasonal fleet structure for two regions, north and south, splitting the Triennial Survey into an early and late time period, and estimates of selectivity and retention curves for each fleet.

3.2.7 Bridging Analysis

bridging-analysis

The exploration of models began by bridging from the 2015 update assessment to Stock Synthesis version 3.30.03.13, which produced no discernible difference (Figure 45).

3.2.8 Convergence

convergence

Proper convergence was determined by starting the minimization process from dispersed values of the maximum likelihood estimates to determine if the model found a better minimum. Starting parameters were jittered by 10%. This was repeated 50 times and a better minimum was not found (Table 18). The model did not experience convergence issues when provided reasonable starting values. Through the jittering done as explained above and likelihood profiles, we are confident that the base model as presented represents the best fit to the data given the assumptions made. There were no difficulties in inverting the Hessian to obtain estimates of variability, although much of the early model investigation was done without attempting to estimate a Hessian.

3.3 Base Model Results

base-model-results

The base model parameter estimates along with approximate asymptotic standard errors are shown in Table 17 and the likelihood components are shown in Table 19. Estimates of

derived reference points and approximate 95% asymptotic confidence intervals are shown in Table 20. Estimates of stock size over time are shown in Table 21.

3.3.1 Parameter Estimates

parameter-estimates

Natural mortality by sex was estimated directly within the model. Natural mortality was estimated to be 0.162 for female fish and 0.174 for male fish. In comparison the estimates from the 2015 assessment were 0.145 and 0.154 for female and male fish, respectively.

Steepness was also estimated within the model, consistent with the approach applied in the 2013 full and 2015 update assessment. The estimate of steepness from the Beverton-Holt stock recruitment curve was estimated at 0.84. The previous update assessment estimated a steepness of 0.89.

The estimates of maximum length and the von Bertalanffy growth coefficient, k , were less than the external estimates for males and female but were well within the 95% confidence interval given the estimated uncertainty (Table 17). The estimated k for female fish was consistent with the value estimated in the 2015 update assessment (0.135 versus 0.134), but the estimated k for male fish was higher than the value estimated in 2015 (0.216 versus 0.203). The majority of growth for female and male petrale sole growth occurs at younger ages, reaching near maximum length by age 10-15, depending upon sex, with female petrale sole reaching larger maximum lengths (Figure 46). The spawning output estimated was equal to the spawning weight of female fish (Figure 47).

Selectivity curves were estimated for the fishery and survey fleets. The estimated selectivities for the fishery fleets are shown in Figure 48. All fishery selectivities were estimated to be asymptotic, reaching maximum selectivity for fish between 35 and 40 cm. Shifts in selectivities were estimated for each fleet fishery were estimated based on time blocks assumed in the 2013 assessment (Figure 48). The estimated retention curves for each fleet based on the historical time blocks and discarded length composition data are shown in Figure 49. Sex specific survey selectivities were assumed to be asymptotic and are shown in Figure 50.

The catchability for each of the winter CPUE time series were estimated as power functions. The Winter North base catchability value was estimated at 0.006 with the exponent parameter at -0.354. The Winter South base catchability value was estimated at 0.255 with the exponent parameter at -0.849.

Additional survey variability, process error added directly to each year's input variability, for the Triennial Survey, both early and late, was estimated within the model. The model estimated a added variance of 0.341 for the early time period of and 0.236 for the late period.

The time-series of estimated recruitments shows a relationship with the decline in spawning output, punctuated by larger recruitments (Figures 51 and 52) in recent years (2006, 2007,

and 2008). There is little information regarding recruitment prior to 1960 and the uncertainty in those estimates is expressed in the model. The five largest estimated recruitment estimated with the model (in ascending order) occurred in 2006, 2007, 1998, 1966, and 2008. The four lowest recruitments estimated within the model (in ascending order) occurred in 1986, 1992, 1973, and 1987.

3.3.2 Fits to the Data

fits-to-the-data

There are numerous types of data for which the fits are discussed: fishery CPUE, survey abundance indices, discard data (rates, mean body weights, and length compositions), length-composition data for the fisheries and surveys, marginal age compositions for the fisheries, and conditional age-at-length observations for the NWFSC West Coast Groundfish Bottom Trawl Survey.

The fit to the CPUE for the winter fisheries is show in Figures 54, 55, 56, and 57. The model fits both of the CPUE time-series relatively well. The fits to the survey indices are shown in Figures 58, 59, and 60. In order to fit the early and the late periods of the Triennial Survey extra standard error was required. The trend in the early time-series of the Triennial Survey was generally not consistent with other data within the model. The final year, 2004, in the late period of the Triennial Survey was under fit by the model. The petrale sole survey index from the NWFSC West Coast Groundfish Bottom Trawl Survey was generally fit well. However, the most recent year, 2018 data point was over fit by the model.

The observed WCGOP discard rates (Figures 61 - 64) were fit by each fishery using time blocks. The time blocks on the discard data was based on those define in the 2013 assessment (Haltuch et al. 2013a) with the final block starting in 2011 being extended through the final model year. The discarding rates over time by each fleet are shown in Figure 65. Fits to the discard rates for the northern fleets from the Pikitch data in 1985-1987 were either under (Figure 61) or over fit (Figure 62) which is consistent to the estimates from the 2015 update assessment. Fits to the WCGOP observed mean body weights are shown in Figures 66 - 69. The fits to the discard mean body weights to the summer fleets were generally better than the data from the winter fisheries which had more variable observations and lower number of observations (hence larger annual uncertainties).

Fits to the length data are shown based on the proportions of lengths observed by year and the Pearson residuals-at-length for all fleets. Detailed fits to the length data by year and fleet are provided in Appendix A, section 11. Aggregate fits by fleet are shown in Figure 70. There are a few things that stand out when examining the aggregated length composition data. First, the sexed discard lengths from the Pikitch study appear to be poorly fit by the model but this is related to small sample sizes. However, the unsexed discard lengths from the WCGOP data for each fleet were fit well by the model.

Discard lengths from WCGOP were fit well by the model and show no obvious pattern in the residuals (Figures 71 - 74). The residuals to the fishery lengths clearly showed the growth

differential between males and females where the majority of positive residuals at larger sizes were from female fish (Figures 75 - 78). Notably, the Summer North fishery has a large positive residual pattern for male fish between 1966-1980. A similar pattern in the Pearson residuals was observed in the 2013 full and the 2015 update assessment (Haltuch et al. 2013b, Stawitz et al. 2015). The residuals for each of the surveys are shown in Figures 79, 80, and 81. The Pearson residuals from the NWFSC West Coast Groundfish Bottom Trawl Survey shows indications of the 2008 cohort moving through the population. Length data were weighted according to the McAllister Ianelli Harmonic mean weights. The relationship between the observed (input) sample size to the effective sample sizes after weighting are shown in Figures 85 - 88.

Age data were fitted to as marginal age compositions for the fishery fleets. The NWFSC West Coast Groundfish Bottom Trawl Survey ages were treated as conditional age-at-length data to facilitate the estimation of growth within the model. The aggregated fits to the marginal age data are shown in Figure 89. The aggregated age data were general fit well for the fishery fleets, however, the peaks of each of the age data were often under fit by the model which was also observed in the 2013 assessment (Haltuch et al. 2013b). Detailed fits to the age data by year and fleet are provided in Appendix B, section 12. The Pearson residuals for the fishery fleets are shown in Figures 90 - ??.

The observed and expected conditional age-at-length fits for NWFSC West Coast Groundfish Bottom Trawl Survey are shown in Figures 94 - ???. The fits generally match the observations. The Pearson residuals are shown in Figure 98 and 99.

The age data were also weighted according to the McAllister Ianelli Harmonic mean weights. The relationship between the observed (input) sample size to the effective sample sizes after weighting are shown in Figures 103 - 103.

3.3.3 Population Trajectory

[population-trajectory](#)

The predicted spawning biomass is given in Table 21 and plotted in Figure 104. The predicted spawning biomass time series shows a strong decline from the late-1930s through the mid-1960s, followed by a small recovery through the mid-1970s, and another decline to its lowest point during the early 1990s. This general pattern of stock decline is coincident with increasing catches and the movement of the fishery from the south to the north, and from summer fishing in shallow waters to winter fishing on spawning aggregations in deeper waters. From the mid-1990s through 2005 the stock increased slightly, then declined through 2010 (Figure 104). The stock has increased strongly since 2010 in response to reduced catches and in response to above average recruitment in 2006, 2007, and 2008. The estimated total biomass follows the same general trend as observed in the spawning biomass (Figure 105). The 2019 estimated spawning biomass relative to unfished equilibrium spawning biomass is above the target of 25% of unfished spawning biomass at 32.3% (Figure 106). Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in

the estimated spawning biomass is generally low. The standard deviation of the log of the spawning output in 2019 is 0.11.

Recruitment deviations were estimated for the entire time-series that was modeled (Figure 51 and discussed in Section 3.3.1) and provide a realistic portrayal of uncertainty. The time series of estimated recruitments shows a relationship with the decline in spawning output, punctuated by larger recruitments in 2006, 2007, and 2008. The five largest estimated recruitment estimated with the model (in ascending order) occurred in 2006, 2007, 1998, 1966, and 2008. The four lowest recruitments estimated within the model (in ascending order) occurred in 1986, 1992, 1973, and 1987. The stock-recruit curve resulting from a value of estimated steepness, 0.84, is shown in Figure 107 with estimated recruitments also shown.

3.3.4 Sensitivity Analyses

sensitivity-analyses

A number of sensitivity analyses were conducted. Each of the sensitivities conducted was a single exploration from the base model assumptions and/or data, and were not performed in a cumulative fashion.

1. Fix natural mortality value for female fish at a lower value of ??.
2. Fix natural mortality value for female fish at a higher value of ??.
3. Use the natural mortality prior for female and male fish used in the 2015 update assessment, natural mortality estimated.
4. Use the coastwide fecundity relationship for petrale sole estimated by Lefebvre et al. (in press).
5. Estimate the sex ratio between female and male fish within the model.
6. Data weight according to the Francis method using the weighting values shown in Table 23.
7. Data weight according to the Dirichlet method where the estimated parameters are shown in Table 24.

Likelihood values and estimates of key parameters from each sensitivity are available in Table 22. Plots of the estimated time-series of spawning biomass and relative spawning biomass are shown in Figures 108 and 109

3.3.5 Retrospective Analysis

retrospective-analysis

A five-year retrospective analysis was conducted by running the model using data only through 2014, 2015, 2016, 2017 and 2018 (Figures 110, 111, and 112). The initial scale of the spawning biomass trended upward relative to the base model. Overall, no alarming patterns were present in the retrospective analysis.

3.3.6 Historical Analysis

historical-analysis

The estimated summary biomass from previous assessments since 2005 are shown in Figure 113. The current assessment estimated a slight increase in initial spawning biomass compared to previous assessments.

3.3.7 Likelihood Profiles

likelihood-profiles

Likelihood profiles were conducted for R_0 , steepness, and female natural mortality values separately. These likelihood profiles were conducted by fixing the parameter at specific values and estimated the remaining parameters based on the fixed parameter value.

For steepness, the negative log-likelihood supported values between 0.70 - 0.95 (Figure 114). Likelihood components by data source show that the age data support a higher steepness value. The surveys generally provide very little information concerning steepness. The relative spawning biomass for petrale sole diverges most during the middle of the time series based on the assumed values of steepness with the final status generally being above the management target biomass (Figure 115).

The negative log-likelihood was minimized at a female natural mortality value of ??, but the 95% confidence interval extends over values ranging from ?? - ?. Male natural mortality was estimated in the likelihood profile. The age and length data likelihood contribution was minimized at natural mortality values ranging from ??-?? (Figure ??). The relative spawning biomass for petrale sole widely varied across alternative values of natural mortality (Figure 117).

In regards to values of R_0 , the negative log-likelihood was minimized at approximately $\log(R_0)$ of 9.83 (Figure 118).

3.3.8 Reference Points

reference-points-1

Reference points were calculated using the estimated selectivities and catch distributions among fleets in the most recent year of the model (2018). Sustainable total yields (landings

plus discards) were 2,819.8 mt when using an $SPR_{30\%}$ reference harvest rate and with a 95% confidence interval of 2,590.1 mt based on estimates of uncertainty. The spawning biomass equivalent to 25% of the unfished spawning output ($SB_{25\%}$) was 7,638.7.

The predicted spawning biomass from the base model generally showed a decline beginning during the 1950s and reaching a low in spawning biomass in 1993 with the stock declining to 5.8% relative stock size (Figures 104 and 106). Since 20010, the spawning biomass has been increasing due to small catches and above average recruitment. The 2019 spawning biomass relative to unfished equilibrium spawning biomass is above the target of 25% of unfished (Figure 106). The fishing intensity, 1-SPR, exceeded the current harvest rate limit ($SPR_{30\%}$) throughout the late 1970s until approximately 2010 as seen in Figure 119. Recent exploitation rates on petrale sole were estimated to be less than target levels.

Table 20 shows the full suite of estimated reference points for the base model and Figure 120 shows the equilibrium curve based on a steepness value estimated at 0.84.

4 Harvest Projections and Decision Tables

harvest-projections-and-decision-tables

5 Regional Management Considerations

regional-management-considerations

Currently petrale sole are managed using a coast-wide harvest; therefore this assessment does not provide a recommended method for allocating harvests regionally. The resource is modeled as a single stock. There is currently no genetic evidence that there are distinct biological stocks of petrale sole off the U.S. coast and the limited tagging data that describes adult movement suggests that movement may be significant across depth and latitude.

6 Research Needs

research-needs

There are many areas of research that could be improved to benefit the understanding and assessment of petrale sole. Below, are issues that are considered of importance.

1. In the past many assessments have derived historical catches independently. The states of California and Oregon have completed comprehensive historical catch reconstructions. At the time of this assessment, a comprehensive historical catch reconstruction is not available for Washington. Completion of a Washington catch reconstruction would provide the best possible estimated catch series that accounts for all the catch and better resolves historical catch uncertainty for flatfish as a group.

2. Due to limited data, new studies on both the maturity and fecundity relationships for petrale sole would be beneficial.
3. The effect of the implementation of the IFQ (catch shares) program that began during 2011 on fleet behavior, including impacts on discards, fishery selectivity, and fishing locations would benefit from further study.
4. Studies on stock structure and movement of petrale sole, particularly with regard to the winter-summer spawning migration of petrale sole and the likely trans-boundary movement of petrale sole between U.S. and Canadian waters seasonally.
5. The extent of spatial variability on productivity processes such as growth, recruitment, and maturity is currently unknown and would benefit from further research.

7 Acknowledgments

acknowledgments

Many people were instrumental in the successful completion of this assessment and their contribution is greatly appreciated.

8 References

references

Alderdice, D., and Forrest, C. 1971. Effects of salinity and temperature on embryonic development of the petrale sole(*Eopsetta jordani*). Journal of Fisheries Research Board Canada **28**: 727–744.

Alverson, D., and Chatwin, B. 1957. Results from tagging experiments on a spawning stock of petrale sole, *Eopsetta jordani* (Lockington). Journal of Fisheries Research Board Canada **14**: 953–974.

Anon. 2001. Fish stocks of the Pacific coast. Fisheries; Oceans Canada.

Best, E. 1960. Petrale Sole. In: California ocean fisheries resources to the year 1960. California Department of Fish; Game.

Best, E. 1963. Movements of petrale sole, *Eopsetta jordani*, tagged off of California. Pacific Marine Fisheries Commission Bulletin **6**: 24–38.

Bradburn, M., Keller, A., and Horness, B. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.

Castillo, G. 1992. Fluctuations of year-class strength of petrale sole (*Eopsetta jordani*) and their relation to environmental factors. Master's thesis, Oregon State University.

Castillo, G. 1995. Latitudinal patterns in reproductive life history traits of northeast Pacific flatfish. In Proceedings of the International Symposium on North Pacific Flatfish. Alaska Sea Grant, University of Alaska Fairbanks. pp. 51–72.

Castillo, G., Li, H., and Golden, J. 1993. Environmental induced recruitment variation in petrale sole, *Eopsetta jordani*. Fisheries Bulletin **92**: 481–493.

Cook, R., He, X., Maguire, J., and Tsou, T. 2009. Petrale sole STAR panel report. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Demory, R. 1984. Progress report on the status of petrale sole in the INPFC Columbia-Vancouver areas in 1984. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Eschmeyer, W., and Herald, E. 1983. A field guide of Pacific coaast fishes North America. Houghton Mifflin CO, Boston, MA.

Fargo, J. 1997. Flatfish stock assessments for the West Coast of Canada for 1997 and

recommended yield options for 1998. Can. Stock Assess. Sec. Res. Doc.

Gates, D., and Frey, H. 1974. Designated common names of certain marine organisms of California. Fish Buletin **161**: 55–90.

Gregory, P., and Jow, T. 1976. The validity of otoliths as indicators of age of petrale sole from California. California Department of Fish and Game **62**(2): 132–140.

Haltuch, M.A., and Hicks, A.C. 2009. Status of the U.S. petrale sole resource ien 2008. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Haltuch, M.A., Hamel, O.S., Piner, K.R., McDonald, P., Kastelle, C.R., and Field, J.C. 2013a. A California Current bomb radiocarbon reference chronology and petrale sole (*Eopsetta jordani*) age validation. Canadian Journal of Fisheries and Aquatic Sciences **70**(1): 22–31. doi: [10.1139/cjfas-2011-0504](https://doi.org/10.1139/cjfas-2011-0504).

Haltuch, M.A., Ono, K., and Valero, J.L. 2013b. Status of the U.S. petrale sole resource in 2012. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Hamel, O.S. 2015. A method for calculating a meta-analytical prior for the natural mortality rate using multiple life history correlates. ICES Journal of Marine Science: Journal du Conseil **72**(1): 62–69. doi: [10.1093/icesjms/fsu131](https://doi.org/10.1093/icesjms/fsu131).

Hannah, R., Parker, S., and Fruth, E. 2002. Length and age at maturity of female petrale sole (*Eopsetta jordani*) determined from samples collected prior to spawning aggregation. U.S. Fish Bulletin **100**: 711–719.

Harry, G. 1956. Analysis and history of the Oregon otter-trawl fishery. PhD Thesis, University of Washington, Seattle, WA.

Harry, G. 1959. Time of spawning, length at maturity, and fecundity of the English, petrale, and dover soles (*Parophrys vetulus*, *Eopsetta jordani*, and *Microstomus pacificus*, respectively). Fisheries Commission of Oregon, Research Briefs **7**(1): 5–13.

Heimann, R., and Carlisle, J. 1970. Pacific Fishes of Canada. California Department of Fish and Game Fish Bulletin **149**.

Helser, T., Punt, A.E., and Methot, R.D. 2004. A generalized linear mixed model analysis of a multi-vessel fishery resource survey. **70**: 251–264.

Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. Fishery Bulletin **82**: 898–903.

Karnowski, M., Gertseva, V., and Stephens, A. 2014. Historical Reconstruction of Oregon's

Commercial Fisheries Landings. Oregon Department of Fish; Wildlife, Salem, OR.

Ketchen, K., and Forrester, C. 1966. Population dynamics of the petrale sole, *Eopsetta jordani*, in waters off western Canada. Fish. Res. Bd. Canada Bull.

Kramer, D., Barss, W., Paust, B., and Bracken, B. 1995. Guide to northeast Pacific flatfishes: Families Bothidae, Cynoglossidae, and Pleuronectidae. Alaska Sea Grant, University of Alaska Fairbanks.

Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H.J., and Bell, B. 2016. TMB: Automatic Differentiation and Laplace Approximation. Journal of Statistical Software **70**: 1–21.

Lai, H., Haltuch, M.A., Punt, A.E., and Cope, J. 2005. Stock assessment of petrale sole: 2004. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Lefebvre, L.S., Friedlander, C., and Field, J.C. (n.d.). Reproductive ecology and size-dependent fecundity in the petrale sole, *Eopsetta jordani*, in California, Oregon, and Washington waters. Fishery Bulletin.

Love, M. 1996. Probably more than you want to know about the fishes of the Pacific Coast. Really Big Press, Santa Barbara, California.

Love, M., Mecklenburg, C., Mecklenburg, T., and Thorsteinson, L. 2005. Resource inventory of marine and estuarine fishes of the West Coast and Alaska: A checklist of north Pacific and arctic ocean species from Baja California to the Alsaka-Yukon border. USGS, Seattle, WA.

Mantua, N.J., Hare, S.R., Zhang, Y., Wallace, J.R., and Francis, R.C. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. Bull. Am. Meteorol. Soc. **78**: 1069–1080.

McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling - importance resampling algorithm. Canadian Journal of Fisheries and Aquatic Sciences **54**: 284–300.

Methot, R.D., and Wetzel, C.R. 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. Fisheries Research **142**: 86–99. doi: [10.1016/j.fishres.2012.10.012](https://doi.org/10.1016/j.fishres.2012.10.012).

Myers, R.A., Bowen, K.G., and Barrowman, N. 1999. Maximum reproductive rate of fish at low population sizes. Canadian Journal of Fisheries and Aquatic Sciences **56**: 2404–2419. Available from <http://www.nrcresearchpress.com/doi/pdf/10.1139/f99-201> [accessed 4 October 2016].

Pedersen, M. 1975. Movements and growth of petrale sole (*Eopsetta jordani*) tagged off

Washington and southwest Vancouver Island. Fishery Research Board of Canada Progress Report.

PFMC. 1979. Data series, groundfish section. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Pikitch, E.K., Erikson, D., and Wallace, J.R. 1988. An evaluation of the effectiveness of trip limits as a management tool. NOAA, NMFSC.

Porter, P. 1964. Notes on fecundity, spawning, and early life history of petrale sole (*Eopsetta jordani*) with descriptions of flatfish larvae collected in the Pacific Ocean off Humboldt Bay, California. Master's thesis, Humboldt State College.

Ralston, S., Pearson, D.E., Field, J.C., and Key, M. 2010. Documentation of the California catch reconstruction project. US Department of Commerce, National Oceanic; Atmospheric Adminstration, National Marine.

Reilly, P., Wilson-Vandenberg, D., Lea, R., Wilson, C., and Sullivan, M. 1994. Recreational angler's guide to the common nearshore fishes of Northern and Central California. California Department of Fish; Game.

Rogers, J.B., and Pikitch, E.K. 1992. Numerical definition of groundfish assemblages caught off the coasts of Oregon and Washington using commercial fishing strategies. Canadian Journal of Fisheries and Aquatic Sciences **49**(12): 2648–2656.

Sampson, D., and Lee, Y. 1999. An assessment of the stocks of petrale sole off Washington, Oregon, and Northern California in 1998. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Scofield, W. 1948. Trawling gear in California. California Fish and Game Fish Bulletin **72**: 1–60.

Smith, R. 1937. Report on the Puget Sound otter trawl investigations. Master's thesis, University of Washington.

Starr, P.J., and Fargo, J. 2004. Petrale sole stock assessment for 2003 and recommendations for management in 2004. CSAS Res. Doc 2004/036.

Stawitz, C.C., Hurtado-Ferro, F., Kuriyama, P.T., Trochta, J., Johnson, K.F., Haltuch, M.A., and Hamel, O.S. 2015. Stock assessment update: Status of the U.S. petrale sole resource in 2014. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Then, A.Y., Hoenig, J.M., Hall, N.G., and Hewitt, D.A. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200

fish species. ICES Journal of Marine Science **72**(1): 82–92. doi: [10.1093/icesjms/fsu136](https://doi.org/10.1093/icesjms/fsu136).

Thorson, J.T. 2019. Guidance for decisions using the Vector Autoregressive Spatio-Temporal (VAST) package in stock, ecosystem, habitat and climate assessments. Fisheries Research **210**: 143–161. doi: [10.1016/j.fishres.2018.10.013](https://doi.org/10.1016/j.fishres.2018.10.013).

Thorson, J.T., and Barnett, L.A.K. 2017. Comparing estimates of abundance trends and distribution shifts using single- and multispecies models of fishes and biogenic habitat. ICES Journal of Marine Science: Journal du Conseil **74**(5): 1311–1321. doi: [10.1093/icesjms/fsw193](https://doi.org/10.1093/icesjms/fsw193).

Thorson, J.T., and Kristensen, K. 2016. Implementing a generic method for bias correction in statistical models using random effects, with spatial and population dynamics examples. Fisheries Research **175**: 66–74. doi: [10.1016/j.fishres.2015.11.016](https://doi.org/10.1016/j.fishres.2015.11.016).

Thorson, J.T., and Ward, E.J. 2013. Accounting for space–time interactions in index standardization models. Fisheries Research **147**: 426–433. doi: [10.1016/j.fishres.2013.03.012](https://doi.org/10.1016/j.fishres.2013.03.012).

Turnock, J., Wilkins, M., Saelens, M., and Wood, C. 1993. Status of West Coast petrale sole in 1993. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.

Weinberg, K., Wilkins, M., Shaw, F., and Zimmermann, M. 2002. The 2001 Pacific West Coast bottom trawl survey of groundfish resources: Estimates of distribution, abundance and length and age composition. NOAA Technical Memorandum, U.S. Department of Commerce.

9 Tables

tables

Table 1: Landings for each fleet for the modeled years.

Year	Winter North	Summer North	Winter South	Summer South	tab:Comm_Catch
1875	0	0	0	0	
1876	0	0	0	1	
1877	0	0	0	1	
1878	0	0	0	1	
1879	0	0	0	1	
1880	0	0	0	12	
1881	0	0	0	22	
1882	0	0	0	33	
1883	0	0	0	43	
1884	0	0	0	54	
1885	0	0	0	64	
1886	0	0	0	75	
1887	0	0	0	85	
1888	0	0	0	96	
1889	0	0	0	106	
1890	0	0	0	117	
1891	0	0	0	128	
1892	0	0	0	138	
1893	0	0	0	149	
1894	0	0	0	159	
1895	0	0	0	170	
1896	0	0	0	180	
1897	0	0	0	191	
1898	0	0	0	201	
1899	0	0	0	212	
1900	0	0	0	223	
1901	0	0	0	233	
1902	0	0	0	244	
1903	0	0	0	254	
1904	0	0	0	265	
1905	0	0	0	275	
1906	0	0	0	286	
1907	0	0	0	296	
1908	0	0	0	307	
1909	0	0	0	318	
1910	0	0	0	328	
1911	0	0	0	339	
1912	0	0	0	349	
1913	0	0	0	360	
1914	0	0	0	370	

Year	Winter North	Summer North	Winter South	Summer South
1915	0	0	0	381
1916	0	0	0	386
1917	0	0	0	526
1918	0	0	0	424
1919	0	0	0	333
1920	0	0	0	230
1921	0	0	0	294
1922	0	0	0	425
1923	0	0	0	427
1924	0	0	0	533
1925	0	0	0	528
1926	0	0	0	522
1927	0	0	0	632
1928	0	0	0	620
1929	0	2	0	706
1930	0	1	0	659
1931	0	81	63	531
1932	2	251	36	520
1933	6	408	39	392
1934	10	568	139	896
1935	14	650	155	777
1936	16	770	95	432
1937	20	1051	75	741
1938	27	1187	48	890
1939	35	1545	31	1029
1940	39	1737	162	597
1941	41	1803	111	331
1942	46	2919	24	216
1943	51	2867	72	345
1944	55	2047	86	447
1945	60	1866	102	439
1946	64	2492	72	1116
1947	69	1778	154	1093
1948	74	2315	273	1778
1949	76	1809	617	1812
1950	156	2322	424	1638
1951	118	1666	208	993
1952	131	1390	326	882
1953	46	737	533	981
1954	27	903	801	1073

Year	Winter North	Summer North	Winter South	Summer South
1955	57	863	526	1052
1956	137	759	508	801
1957	171	1103	527	1027
1958	99	1152	568	957
1959	332	947	379	723
1960	241	1374	520	644
1961	217	1547	542	1029
1962	295	1512	515	859
1963	663	1038	534	978
1964	282	1090	378	927
1965	370	950	374	853
1966	366	972	325	925
1967	409	793	532	874
1968	284	811	361	871
1969	190	887	421	848
1970	412	1081	472	1071
1971	743	883	540	1016
1972	730	1017	703	1000
1973	497	1272	417	742
1974	517	1611	665	893
1975	539	1559	561	901
1976	506	951	713	737
1977	682	743	484	495
1978	746	1098	419	801
1979	734	1086	353	945
1980	382	976	518	680
1981	761	468	360	895
1982	1041	771	262	502
1983	696	935	273	361
1984	416	739	260	329
1985	392	553	273	471
1986	474	714	403	355
1987	855	573	311	556
1988	743	610	349	411
1989	696	583	393	415
1990	641	460	319	373
1991	793	397	448	310
1992	640	366	272	307
1993	685	392	237	234
1994	518	355	246	299

Year	Winter North	Summer North	Winter South	Summer South
1915	0	0	0	381
1916	0	0	0	386
1917	0	0	0	526
1918	0	0	0	424
1919	0	0	0	333
1920	0	0	0	230
1921	0	0	0	294
1922	0	0	0	425
1923	0	0	0	427
1924	0	0	0	533
1925	0	0	0	528
1926	0	0	0	522
1927	0	0	0	632
1928	0	0	0	620
1929	0	2	0	706
1930	0	1	0	659
1931	0	81	63	531
1932	2	251	36	520
1933	6	408	39	392
1934	10	568	139	896
1935	14	650	155	777
1936	16	770	95	432
1937	20	1051	75	741
1938	27	1187	48	890
1939	35	1545	31	1029
1940	39	1737	162	597
1941	41	1803	111	331
1942	46	2919	24	216
1943	51	2867	72	345
1944	55	2047	86	447
1945	60	1866	102	439
1946	64	2492	72	1116
1947	69	1778	154	1093
1948	74	2315	273	1778
1949	76	1809	617	1812
1950	156	2322	424	1638
1951	118	1666	208	993
1952	131	1390	326	882
1953	46	737	533	981
1954	27	903	801	1073

Year	Winter North	Summer North	Winter South	Summer South
1995	591	454	236	287
1996	591	440	406	394
1997	621	430	448	442
1998	522	577	221	300
1999	463	504	287	267
2000	610	586	372	241
2001	691	597	308	260
2002	667	714	335	195
2003	544	713	256	180
2004	1010	750	177	271
2005	964	1069	343	533
2006	537	1012	125	454
2007	930	536	404	475
2008	842	354	519	414
2009	847	642	470	250
2010	264	292	78	121
2011	224	427	40	78
2012	410	494	124	108
2013	513	1013	130	280
2014	853	860	273	386
2015	1040	1077	215	354
2016	865	1168	237	235
2017	1142	1271	201	393
2018	957	1262	218	402

Table 2: Recent trend in estimated total catch relative to management guidelines. The estimated total catch includes the total landings plus the model estimated discard mortality based upon discard rate data.

Year	OFL (mt; ABC prior to 2011)	ACL (mt; OY prior to 2011)	Total landings (mt)	Estimated total catch (mt)
2009	2811	2433	2209	2329
2010	2751	1200	755	867
2011	1021	976	768	791
2012	1275	1160	1135	1159
2013	2711	2592	1936	1973
2014	2774	2652	2373	2398
2015	3073	2816	2686	2727
2016	3208	2910	2506	2543
2017	3208	3136	3008	3050
2018	3152	3013	2840	2882

Table 3: Description of the strata used to create the indices for the NWFSC Shelf-Slope survey.

Strata	Depth Lower Bound	Depth Upper Bound	Latitude South	Latitude North	tab:strata_nwfsc
Shallow Vancouver	55	100	47.5	49.0	
Shallow Columbia	55	100	43.0	47.5	
Shallow Eureka	55	100	40.5	43.0	
Shallow Monterey	55	100	36.0	40.5	
Shallow Conception	55	100	34.5	36.0	
Mid Vancouver	100	183	47.5	49.0	
Mid Columbia	100	183	43.0	47.5	
Mid Eureka	100	183	40.5	43.0	
Mid Monterey	100	183	36.0	40.5	
Mid Conception	100	183	34.5	36.0	
Deep Van/Col/Eur	183	549	40.5	49.0	
Deep Monterey	183	549	36.0	40.5	
Deep Conception	183	549	32.0	36.0	

Table 4: Summary of the fishery-independent biomass/abundance time-series used in the stock assessment. The standard error includes the input annual standard error and model estimated added variance.

Year	Winter N.		Winter S.		Triennial Early		Triennial Late		NWFSC Combo		tab:Index_Summary
	Obs	SE	Obs	SE	Obs	SE	Obs	SE	Obs	SE	
1980	-	-	-	-	1512	0.49	-	-	-	-	-
1983	-	-	-	-	2380	0.45	-	-	-	-	-
1986	-	-	-	-	2249	0.46	-	-	-	-	-
1987	1.09	0.28	1.08	0.56	-	-	-	-	-	-	-
1988	1.16	0.27	0.91	0.33	-	-	-	-	-	-	-
1989	0.92	0.27	0.53	0.43	3569	0.43	-	-	-	-	-
1990	0.76	0.28	0.96	0.46	-	-	-	-	-	-	-
1991	0.86	0.27	0.90	0.36	-	-	-	-	-	-	-
1992	0.56	0.28	0.59	0.68	2226	0.43	-	-	-	-	-
1993	0.56	0.27	0.86	0.35	-	-	-	-	-	-	-
1994	0.50	0.28	0.71	0.30	-	-	-	-	-	-	-
1995	0.66	0.28	0.90	0.30	-	-	2636	0.33	-	-	-
1996	0.77	0.29	1.25	0.30	-	-	-	-	-	-	-
1997	0.85	0.28	0.82	0.28	-	-	-	-	-	-	-
1998	1.01	0.29	0.93	0.31	-	-	3836	0.32	-	-	-
1999	0.71	0.29	0.83	0.29	-	-	-	-	-	-	-
2000	0.67	0.28	0.62	0.29	-	-	-	-	-	-	-
2001	0.83	0.27	0.66	0.29	-	-	4362	0.33	-	-	-
2002	0.93	0.28	0.80	0.29	-	-	-	-	-	-	-
2003	1.02	0.28	0.85	0.29	-	-	-	-	19970	0.13	-
2004	1.63	0.28	1.71	0.31	-	-	10662	0.33	26767	0.14	-
2005	1.85	0.28	1.93	0.29	-	-	-	-	26795	0.12	-
2006	2.01	0.28	1.58	0.29	-	-	-	-	22288	0.12	-
2007	2.04	0.28	2.07	0.28	-	-	-	-	21003	0.13	-
2008	1.96	0.27	1.62	0.28	-	-	-	-	17597	0.12	-
2009	2.12	0.27	1.76	0.28	-	-	-	-	18270	0.12	-
2010	-	-	-	-	-	-	-	-	26860	0.12	-
2011	-	-	-	-	-	-	-	-	36324	0.12	-
2012	-	-	-	-	-	-	-	-	42005	0.12	-
2013	-	-	-	-	-	-	-	-	58743	0.13	-
2014	-	-	-	-	-	-	-	-	65532	0.12	-
2015	-	-	-	-	-	-	-	-	59015	0.12	-
2016	-	-	-	-	-	-	-	-	64522	0.12	-
2017	-	-	-	-	-	-	-	-	66101	0.12	-
2018	-	-	-	-	-	-	-	-	46594	0.12	-

Table 5: Summary of NWFSC shelf-slope survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size	tab:NWcombo_Lengths
2003	197	2837	589	
2004	212	3346	674	
2005	278	4555	907	
2006	247	3668	753	
2007	257	3409	727	
2008	257	3047	677	
2009	277	3387	744	
2010	325	6052	1160	
2011	320	6176	1172	
2012	295	5372	1036	
2013	218	3445	693	
2014	332	4822	997	
2015	312	4236	897	
2016	309	4385	914	
2017	314	4261	902	
2018	291	3783	813	

Table 6: Summary of NWFSC shelf-slope survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size	tab:NWcombo_Ages
2003	173	765	279	
2004	167	723	267	
2005	237	752	341	
2006	236	774	343	
2007	196	690	291	
2008	225	746	328	
2009	258	777	365	
2010	297	801	408	
2011	289	799	399	
2012	269	777	376	
2013	217	843	333	
2014	318	766	424	
2015	291	751	395	

Table 7: Description of the strata used to create the indices for the Triennial Early (1980 - 1992) survey.

Strata	Depth Lower Bound	Depth Upper Bound	Latitude South	Latitude North	tab:strata_tri_early
Shallow Van/Col	55	100	43.0	49.0	
Shallow Eureka	55	100	40.5	43.0	
Shallow Mon/Con	55	100	32.0	40.5	
Deep Van/Col/Eur	100	400	40.5	49.0	
Deep Mon/Con	100	400	32.0	40.5	

Table 8: Description of the strata used to create the indices for the Triennial Late (1995-2004) survey.

Strata	Depth Lower Bound	Depth Upper Bound	Latitude South	Latitude North	tab:strata_tri_late
Shallow Van/Col	55	100	43.0	49.0	
Shallow Eureka	55	100	40.5	43.0	
Shallow Mon/Con	55	100	32.0	40.5	
Deep Van/Col	100	500	43.0	49.0	
Deep Eureka	100	500	40.5	43.0	
Deep Mon/Con	100	500	36.0	40.5	
Deep Con	100	500	32.0	36.0	

Table 9: Summary of Triennial survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1980	1	16	3
1983	2	30	6
1986	36	540	111
1989	141	1419	337
1992	116	1015	256
1995	145	1369	334
1998	236	2624	598
2001	254	3016	670
2004	239	4676	884

Table 10: Summary of discard rates used in the model by each data source (continued on next page).

Year	Fleet	Discard Rate	Standard Error	Data Source	tab:Discard
1985	WinterN	0.022	0.110	Pikitch	
1986	WinterN	0.021	0.116	Pikitch	
1987	WinterN	0.027	0.119	Pikitch	
2002	WinterN	0.008	0.001	WGCOP	
2003	WinterN	0.004	0.002	WGCOP	
2004	WinterN	0.003	0.002	WGCOP	
2005	WinterN	0.002	0.001	WGCOP	
2006	WinterN	0.006	0.003	WGCOP	
2007	WinterN	0.012	0.005	WGCOP	
2008	WinterN	0.022	0.012	WGCOP	
2009	WinterN	0.027	0.014	WGCOP	
2010	WinterN	0.119	0.023	WGCOP	
2011	WinterN	0.002	0.015	WGCOP	
2012	WinterN	0.001	0.015	WGCOP	
2013	WinterN	0.001	0.015	WGCOP	
2014	WinterN	0.003	0.015	WGCOP	
2015	WinterN	0.001	0.015	WGCOP	
2016	WinterN	0.001	0.015	WGCOP	
2017	WinterN	0.003	0.015	WGCOP	
2018	WinterN	0.001	0.015	WGCOP	
2019	WinterN	0.001	0.015	WGCOP	
1985	SummerN	0.035	0.042	Pikitch	
1986	SummerN	0.034	0.043	Pikitch	
1987	SummerN	0.032	0.045	Pikitch	
2002	SummerN	0.186	0.023	WGCOP	
2003	SummerN	0.105	0.022	WGCOP	
2004	SummerN	0.083	0.023	WGCOP	
2005	SummerN	0.042	0.008	WGCOP	
2006	SummerN	0.078	0.015	WGCOP	
2007	SummerN	0.116	0.021	WGCOP	
2008	SummerN	0.051	0.016	WGCOP	
2009	SummerN	0.206	0.067	WGCOP	
2010	SummerN	0.099	0.029	WGCOP	
2011	SummerN	0.037	0.015	WGCOP	
2012	SummerN	0.022	0.015	WGCOP	
2013	SummerN	0.017	0.015	WGCOP	
2014	SummerN	0.026	0.015	WGCOP	
2015	SummerN	0.006	0.015	WGCOP	
2016	SummerN	0.017	0.015	WGCOP	
2017	SummerN	0.007	0.015	WGCOP	

Year	Fleet	Discard Rate	Standard Error	Data Source
2002	WinterS	0.035	0.016	WGCOP
2003	WinterS	0.012	0.001	WGCOP
2004	WinterS	0.013	0.033	WGCOP
2005	WinterS	0.033	0.004	WGCOP
2006	WinterS	0.071	0.035	WGCOP
2007	WinterS	0.012	0.003	WGCOP
2008	WinterS	0.013	0.010	WGCOP
2009	WinterS	0.024	0.009	WGCOP
2010	WinterS	0.052	0.031	WGCOP
2011	WinterS	0.001	0.015	WGCOP
2012	WinterS	0.001	0.015	WGCOP
2013	WinterS	0.003	0.015	WGCOP
2014	WinterS	0.001	0.015	WGCOP
2015	WinterS	0.001	0.015	WGCOP
2016	WinterS	0.003	0.015	WGCOP
2017	WinterS	0.006	0.015	WGCOP
2018	WinterS	0.001	0.015	WGCOP
2019	WinterS	0.001	0.015	WGCOP
2002	SummerS	0.058	0.016	WGCOP
2003	SummerS	0.033	0.011	WGCOP
2004	SummerS	0.033	0.014	WGCOP
2005	SummerS	0.012	0.003	WGCOP
2006	SummerS	0.038	0.014	WGCOP
2007	SummerS	0.065	0.023	WGCOP
2008	SummerS	0.026	0.014	WGCOP
2009	SummerS	0.023	0.006	WGCOP
2010	SummerS	0.056	0.007	WGCOP
2011	SummerS	0.041	0.015	WGCOP
2012	SummerS	0.013	0.015	WGCOP
2013	SummerS	0.004	0.015	WGCOP
2014	SummerS	0.004	0.015	WGCOP
2015	SummerS	0.010	0.015	WGCOP
2016	SummerS	0.004	0.015	WGCOP
2017	SummerS	0.008	0.015	WGCOP

Table 11: Summary of fishery length samples used in the stock assessment (continued on next page).

Year	Winter N.		Summer N.		Winter S.		Summer S.		tab:Fishery_Lengths
	Trips	Fish	Trips	Fish	Trips	Fish	Trips	Fish	
1948	0	0	0	0	4	202	4	203	
1949	0	0	0	0	6	275	4	183	
1955	1	507	0	0	0	0	0	0	
1956	0	0	1	534	0	0	0	0	
1960	0	0	1	644	0	0	0	0	
1962	0	0	0	0	0	0	3	150	
1964	0	0	0	0	2	73	22	897	
1965	0	0	0	0	1	25	14	583	
1966	0	0	2	463	20	852	33	1396	
1967	0	0	3	485	12	481	44	1815	
1968	0	0	7	1842	13	499	87	3414	
1969	1	328	4	992	19	705	49	1907	
1970	1	237	5	1309	6	226	29	920	
1971	3	721	6	1481	12	519	37	1180	
1972	2	516	14	3255	21	747	39	1435	
1973	2	440	4	829	18	752	40	1460	
1974	3	768	25	7196	28	974	35	1133	
1975	9	1978	12	3509	8	325	19	873	
1976	1	379	4	1054	10	475	26	1255	
1977	2	321	21	2410	16	739	38	1816	
1978	3	678	2	570	9	448	33	1649	
1979	6	618	23	2438	5	247	13	601	
1980	4	573	22	2287	20	999	81	4042	
1981	4	400	0	0	31	1522	65	3134	
1982	0	0	0	0	30	1496	34	1434	
1983	0	0	0	0	17	851	33	1600	
1984	0	0	0	0	13	627	19	943	
1985	0	0	0	0	8	400	17	825	
1986	0	0	0	0	22	1100	32	1602	
1987	6	300	16	805	12	600	29	1450	
1988	10	499	8	401	10	500	12	532	
1989	3	151	13	652	16	783	18	900	
1990	5	251	11	552	10	428	2	76	
1991	10	356	7	277	22	754	2	82	
1992	8	313	11	428	6	176	0	0	
1993	8	236	8	296	0	0	0	0	
1994	6	258	9	371	1	1	0	0	
1995	6	230	2	66	0	0	0	0	
1996	2	67	4	168	0	0	0	0	
1997	8	284	11	417	0	0	0	0	
1998	5	201	22	1004	0	0	0	0	
1999	11	413	15	703	0	0	0	0	
2000	17	638	24	1012	0	0	0	0	
2001	12	468	18	786	10	305	9	289	
2002	13	551	31	1259	7	209	10	252	

Year	Winter N.		Summer N.		Winter S.		Summer S.	
	Trips	Fish	Trips	Fish	Trips	Fish	Trips	Fish
2003	28	872	35	1370	10	254	30	475
2004	22	720	30	1328	10	228	15	431
2005	18	628	35	1493	9	169	36	966
2006	26	1106	51	2639	37	1040	47	1059
2007	42	1680	46	2402	58	1656	103	2971
2008	65	2059	36	2127	66	2023	97	2442
2009	32	1220	66	2860	34	749	62	1597
2010	49	1614	59	1795	29	655	52	1356
2011	26	855	47	2019	33	1170	23	400
2012	32	1059	44	1954	28	1099	40	1125
2013	55	2145	52	2300	40	1753	43	1930
2014	59	2158	64	2421	35	1292	49	1672
2015	61	1929	60	2386	34	1062	62	2026
2016	31	1045	39	1071	34	1311	70	2306
2017	57	1816	74	2790	33	1289	85	2489
2018	50	1386	93	2654	19	823	77	2663
2019	5	225	6	158	0	0	0	0

Table 12: Summary of fishery age samples used in the stock assessment (continued on next page).

tab:Fishery_Ages

Year	Winter		Summer		Winter		Summer	
	N. Trips	Fish	N. Trips	Fish	S. Trips	Fish	S. Trips	Fish
1960	0	0	1	168	0	0	0	0
1966	0	0	2	340	19	441	27	649
1967	0	0	3	482	2	50	11	273
1968	0	0	3	663	4	64	56	1340
1969	1	100	2	192	12	293	31	765
1970	1	116	4	499	5	126	29	709
1971	2	318	5	785	12	294	37	930
1972	2	349	13	1984	21	512	38	962
1973	2	393	4	684	16	425	37	951
1974	3	295	20	2032	27	642	34	837
1975	8	765	10	1012	7	175	18	473
1976	1	99	4	400	10	250	23	575
1977	2	197	19	1853	10	241	33	822
1978	3	308	2	387	6	150	32	799
1979	4	350	21	2214	4	99	11	270
1980	2	177	16	1568	12	300	50	1243
1981	2	195	0	0	10	250	27	677
1982	0	0	0	0	7	175	18	352
1983	0	0	0	0	9	276	8	191
1984	0	0	0	0	2	49	3	74
1985	0	0	0	0	2	50	4	100
1986	0	0	0	0	11	261	16	376
1987	6	173	16	573	5	117	12	291
1988	10	379	8	256	5	123	6	149
1989	3	144	12	507	0	0	0	0
1990	5	160	11	272	10	294	1	38
1991	10	201	7	151	8	245	0	0
1992	8	313	11	424	0	0	0	0
1993	8	234	8	296	0	0	0	0
1994	6	256	9	371	0	0	0	0
1995	6	228	2	66	0	0	0	0
1996	2	67	4	165	0	0	0	0
1997	8	283	10	375	0	0	0	0
1998	5	201	22	999	0	0	0	0
1999	6	256	13	608	0	0	0	0
2000	6	258	12	560	0	0	0	0
2001	5	250	11	498	0	0	0	0
2002	8	346	20	834	0	0	0	0
2003	20	665	26	1071	2	41	5	55
2004	7	313	24	1059	2	57	4	96
2005	6	294	18	874	3	55	10	217
2006	4	197	14	697	2	51	7	154
2007	14	536	24	1018	4	78	5	97
2008	11	337	26	1079	7	97	18	300
2009	28	400	39	684	0	0	3	78
2010	19	353	34	542	0	0	0	0
2011	24	326	42	845	8	185	8	26
2012	31	385	40	835	4	118	1	34
2013	48	723	46	831	1	39	3	100
2014	29	678	24	616	0	0	0	0

Year	Winter N.		Summer N.		Winter S.		Summer S.	
	Trips	Fish	Trips	Fish	Trips	Fish	Trips	Fish
2015	56	584	48	811	0	0	0	0
2016	2	95	0	0	0	0	0	0
2017	5	246	7	339	0	0	0	0
2018	5	249	23	494	0	0	0	0

Table 13: Estimated ageing error vectors applied to ages read by the Cooperative Aging Project lab used in the assessment model.

`tab:age_error1`

True Age	Break and Burn		Surface		Combo		Surface Pre-1990	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
0	0.26	0.17	0.16	0.12	0.47	0.13	0.00	0.00
1	1.35	0.17	1.27	0.12	1.42	0.13	0.71	0.00
2	2.41	0.23	2.35	0.18	2.37	0.25	2.02	0.08
3	3.44	0.29	3.41	0.25	3.32	0.38	3.24	0.17
4	4.45	0.36	4.43	0.32	4.27	0.51	4.38	0.26
5	5.44	0.44	5.42	0.40	5.22	0.64	5.44	0.35
6	6.41	0.52	6.39	0.49	6.17	0.76	6.44	0.46
7	7.35	0.61	7.33	0.59	7.12	0.89	7.36	0.56
8	8.28	0.71	8.25	0.70	8.07	1.02	8.22	0.67
9	9.18	0.81	9.14	0.82	9.02	1.14	9.03	0.79
10	10.06	0.92	10.01	0.96	9.97	1.27	9.78	0.92
11	10.92	1.04	10.85	1.11	10.92	1.40	10.48	1.05
12	11.76	1.18	11.67	1.27	11.87	1.53	11.14	1.19
13	12.58	1.32	12.47	1.45	12.82	1.65	11.75	1.34
14	13.38	1.48	13.24	1.66	13.77	1.78	12.32	1.49
15	14.17	1.64	14.00	1.88	14.72	1.91	12.85	1.66
16	14.94	1.82	14.73	2.12	15.67	2.03	13.35	1.83
17	15.68	2.02	15.45	2.39	16.62	2.16	13.81	2.01

Table 14: Estimated ageing error vectors applied to ages read by Washington Department of Fish and Wildlife used in the assessment model.

`tab:age_error2`

True Age	Combo		Surface		Break and Burn	
	Mean	SD	Mean	SD	Mean	SD
0	0.49	0.13	0.13	0.10	0.50	0.15
1	1.46	0.13	1.32	0.10	1.51	0.15
2	2.44	0.27	2.47	0.21	2.52	0.30
3	3.42	0.40	3.58	0.31	3.52	0.45
4	4.39	0.53	4.64	0.41	4.53	0.60
5	5.37	0.67	5.67	0.52	5.53	0.75
6	6.35	0.80	6.66	0.62	6.54	0.90
7	7.32	0.93	7.62	0.72	7.55	1.05
8	8.30	1.07	8.54	0.83	8.55	1.20
9	9.28	1.20	9.43	0.93	9.56	1.35
10	10.25	1.33	10.28	1.03	10.57	1.51
11	11.23	1.47	11.11	1.13	11.57	1.66
12	12.21	1.60	11.90	1.24	12.58	1.81
13	13.18	1.74	12.67	1.34	13.59	1.96
14	14.16	1.87	13.41	1.44	14.59	2.11
15	15.14	2.00	14.12	1.55	15.60	2.26
16	16.11	2.14	14.81	1.65	16.60	2.41
17	17.09	2.27	15.47	1.75	17.61	2.56

Table 15: Specifications of the model for petrale sole.

Model Specification	tab:Model_setup
Starting year	1876
<u>Population characteristics</u>	
Maximum age	40
Gender	2
Population lengths	4-78 cm by 2 cm bins
Summary biomass (mt)	Age 3+
<u>Data characteristics</u>	
Data lengths	12-62 cm by 2 cm bins
Data ages	1-17 ages
Minimum age for growth calculations	2
Maximum age for growth calculations	17
First mature age	3
<u>Fishery characteristics</u>	
Fishing mortality method	Hybrid
Maximum F	3
Catchability - Fishery	Power
Catchability - Survey	Analytical estimate
Winter North selectivity	Double Normal
Summer North selectivity	Double Normal
Winter South selectivity	Double Normal
Summer South selectivity	Double Normal
AFSC/NWFSC West Coast Triennial	Double Normal
Shelf Survey - early	
AFSC/NWFSC West Coast Triennial	Double Normal
Shelf Survey - late	
NWFSC West Coast Groundfish Bottom	Double Normal
Trawl Survey	
<u>Fishery time blocks</u>	
Fishery selectivity	1876-1972, 1973-1982, 1983-1992, 1993-2002, 2003-2010, 2011-2018
Winter retention	1876-2002, 2003-2009, 2010, 2011-2018
Summer retention	1876-2002, 2003-2008, 2009-2010, 2011-2018

Table 16: Data weights applied when using harmonic data weighting.

Fleet	Lengths	Ages	tab:harm
Winter North			
Summer North			
Winter South			
Summer South			
Triennial Early survey		-	
Triennial Late survey		-	
NWFSC shelf-slope survey			

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
NatM_p_1_Fem_GP_1	0.161992	6	(0.005, 0.5)	OK	0.02	Log_Norm (-1.7793, 0.438)
L_at_Amin_Fem_GP_1	15.8915	2	(10, 45)	OK	0.42	None
L_at_Amax_Fem_GP_1	53.9424	3	(35, 80)	OK	0.40	None
VonBert_K_Fem_GP_1	0.135371	2	(0.04, 0.5)	OK	0.01	None
SD_young_Fem_GP_1	0.189896	3	(0.01, 1)	OK	0.01	None
SD_old_Fem_GP_1	0.0250563	4	(0.01, 1)	OK	0.01	None
Wtlen_1_Fem_GP_1	0.00000208	-3	(-3, 3)			Normal (0.00000208, 0.8)
Wtlen_2_Fem_GP_1	3.4737	-3	(1, 5)			Normal (3.4737, 0.8)
Mat50%_Fem_GP_1	33.1	-3	(10, 50)			Normal (33.1, 0.8)
Mat_slope_Fem_GP_1	-0.743	-3	(-3, 3)			Normal (-0.743, 0.8)
Eggs/kg_inter_Fem_GP_1	1	-3	(-3, 3)			Normal (1, 1)
Eggs/kg_slope_wt_Fem_GP_1	0	-3	(-3, 3)			Normal (0, 1)
NatM_p_1_Mal_GP_1	0.174411	6	(0.005, 0.6)	OK	0.02	Log_Norm (-1.6809, 0.438)
L_at_Amin_Mal_GP_1	16.4779	2	(10, 45)	OK	0.33	None
L_at_Amax_Mal_GP_1	42.1411	3	(35, 80)	OK	0.41	None
VonBert_K_Mal_GP_1	0.216	2	(0.04, 0.5)	OK	0.01	None
SD_young_Mal_GP_1	0.134634	3	(0.01, 1)	OK	0.01	None
SD_old_Mal_GP_1	0.053	4	(0.01, 1)	OK	0.01	None
Wtlen_1_Mal_GP_1	0.00000305	-3	(-3, 3)			Normal (0.00000305, 0.8)
Wtlen_2_Mal_GP_1	3.36054	-3	(-3, 5)			Normal (3.36054, 0.8)
CohortGrowDev	1	-4	(0, 1)			None
FracFemale_GP_1	0.5	-99	(0.01, 0.99)			None
SR_LN(R0)	9.83235	1	(5, 20)	OK	0.20	None
SR_BH_stEEP	0.839916	5	(0.2, 1)	OK	0.05	Normal (0.8, 0.09)
SR_sigmaR	0.4	-99	(0, 2)			Normal (0.9, 5)
SR_regime	0	-2	(-5, 5)			Normal (0, 0.2)
SR_autocorr	0	-99	(0, 0)			None
Early_InitAge_31	0.000000278911	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_30	0.000000327699	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_29	0.000000384942	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_28	0.000000452078	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_27	0.000000530781	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_26	0.000000622998	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_25	0.000000730985	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_24	0.000000857355	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_23	0.00000100513	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_22	0.00000117777	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_21	0.00000137929	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_20	0.00000161425	3	(-4, 4)	act	0.40	dev (NA, NA)

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
Early_InitAge_19	0.00000188786	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_18	0.00000220604	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_17	0.00000257549	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_16	0.00000300378	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_15	0.00000349937	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_14	0.00000407165	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_13	0.00000473103	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_12	0.00000548896	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_11	0.00000635795	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_10	0.00000735162	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_9	0.00000848451	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_8	0.00000977139	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_7	0.00001122258	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_6	0.0000128609	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_5	0.0000146966	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_4	0.0000167694	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_3	0.0000191253	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_2	0.0000218083	3	(-4, 4)	act	0.40	dev (NA, NA)
Early_InitAge_1	0.0000248635	3	(-4, 4)	act	0.40	dev (NA, NA)
LnQ_base_WinterN(1)	-5.11324	1	(-20, 5)	OK	3.14	None
Q_power_WinterN(1)	-0.353898	3	(-5, 5)	OK	0.41	None
LnQ_base_WinterS(3)	-1.36603	1	(-20, 5)	OK	2.34	None
Q_power_WinterS(3)	-0.849303	3	(-5, 5)	OK	0.30	None
LnQ_base_TriEarly(5)	-0.803067	-1	(-15, 15)			None
Q_extraSD_TriEarly(5)	0.340847	5	(0.001, 2)	OK	0.15	None
LnQ_base_TriLate(6)	-0.254844	-1	(-15, 15)			None
Q_extraSD_TriLate(6)	0.236348	4	(0.001, 2)	OK	0.12	None
LnQ_base_NWFSC(7)	1.33092	-1	(-15, 15)			None
LnQ_base_WinterN(1)_BLK5add_2004	0.547816	3	(-0.99, 0.99)	OK	0.22	Normal (0, 0.5)
LnQ_base_WinterS(3)_BLK5add_2004	0.623155	3	(-0.99, 0.99)	OK	0.22	Normal (0, 0.5)
Size_DblN_peak_WinterN(1)	49.1595	1	(15, 75)	OK	0.92	None
Size_DblN_top_logit_WinterN(1)	3	-3	(-5, 3)			None
Size_DblN_ascend_se_WinterN(1)	4.28513	2	(-4, 12)	OK	0.13	None
Size_DblN_descend_se_WinterN(1)	14	-3	(-2, 15)			None
Size_DblN_start_logit_WinterN(1)	-999	-4	(-15, 5)			None
Size_DblN_end_logit_WinterN(1)	-999	-4	(-5, 5)			None
Retain_L_infl_WinterN(1)	24.2848	1	(10, 40)	OK	1.42	None
Retain_L_width_WinterN(1)	1.44101	2	(0.1, 10)	OK	0.19	None
Retain_L_asymptote_logit_WinterN(1)	9.99652	4	(-10, 10)	HI	0.11	None

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior	(Exp.Val, SD)
Retain_L_maleoffset_WinterN(1)	0	-2	(-10, 10)			None	
SzSel_Male_Peak_WinterN(1)	-11.5865	3	(-15, 15)	OK	0.78	None	
SzSel_Male_Ascend_WinterN(1)	-1.51838	4	(-15, 15)	OK	0.18	None	
SzSel_Male_Descend_WinterN(1)	0	-4	(-15, 15)			None	
SzSel_Male_Final_WinterN(1)	0	-4	(-15, 15)			None	
SzSel_Male_Scale_WinterN(1)	1	-4	(-15, 15)			None	
Size_DblN_peak_SummerN(2)	54.4414	1	(15, 75)	OK	1.27	None	
Size_DblN_top_logit_SummerN(2)	3	-3	(-5, 3)			None	
Size_DblN_ascend_se_SummerN(2)	5.43766	2	(-4, 12)	OK	0.10	None	
Size_DblN_descend_se_SummerN(2)	14	-3	(-2, 15)			None	
Size_DblN_start_logit_SummerN(2)	-999	-4	(-15, 5)			None	
Size_DblN_end_logit_SummerN(2)	-999	-4	(-5, 5)			None	
Retain_L_infl_SummerN(2)	30.2865	1	(10, 40)	OK	0.36	None	
Retain_L_width_SummerN(2)	1.30185	2	(0.1, 10)	OK	0.11	None	
Retain_L_asymptote_logit_SummerN(2)	5.74848	4	(-10, 10)	OK	0.35	None	
Retain_L_maleoffset_SummerN(2)	0	-2	(-10, 10)			None	
SzSel_Male_Peak_SummerN(2)	-14.0633	3	(-20, 15)	OK	0.97	None	
SzSel_Male_Ascend_SummerN(2)	-1.9821	4	(-15, 15)	OK	0.18	None	
SzSel_Male_Descend_SummerN(2)	0	-4	(-15, 15)			None	
SzSel_Male_Final_SummerN(2)	0	-4	(-15, 15)			None	
SzSel_Male_Scale_SummerN(2)	1	-4	(-15, 15)			None	
Size_DblN_peak_WinterS(3)	44.2878	1	(15, 75)	OK	1.74	None	
Size_DblN_top_logit_WinterS(3)	3	-3	(-5, 3)			None	
Size_DblN_ascend_se_WinterS(3)	4.4519	2	(-4, 12)	OK	0.27	None	
Size_DblN_descend_se_WinterS(3)	14	-3	(-2, 15)			None	
Size_DblN_start_logit_WinterS(3)	-999	-4	(-15, 5)			None	
Size_DblN_end_logit_WinterS(3)	-999	-4	(-5, 5)			None	
Retain_L_infl_WinterS(3)	28.9561	1	(10, 40)	OK	0.57	None	
Retain_L_width_WinterS(3)	1.44014	2	(0.1, 10)	OK	0.29	None	
Retain_L_asymptote_logit_WinterS(3)	4.86774	4	(-10, 10)	OK	0.94	None	
Retain_L_maleoffset_WinterS(3)	0	-2	(-10, 10)			None	
SzSel_Male_Peak_WinterS(3)	-12.2823	3	(-15, 15)	OK	1.77	None	
SzSel_Male_Ascend_WinterS(3)	-1.8526	4	(-15, 15)	OK	0.50	None	
SzSel_Male_Descend_WinterS(3)	0	-4	(-15, 15)			None	
SzSel_Male_Final_WinterS(3)	0	-4	(-15, 15)			None	
SzSel_Male_Scale_WinterS(3)	1	-4	(-15, 15)			None	
Size_DblN_peak_SummerS(4)	45.2206	1	(15, 75)	OK	1.29	None	
Size_DblN_top_logit_SummerS(4)	3	-3	(-5, 3)			None	
Size_DblN_ascend_se_SummerS(4)	4.89214	2	(-4, 12)	OK	0.15	None	

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior	(Exp.Val, SD)
Size_DblN_descend_se_SummerS(4)	14	-3	(-2, 15)			None	
Size_DblN_start_logit_SummerS(4)	-999	-4	(-15, 5)			None	
Size_DblN_end_logit_SummerS(4)	-999	-4	(-5, 5)			None	
Retain_L_infl_SummerS(4)	29.2437	1	(10, 40)	OK	0.33	None	
Retain_L_width_SummerS(4)	1.4909	2	(0.1, 10)	OK	0.12	None	
Retain_L_asymptote_logit_SummerS(4)	5.79317	4	(-10, 10)	OK	0.89	None	
Retain_L_maleoffset_SummerS(4)	0	-2	(-10, 10)			None	
SzSel_Male_Peak_SummerS(4)	-11.8309	3	(-15, 15)	OK	1.28	None	
SzSel_Male_Ascend_SummerS(4)	-1.83638	4	(-15, 15)	OK	0.29	None	
SzSel_Male_Descend_SummerS(4)	0	-4	(-15, 15)			None	
SzSel_Male_Final_SummerS(4)	0	-4	(-15, 15)			None	
SzSel_Male_Scale_SummerS(4)	1	-4	(-15, 15)			None	
Size_DblN_peak_TriEarly(5)	34.4179	1	(15, 61)	OK	1.10	None	
Size_DblN_top_logit_TriEarly(5)	3	-2	(-5, 3)			None	
Size_DblN_ascend_se_TriEarly(5)	4.05668	1	(-4, 12)	OK	0.20	None	
Size_DblN_descend_se_TriEarly(5)	14	-2	(-2, 15)			None	
Size_DblN_start_logit_TriEarly(5)	-999	-4	(-15, 5)			None	
Size_DblN_end_logit_TriEarly(5)	-999	-4	(-5, 5)			None	
SzSel_Male_Peak_TriEarly(5)	-3.30159	2	(-15, 15)	OK	1.07	None	
SzSel_Male_Ascend_TriEarly(5)	-0.488328	2	(-15, 15)	OK	0.25	None	
SzSel_Male_Descend_TriEarly(5)	0	-3	(-15, 15)			None	
SzSel_Male_Final_TriEarly(5)	0	-3	(-15, 15)			None	
SzSel_Male_Scale_TriEarly(5)	1	-4	(-15, 15)			None	
Size_DblN_peak_TriLate(6)	36.7376	1	(15, 61)	OK	0.90	None	
Size_DblN_top_logit_TriLate(6)	3	-2	(-5, 3)			None	
Size_DblN_ascend_se_TriLate(6)	4.65333	1	(-4, 12)	OK	0.11	None	
Size_DblN_descend_se_TriLate(6)	14	-2	(-2, 15)			None	
Size_DblN_start_logit_TriLate(6)	-999	-4	(-15, 5)			None	
Size_DblN_end_logit_TriLate(6)	-999	-4	(-5, 5)			None	
SzSel_Male_Peak_TriLate(6)	-2.59651	2	(-15, 15)	OK	0.92	None	
SzSel_Male_Ascend_TriLate(6)	-0.0842824	2	(-15, 15)	OK	0.14	None	
SzSel_Male_Descend_TriLate(6)	0	-3	(-15, 15)			None	
SzSel_Male_Final_TriLate(6)	0	-3	(-15, 15)			None	
SzSel_Male_Scale_TriLate(6)	1	-4	(-15, 15)			None	
Size_DblN_peak_NWFSC(7)	43.5751	1	(15, 61)	OK	0.83	None	
Size_DblN_top_logit_NWFSC(7)	3	-2	(-5, 3)			None	
Size_DblN_ascend_se_NWFSC(7)	5.18139	1	(-4, 12)	OK	0.07	None	
Size_DblN_descend_se_NWFSC(7)	14	-2	(-2, 15)			None	
Size_DblN_start_logit_NWFSC(7)	-999	-4	(-15, 5)			None	

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior	(Exp.Val, SD)
Size_DbLN_end_logit_NWFSC(7)	-999	-4	(-5, 5)			None	
SzSel_Male_Peak_NWFSC(7)	-5.93554	2	(-15, 15)	OK	0.71	None	
SzSel_Male_Ascend_NWFSC(7)	-0.502877	2	(-15, 15)	OK	0.08	None	
SzSel_Male_Descend_NWFSC(7)	0	-3	(-15, 15)			None	
SzSel_Male_Final_NWFSC(7)	0	-3	(-15, 15)			None	
SzSel_Male_Scale_NWFSC(7)	1	-4	(-15, 15)			None	
Size_DbLN_peak_WinterN(1)_BLK1add_1973	-0.201894	4	(-31.6, 28.4)	OK	0.88	Normal (0, 14.2)	
Size_DbLN_peak_WinterN(1)_BLK1add_1983	-1.23738	4	(-31.6, 28.4)	OK	0.81	Normal (0, 14.2)	
Size_DbLN_peak_WinterN(1)_BLK1add_1993	-1.32607	4	(-31.6, 28.4)	OK	0.63	Normal (0, 14.2)	
Size_DbLN_peak_WinterN(1)_BLK1add_2003	-0.388152	4	(-31.6, 28.4)	OK	0.46	Normal (0, 14.2)	
Size_DbLN_peak_WinterN(1)_BLK1add_2011	-0.119181	4	(-31.6, 28.4)	OK	0.48	Normal (0, 14.2)	
Retain_L_infl_WinterN(1)_BLK2add_2003	0.477145	4	(-16.19, 13.81)	OK	3.61	Normal (0, 6.905)	
Retain_L_infl_WinterN(1)_BLK2add_2010	5.78079	4	(-16.19, 13.81)	OK	2.87	Normal (0, 6.905)	
Retain_L_infl_WinterN(1)_BLK2add_2011	2.22632	4	(-16.19, 13.81)	OK	1.73	Normal (0, 6.905)	
Retain_L_width_WinterN(1)_BLK2add_2003	0.437634	4	(-1.601, 8.299)	OK	0.33	Normal (0, 0.8005)	
Retain_L_width_WinterN(1)_BLK2add_2010	0.571584	4	(-1.601, 8.299)	OK	0.74	Normal (0, 0.8005)	
Retain_L_width_WinterN(1)_BLK2add_2011	-0.510919	4	(-1.601, 8.299)	OK	0.20	Normal (0, 0.8005)	
Retain_L_asymptote_logit_WinterN(1)_BLK2repl_2003	7.37374	4	(-10, 10)	OK	1.70	None	
Retain_L_asymptote_logit_WinterN(1)_BLK2repl_2010	2.08428	4	(-10, 10)	OK	0.40	None	
Retain_L_asymptote_logit_WinterN(1)_BLK2repl_2011	9.02851	4	(-10, 10)	OK	1.02	None	
Size_DbLN_peak_SummerN(2)_BLK1add_1973	-2.75604	4	(-38.8, 21.2)	OK	0.85	Normal (0, 10.6)	
Size_DbLN_peak_SummerN(2)_BLK1add_1983	-4.93864	4	(-38.8, 21.2)	OK	1.12	Normal (0, 10.6)	
Size_DbLN_peak_SummerN(2)_BLK1add_1993	-5.7762	4	(-38.8, 21.2)	OK	1.16	Normal (0, 10.6)	
Size_DbLN_peak_SummerN(2)_BLK1add_2003	-4.12696	4	(-38.8, 21.2)	OK	0.67	Normal (0, 10.6)	
Size_DbLN_peak_SummerN(2)_BLK1add_2011	-1.53447	4	(-38.8, 21.2)	OK	0.62	Normal (0, 10.6)	
Retain_L_infl_SummerN(2)_BLK3add_2003	0.15738	4	(-20.679, 9.321)	OK	0.52	Normal (0, 4.6605)	
Retain_L_infl_SummerN(2)_BLK3add_2009	1.77908	4	(-20.679, 9.321)	OK	0.55	Normal (0, 4.6605)	
Retain_L_infl_SummerN(2)_BLK3add_2011	-1.06712	4	(-20.679, 9.321)	OK	0.51	Normal (0, 4.6605)	
Retain_L_width_SummerN(2)_BLK3add_2003	0.106445	4	(-1.0278, 8.8722)	OK	0.22	Normal (0, 0.5139)	
Retain_L_width_SummerN(2)_BLK3add_2009	0.142802	4	(-1.0278, 8.8722)	OK	0.23	Normal (0, 0.5139)	
Retain_L_width_SummerN(2)_BLK3add_2011	0.228908	4	(-1.0278, 8.8722)	OK	0.16	Normal (0, 0.5139)	
Retain_L_asymptote_logit_SummerN(2)_BLK3repl_2003	5.51708	4	(-10, 10)	OK	0.97	None	
Retain_L_asymptote_logit_SummerN(2)_BLK3repl_2009	6.86695	4	(-10, 10)	OK	6.44	None	
Retain_L_asymptote_logit_SummerN(2)_BLK3repl_2011	5.96198	4	(-10, 10)	OK	0.58	None	
Size_DbLN_peak_WinterS(3)_BLK1add_1973	-6.07356	4	(-25.422, 34.578)	OK	2.15	Normal (0, 12.711)	
Size_DbLN_peak_WinterS(3)_BLK1add_1983	-0.393954	4	(-25.422, 34.578)	OK	1.18	Normal (0, 12.711)	
Size_DbLN_peak_WinterS(3)_BLK1add_1993	2.96858	4	(-25.422, 34.578)	OK	1.66	Normal (0, 12.711)	
Size_DbLN_peak_WinterS(3)_BLK1add_2003	0.859894	4	(-25.422, 34.578)	OK	0.85	Normal (0, 12.711)	
Size_DbLN_peak_WinterS(3)_BLK1add_2011	1.91596	4	(-25.422, 34.578)	OK	0.99	Normal (0, 12.711)	

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp. Val, SD)
Retain_L.infl.WinterS(3).BLK2add.2003	-2.31652	4	(-18.816, 11.184)	OK	1.34	Normal (0, 5.592)
Retain_L.infl.WinterS(3).BLK2add.2010	1.40756	4	(-18.816, 11.184)	OK	1.65	Normal (0, 5.592)
Retain_L.infl.WinterS(3).BLK2add.2011	-3.95341	4	(-18.816, 11.184)	OK	2.49	Normal (0, 5.592)
Retain_L.width.WinterS(3).BLK2add.2003	0.34367	4	(-1.0443, 8.8557)	OK	0.37	Normal (0, 0.52215)
Retain_L.width.WinterS(3).BLK2add.2010	0.118141	4	(-1.0443, 8.8557)	OK	0.46	Normal (0, 0.52215)
Retain_L.width.WinterS(3).BLK2add.2011	-0.387615	4	(-1.0443, 8.8557)	OK	0.48	Normal (0, 0.52215)
Retain_L.asymptote.logit.WinterS(3).BLK2repl.2003	8.47752	4	(-10, 10)	OK	9.50	None
Retain_L.asymptote.logit.WinterS(3).BLK2repl.2010	5.69214	4	(-10, 10)	OK	8.51	None
Retain_L.asymptote.logit.WinterS(3).BLK2repl.2011	6.98104	4	(-10, 10)	OK	1.60	None
Size_DblN_peak_SummerS(4).BLK1add.1973	-8.19143	4	(-28.0793, 31.9207)	OK	2.13	Normal (0, 14.0397)
Size_DblN_peak_SummerS(4).BLK1add.1983	-8.6809	4	(-28.0793, 31.9207)	OK	2.90	Normal (0, 14.0397)
Size_DblN_peak_SummerS(4).BLK1add.1993	-0.797364	4	(-28.0793, 31.9207)	OK	1.39	Normal (0, 14.0397)
Size_DblN_peak_SummerS(4).BLK1add.2003	0.932051	4	(-28.0793, 31.9207)	OK	0.74	Normal (0, 14.0397)
Size_DblN_peak_SummerS(4).BLK1add.2011	0.680221	4	(-28.0793, 31.9207)	OK	0.85	Normal (0, 14.0397)
Retain_L.infl.SummerS(4).BLK3add.2003	-1.86903	4	(-19.055, 10.945)	OK	0.84	Normal (0, 5.4725)
Retain_L.infl.SummerS(4).BLK3add.2009	-2.00786	4	(-19.055, 10.945)	OK	1.14	Normal (0, 5.4725)
Retain_L.infl.SummerS(4).BLK3add.2011	-2.12954	4	(-19.055, 10.945)	OK	0.90	Normal (0, 5.4725)
Retain_L.width.SummerS(4).BLK3add.2003	0.305622	4	(-0.876, 9.024)	OK	0.22	Normal (0, 0.438)
Retain_L.width.SummerS(4).BLK3add.2009	0.176878	4	(-0.876, 9.024)	OK	0.26	Normal (0, 0.438)
Retain_L.width.SummerS(4).BLK3add.2011	0.0539534	4	(-0.876, 9.024)	OK	0.21	Normal (0, 0.438)
Retain_L.asymptote.logit.SummerS(4).BLK3repl.2003	9.13443	4	(-10, 10)	OK	12.97	None
Retain_L.asymptote.logit.SummerS(4).BLK3repl.2009	9.7475	4	(-10, 10)	OK	7.09	None
Retain_L.asymptote.logit.SummerS(4).BLK3repl.2011	9.9577	4	(-10, 10)	HI	1.31	None

Table 18: Results from 50 jitters from the base model.

Status	Base.Model	tab:jitter
Returned to base case		
Found local minimum		
Found better solution		
Total	50	

Table 19: Likelihood components from the base model

Likelihood Component	Value	tab:like
Total	1455.79	
Survey	-77.56	
Discard	-225.63	
Mean-body weight data	-158.25	
Length-frequency data	904.37	
Age-frequency data	1036.37	
Recruitment	-28.68	
Forecast Recruitment	0.03	
Parameter Priors	5.09	
Parameter Softbounds	0.04	

Table 20: Summary of reference points and management quantities for the base case.

Quantity	Estimate	~2.5%	~97.5%	tab:Ref_pts
	Confidence Interval	Confidence Interval		
Unfished spawning biomass (mt)	30554.7	24634.6	36474.8	
Unfished age 3+ biomass (mt)	49439.6	41597	57282.2	
Unfished recruitment (R0, thousands)	18626.7	12518.1	27716.1	
Spawning biomass(2019 mt)	9867.3	7682.4	12052.2	
Depletion (2019)	0.323	0.219	0.426	
Reference points based on SB_{40%}				
Proxy spawning biomass ($B_{25\%}$)	7638.7	6158.7	9118.7	
SPR resulting in $B_{25\%}$ ($SPR_{B25\%}$)	0.286	0.258	0.313	
Exploitation rate resulting in $B_{25\%}$	0.182	0.163	0.2	
Yield with $SPR_{B25\%}$ at $B_{25\%}$ (mt)	2830.3	2624.2	3036.4	
Reference points based on SPR proxy for MSY				
Spawning biomass	8096.3	6199.3	9993.3	
SPR_{proxy}				
Exploitation rate corresponding to SPR_{proxy}	0.173	0.145	0.2	
Yield with SPR_{proxy} at SB_{SPR} (mt)	2819.8	2590.1	3049.5	
Reference points based on estimated MSY values				
Spawning bioamss at MSY (SB_{MSY})	7005.9	5242.2	8769.6	
SPR_{MSY}				
Exploitation rate at MSY	0.195	0.164	0.225	
MSY (mt)	2835.9	2641.9	3029.9	

Table 21: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning biomass (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1876	49,440	30,555	48,800	1.00	18,627	1	0	0
1877	49,439	30,554	48,800	1.00	18,627	1	0	0
1878	49,438	30,554	48,799	1.00	18,627	1	0	0
1879	49,437	30,553	48,798	1.00	18,628	1	0	0
1880	49,437	30,553	48,797	1.00	18,628	12	0	0
1881	49,426	30,545	48,786	1.00	18,628	23	0	0
1882	49,406	30,532	48,766	1.00	18,627	33	0.003	0.001
1883	49,377	30,512	48,738	1.00	18,627	44	0.003	0.001
1884	49,341	30,487	48,701	1.00	18,626	55	0.003	0.001
1885	49,298	30,457	48,658	1.00	18,626	65	0.003	0.001
1886	49,248	30,422	48,608	1.00	18,625	76	0.006	0.002
1887	49,193	30,384	48,553	0.99	18,624	87	0.006	0.002
1888	49,132	30,342	48,493	0.99	18,623	98	0.006	0.002
1889	49,068	30,297	48,429	0.99	18,622	108	0.006	0.002
1890	49,000	30,248	48,360	0.99	18,621	119	0.006	0.002
1891	48,928	30,198	48,288	0.99	18,620	130	0.009	0.003
1892	48,853	30,145	48,214	0.99	18,619	141	0.009	0.003
1893	48,776	30,090	48,136	0.98	18,618	151	0.009	0.003
1894	48,696	30,034	48,057	0.98	18,617	162	0.009	0.003
1895	48,614	29,976	47,975	0.98	18,616	173	0.009	0.004
1896	48,531	29,916	47,892	0.98	18,615	184	0.012	0.004
1897	48,446	29,856	47,806	0.98	18,614	195	0.012	0.004
1898	48,359	29,794	47,720	0.98	18,613	205	0.012	0.004
1899	48,272	29,732	47,633	0.97	18,613	216	0.012	0.005
1900	48,183	29,669	47,544	0.97	18,612	227	0.012	0.005
1901	48,094	29,605	47,455	0.97	18,612	238	0.015	0.005
1902	48,004	29,540	47,365	0.97	18,611	248	0.015	0.005
1903	47,913	29,476	47,274	0.96	18,611	259	0.015	0.005
1904	47,822	29,410	47,183	0.96	18,612	270	0.015	0.006
1905	47,731	29,345	47,092	0.96	18,612	281	0.018	0.006
1906	47,639	29,279	47,000	0.96	18,613	291	0.018	0.006
1907	47,547	29,213	46,908	0.96	18,614	302	0.018	0.006
1908	47,455	29,146	46,816	0.95	18,616	313	0.018	0.007
1909	47,363	29,080	46,724	0.95	18,618	324	0.018	0.007
1910	47,271	29,014	46,632	0.95	18,621	334	0.021	0.007
1911	47,179	28,947	46,540	0.95	18,624	345	0.021	0.007
1912	47,088	28,881	46,448	0.95	18,628	356	0.021	0.008
1913	46,996	28,814	46,357	0.94	18,633	367	0.021	0.008
1914	46,906	28,748	46,266	0.94	18,638	377	0.024	0.008
1915	46,815	28,682	46,175	0.94	18,645	388	0.024	0.008
1916	46,725	28,617	46,085	0.94	18,652	394	0.024	0.009
1917	46,641	28,555	46,001	0.93	18,661	537	0.03	0.012
1918	46,433	28,409	45,792	0.93	18,668	432	0.027	0.009
1919	46,343	28,341	45,702	0.93	18,679	340	0.021	0.007
1920	46,352	28,341	45,711	0.93	18,694	235	0.015	0.005
1921	46,467	28,411	45,825	0.93	18,712	299	0.018	0.007
1922	46,518	28,441	45,876	0.93	18,731	433	0.027	0.009
1923	46,443	28,387	45,800	0.93	18,749	436	0.027	0.01
1924	46,374	28,336	45,730	0.93	18,768	543	0.033	0.012
1925	46,212	28,222	45,568	0.92	18,787	539	0.033	0.012
1926	46,072	28,120	45,427	0.92	18,808	532	0.033	0.012
1927	45,954	28,032	45,309	0.92	18,832	644	0.039	0.014

Table 21: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning biomass (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1928	45,747	27,884	45,101	0.91	18,856	632	0.036	0.014
1929	45,574	27,757	44,928	0.91	18,882	721	0.042	0.016
1930	45,340	27,588	44,692	0.90	18,910	673	0.039	0.015
1931	45,179	27,467	44,530	0.90	18,946	688	0.039	0.015
1932	45,029	27,350	44,379	0.90	18,989	822	0.048	0.019
1933	44,781	27,163	44,130	0.89	19,042	857	0.048	0.019
1934	44,539	26,973	43,887	0.88	19,131	1640	0.084	0.037
1935	43,600	26,312	42,946	0.86	19,259	1622	0.084	0.038
1936	42,770	25,713	42,112	0.84	19,445	1331	0.072	0.032
1937	42,310	25,351	41,648	0.83	19,678	1914	0.096	0.046
1938	41,384	24,665	40,715	0.81	19,869	2184	0.108	0.054
1939	40,324	23,878	39,647	0.78	19,860	2678	0.126	0.068
1940	38,938	22,855	38,257	0.75	19,434	2569	0.123	0.067
1941	37,821	22,000	37,143	0.72	18,623	2315	0.12	0.062
1942	37,080	21,408	36,420	0.70	17,727	3242	0.147	0.089
1943	35,577	20,317	34,944	0.66	17,085	3377	0.153	0.097
1944	34,051	19,271	33,447	0.63	17,050	2672	0.141	0.08
1945	33,239	18,772	32,653	0.61	17,582	2503	0.138	0.077
1946	32,584	18,413	31,995	0.60	17,963	3805	0.171	0.119
1947	30,712	17,244	30,106	0.56	17,751	3148	0.162	0.105
1948	29,524	16,484	28,909	0.54	17,503	4525	0.195	0.157
1949	27,115	14,878	26,508	0.49	17,257	4412	0.201	0.166
1950	24,942	13,412	24,343	0.44	17,160	4643	0.213	0.191
1951	22,705	11,885	22,113	0.39	17,164	3054	0.189	0.138
1952	22,093	11,425	21,503	0.37	17,273	2797	0.186	0.13
1953	21,790	11,202	21,201	0.37	16,951	2367	0.174	0.112
1954	21,912	11,295	21,322	0.37	16,181	2888	0.189	0.135
1955	21,541	11,088	20,965	0.36	15,105	2570	0.183	0.123
1956	21,451	11,063	20,904	0.36	14,068	2267	0.174	0.108
1957	21,592	11,219	21,082	0.37	13,386	2904	0.192	0.138
1958	21,048	10,988	20,570	0.36	13,872	2851	0.192	0.139
1959	20,457	10,756	19,992	0.35	17,137	2435	0.18	0.122
1960	20,192	10,709	19,690	0.35	21,658	2840	0.192	0.144
1961	19,576	10,334	18,959	0.34	16,766	3413	0.21	0.18
1962	18,561	9,536	17,856	0.31	11,393	3257	0.21	0.182
1963	17,840	8,807	17,301	0.29	12,855	3299	0.216	0.191
1964	17,126	8,186	16,721	0.27	18,499	2765	0.21	0.165
1965	16,893	8,096	16,414	0.26	15,563	2627	0.207	0.16
1966	16,798	8,220	16,173	0.27	32,710	2661	0.207	0.165
1967	16,768	8,233	16,123	0.27	14,915	2682	0.207	0.166
1968	16,972	8,087	15,975	0.26	15,421	2398	0.201	0.15
1969	17,638	8,085	17,122	0.26	15,854	2432	0.201	0.142
1970	18,321	8,242	17,788	0.27	16,417	3153	0.216	0.177
1971	18,290	8,336	17,742	0.27	15,835	3288	0.219	0.185
1972	18,038	8,541	17,480	0.28	13,025	3555	0.222	0.203
1973	17,407	8,459	16,885	0.28	10,902	3132	0.222	0.186
1974	16,897	8,390	16,463	0.27	13,807	3949	0.24	0.24
1975	15,362	7,679	14,967	0.25	14,283	3808	0.243	0.254
1976	13,780	6,899	13,302	0.23	16,377	3122	0.237	0.235
1977	12,768	6,368	12,264	0.21	15,187	2561	0.228	0.209
1978	12,345	6,021	11,794	0.20	11,138	3302	0.249	0.28
1979	11,294	5,226	10,802	0.17	11,032	3423	0.258	0.317

Table 21: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning biomass (million eggs)	Summary biomass 3+	Relative biomass	Age-0 recruits	Estimated total catch (mt)	1-SPR	Exploit. rate
1980	10,128	4,435	9,747	0.15	12,172	2855	0.255	0.293
1981	9,419	4,068	9,033	0.13	10,311	2765	0.258	0.306
1982	8,704	3,803	8,300	0.12	10,034	2770	0.258	0.334
1983	7,955	3,454	7,602	0.11	10,772	2454	0.258	0.323
1984	7,429	3,199	7,075	0.10	17,986	1911	0.249	0.27
1985	7,385	3,170	6,968	0.10	10,962	1871	0.249	0.269
1986	7,448	3,142	6,884	0.10	6,464	2133	0.255	0.31
1987	7,330	2,937	6,985	0.10	8,491	2563	0.267	0.367
1988	6,756	2,558	6,518	0.08	11,405	2360	0.267	0.362
1989	6,260	2,409	5,946	0.08	15,485	2304	0.267	0.387
1990	5,762	2,326	5,342	0.08	15,973	1961	0.264	0.367
1991	5,636	2,230	5,106	0.07	9,723	2133	0.27	0.418
1992	5,511	1,894	5,010	0.06	6,145	1807	0.267	0.361
1993	5,742	1,785	5,431	0.06	11,468	1681	0.264	0.309
1994	6,105	1,943	5,855	0.06	14,126	1544	0.255	0.264
1995	6,556	2,405	6,147	0.08	8,861	1674	0.252	0.272
1996	6,877	2,801	6,428	0.09	10,797	1929	0.252	0.3
1997	6,936	2,863	6,618	0.09	10,373	2051	0.255	0.31
1998	6,865	2,718	6,489	0.09	24,170	1739	0.249	0.268
1999	7,128	2,797	6,679	0.09	16,132	1621	0.243	0.243
2000	7,720	3,002	6,950	0.10	11,710	1915	0.249	0.275
2001	8,292	3,050	7,771	0.10	10,177	1980	0.249	0.255
2002	8,917	3,122	8,525	0.10	11,236	2067	0.249	0.242
2003	9,434	3,434	9,078	0.11	8,955	1795	0.234	0.198
2004	10,066	4,110	9,695	0.13	10,870	2294	0.24	0.237
2005	10,088	4,523	9,766	0.15	11,630	3015	0.252	0.309
2006	9,275	4,306	8,890	0.14	21,088	2215	0.24	0.249
2007	9,107	4,222	8,638	0.14	25,131	2404	0.243	0.278
2008	8,985	3,906	8,227	0.13	34,665	2180	0.24	0.265
2009	9,529	3,681	8,610	0.12	15,169	2329	0.246	0.271
2010	10,536	3,494	9,487	0.11	11,740	867	0.186	0.091
2011	13,224	4,414	12,727	0.14	13,434	791	0.165	0.062
2012	16,022	5,904	15,603	0.19	19,781	1159	0.171	0.074
2013	18,309	7,751	17,807	0.25	12,763	1973	0.189	0.111
2014	19,603	9,284	18,973	0.30	13,826	2398	0.189	0.126
2015	20,194	10,202	19,748	0.33	13,751	2727	0.192	0.138
2016	20,190	10,439	19,716	0.34	13,924	2543	0.186	0.129
2017	20,111	10,531	19,637	0.34	15,071	3050	0.195	0.155
2018	19,406	10,213	18,919	0.33	17,012	2882	0.195	0.152
2019	18,796	9,867	18,265	0.32	16,935	-	-	-

tab:Timeseries_mod1

Table 22: Sensitivity of the base model.

Label	Base	Low M	High M	Old M Prior	Fecundity	Sex Ratio	Francis	Dirichlet
Total Likelihood	1436.480	1436.480	1436.480	1436.480	1436.480	1436.480	1436.480	1436.480
Survey Likelihood	-77.916	-77.916	-77.916	-77.916	-77.916	-77.916	-77.916	-77.916
Discard Likelihood	-227.795	-227.795	-227.795	-227.795	-227.795	-227.795	-227.795	-227.795
Discard Mean Body Wt.	-159.144	-159.144	-159.144	-159.144	-159.144	-159.144	-159.144	-159.144
Length Likelihood	899.982	899.982	899.982	899.982	899.982	899.982	899.982	899.982
Age Likelihood	1023.080	1023.080	1023.080	1023.080	1023.080	1023.080	1023.080	1023.080
Recruitment Likelihood	-28.852	-28.852	-28.852	-28.852	-28.852	-28.852	-28.852	-28.852
Forecast Recruitment Likelihood	0.034	0.034	0.034	0.034	0.034	0.034	0.034	0.034
Parameter Priors Likelihood	7.054	7.054	7.054	7.054	7.054	7.054	7.054	7.054
log(R0)	9.938	9.938	9.938	9.938	9.938	9.938	9.938	9.938
SB Virgin	31106.300	31106.300	31106.300	31106.300	31106.300	31106.300	31106.300	31106.300
SB 2019	9907.260	9907.260	9907.260	9907.260	9907.260	9907.260	9907.260	9907.260
Depletion 2019	0.318	0.318	0.318	0.318	0.318	0.318	0.318	0.318
Total Yield - SPR 30	2990.590	2990.590	2990.590	2990.590	2990.590	2990.590	2990.590	2990.590
Steepness	0.814	0.814	0.814	0.814	0.814	0.814	0.814	0.814
Natural Mortality - Female	0.168	0.168	0.168	0.168	0.168	0.168	0.168	0.168
Length at Amin - Female	15.822	15.822	15.822	15.822	15.822	15.822	15.822	15.822
Length at Amax - Female	53.978	53.978	53.978	53.978	53.978	53.978	53.978	53.978
Von Bert. k - Female	0.136	0.136	0.136	0.136	0.136	0.136	0.136	0.136
SD young - Female	0.190	0.190	0.190	0.190	0.190	0.190	0.190	0.190
SD old - Female	0.024	0.024	0.024	0.024	0.024	0.024	0.024	0.024
Natural Mortality - Male	0.168	0.168	0.168	0.168	0.168	0.168	0.168	0.168
Length at Amin - Male	15.822	15.822	15.822	15.822	15.822	15.822	15.822	15.822
Length at Amax - Male	53.978	53.978	53.978	53.978	53.978	53.978	53.978	53.978
Von Bert. k - Male	0.136	0.136	0.136	0.136	0.136	0.136	0.136	0.136
SD young - Male	0.190	0.190	0.190	0.190	0.190	0.190	0.190	0.190
SD old - Male	0.024	0.024	0.024	0.024	0.024	0.024	0.024	0.024

Table 23: Data weights applied when using Francis data weighting in the base model. The data weights were acquired after a single model weighting iteration.

Fleet	Lengths	Ages	<code>tab:francis</code>
Winter North			
Summer North			
Winter South			
Summer South			
Triennial Early survey		-	
Triennial Late survey		-	
NWFSC shelf-slope survey			

Table 24: Data weights applied when using Dirichlet data weighting.

Fleet	Lengths	Ages	<code>tab:dirichlet</code>
Winter North			
Summer North			
Winter South			
Summer South			
Triennial Early survey		-	
Triennial Late survey		-	
NWFSC shelf-slope survey			

Table 25: Projection of potential OFL, spawning biomass, and depletion for the base case model. The removals in 2017 and 2018 were set at the defined management specification of XXX mt for each year assuming full attainment.

Year	OFL (mt)	ACL (mt)	Spawning Biomass	^{tab:Forecast_mod1} Depletion (%)
2019	4753	4340	5741	83.3
2020	4632	4229	5745	83.4
2021	4499	4108	5723	83.1
2022	4364	3984	5666	82.2
2023	4230	3862	5586	81.1
2024	4105	3748	5494	79.8
2025	3991	3644	5395	78.3
2026	3889	3551	5292	76.8
2027	3797	3467	5188	75.3
2028	3712	3389	5084	73.8

Table 26: Decision table summary of 10-year projections beginning in 2021 for alternate states of nature based on an axis of uncertainty for the base model. The removals in 2017 and 2018 were set at the defined management specification of 281 mt for each year assuming full attainment. Columns range over low, mid, and high states of nature over natural mortality, and rows range over different assumptions of catch levels. An entry of “–” indicates that the stock is driven to very low abundance under the particular scenario.

		tab:Decision_table_mod1_back					
		States of nature					
		M = 0.04725		M = 0.054		M = 0.0595	
		Year	Catch	Spawning Biomass	Depletion (%)	Spawning Biomass	Depletion (%)
ABC	2021						
	2022						
	2023						
	2024						
	2025						
	2026						
	2027						
	2028						
	2029						
	2030						
SPR target = 0.34	2021						
	2022						
	2023						
	2024						
	2025						
	2026						
	2027						
	2028						
	2029						
	2030						

10 Figures

figures




Figure 1: Total catches petrale sole. ^{fig:Catch}




Figure 2: Summary of data sources used in the base model. `fig:data_plot`




Figure 3: Map of the catch-per-unit-effort across by year for the NWFSC West Coast Groundfish Bottom Trawl Survey data. [fig:nw_map](#)




Figure 4: Catch-per-unit-effort (in \log_{space}) by latitude for the NWFSC West Coast Groundfish Bottom Trawl Survey data. fig:nw_cpue_lat




Figure 5: Catch-per-unit-effort (in log space) by depth for the NWFSC West Coast Groundfish Bottom Trawl Survey data. fig:nw_cpue_depth




Figure 6: QQ plot for the NWFSC West Coast Groundfish Bottom Trawl Survey data. `fig:nw_qq`




Figure 7: Pearson residuals for the encounter rate for the NWFSC West Coast Groundfish Bottom Trawl Survey by VAST. fig:nw_enc_76




Figure 8: Pearson residuals for the estimated catch rate for the NWFSC West Coast Groundfish Bottom Trawl Survey by VAST. [?77g:nwCatchRate](#)

NWFSC West Coast Groundfish Bottom Trawl Survey




Figure 9: Estimated index of abundance from the NWFSC West Coast Groundfish Bottom Trawl Survey data compared to the design-based index and the index from the 2015 update assessment. [fig:nw_index](#)




Figure 10: Estimated density of abundance from the NWFSC West Coast Groundfish Bottom Trawl Survey data by VAST. [fig:nw_density79](#)





Figure 11: Length frequency by sex for the NWFSC West Coast Groundfish Bottom Trawl Survey data. [fig:nw_len_freq](#)

Female



Male




Figure 12: Age frequency by sex for the NWFSC West Coast Groundfish Bottom Trawl Survey data. [fig:nw_age_freq](#)





Figure 13: Map of the catch-per-unit-effort across by year for the Triennial Survey data. [fig:tri_map](#)

Early



Late





Figure 14: Catch-per-unit-effort (in log space) by latitude for the Triennial Survey data. `fig:tri_cpue_l`

Early



Late




Figure 15: Catch-per-unit-effort (in log space) by depth (m) for the Triennial Survey data. `fig:tri_cpue`




Figure 16: Length (cm) by depth (m) for the Triennial Survey data. `fig:tri_size_depth`




Figure 17: QQ plot for the Triennial Early Survey data. fig:tri_early_qq




Figure 18: Pearson residuals for the encounter rate for the Triennial Early Survey by VAST. fig:tri_early





Figure 19: Pearson residuals for the estimated catch rate for the Triennial Early Survey by VAST. | [fig:tri_early_catch_rate](#) 88






Figure 21: Pearson residuals for the encounter rate for the Triennial Late Survey by VAST. fig:tri_late_90




Figure 22: Pearson residuals for the estimated catch rate for the Triennial Late Survey by VAST. | [fig:tri_late_catch_rate](#) 91

AFSC/NWFSC West Coast Triennial Shelf Survey




Figure 23: Estimated index of abundance from the Triennial Survey data compared to the design-based index and the index from the 2015 update assessment. [fig:tri_index](#)




Figure 24: Estimated density of abundance from the Triennial Early Survey data by VAST. fig:tri_early
93





Figure 25: Estimated density of abundance from the Triennial Late Survey data by VAST. fig:tri_late_94

Female



Male





Figure 26: Length frequency by sex for the Triennial Early Survey data. [fig:tri_early_len_freq](#)

Female



Male




Figure 27: Length frequency by sex for the Triennial Late Survey data. `fig:tri_late_len_freq`




Figure 28: Northern winter fishery mean body weights of discarded fish for petrale sole. fig:nw_bodywt




Figure 29: Northern summer fishery mean body weights of discarded fish for petrale sole. fig:ns_bodywt




Figure 30: Southern winter fishery mean body weights of discarded fish for petrale sole. `fig:sw_bodywt`




Figure 31: Southern summer fishery mean body weights of discarded fish for petrale sole. `fig:ss_bodywt`




Figure 32: Northern fishery, winter and summer, retained and discarded length frequency distributions for petrale sole. [fig:north_lengths](#)




Figure 33: Southern fishery, winter and summer, retained and discarded length frequency distributions for petrale sole. [fig:south_lengths](#)




Figure 34: Commercial fishery age frequency distributions for petrale sole. `fig:comm_ages`




Figure 35: The Northern Winter fishery catch-per-unit-effort based on logbook data for petrale sole. [fig:north_cpue](#)




Figure 36: The Southern Winter fishery catch-per-unit-effort based on logbook data for petrale sole. | [fig:south_cpue](#)




Figure 37: Estimated maturity-at-length for petrale sole. fig:maturity




Figure 38: Fecundity-at-length assumed in the model for petrale sole. [fig:fecundity_model](#)




Figure 39: Estimated fecundity-at-length for petrale sole based on Lefebvre et al. (in press). fig:fecundity

NWFSC Shelf-Slope




Figure 40: Estimated proportion of female fish collected by the NWFSC shelf-slope survey across all years for petrale sole. [fig:sex_ratio](#)




Figure 41: Estimated weight-at-length for female and male petrale sole. [fig:wt_length](#)




Figure 42: Length-at-age across data sources for female and male petrale sole. fig:length_age




Figure 43: Prior distribution for natural mortality for female and male petrale sole. `fig:m_prior`




Figure 44: Prior distribution for steepness petrale sole. `fig:h_prior`




Figure 45: Comparison of model bridging estimates from Stock Synthesis version 3.30.13 and 3.24U for petrale sole for the 2015 assessment. fig:bridge




Figure 46: Estimated length-at-age for male and female for petrale sole with estimated CV. fig:sizeatage




Figure 47: Estimated spawning output-at-length for female petrale sole. `fig:spawnoutlen`




Figure 48: Estimated selectivity for each commerical fleet over the assessment period for female and male petrale sole. [fig:fish_selex](#)




Figure 49: Estimated retention for each commercial fleet over the assessment period for petrale sole. Retention was not estimated to be sex-specific. [fig:fish_reten](#)




Figure 50: Estimated selectivity for each survey over the assessment period for female and male petrale sole. [fig:survey_selex](#)

Age-0 recruits (1,000s) with ~95% asymptotic intervals




Figure 51: Estimated time-series of recruitment for petrale sole. `fig:recruits`




Figure 52: Estimated time-series of recruitment deviations for petrale sole. `fig:recdevs`




Figure 53: Recruitment bias adjustment in the model. `fig:bias_adjust`




Figure 54: Fit to the Winter North catch-per-unit-effort time series for petrale sole. `fig:fit_wn_cpue`




Figure 55: Catchability to the Winter North catch-per-unit-effort time series. `fig:q_north`




Figure 56: Fit to the Winter South catch-per-unit-effort time series for petrale sole. `fig:fit_ws_cpue`




Figure 57: Catchability to the Winter South catch-per-unit-effort time series. `fig:q_south`




Figure 58: Fit to the Triennial Survey Early time series for petrale sole. [fig:fit_tri_early](#)




Figure 59: Fit to the Triennial Survey Late time series for petrale sole. `fig:fit_tri_late`




Figure 60: Fit to the NWFSC West Coast Groundfish Bottom Trawl Survey time series for petrale sole. [fig:fit_nwfsc_survey](#)




Figure 61: Fit to the discard rates for the Winter North fleet for petrale sole. `fig:fit_wn_discard`




Figure 62: Fit to the discard rates for the Summer North fleet for petrale sole. [`fig:fit_sn_discard`](#)




Figure 63: Fit to the discard rates for the Winter South fleet for petrale sole. `fig:fit_ws_discard`




Figure 64: Fit to the discard rates for the Summer South fleet for petrale sole. fig:fit_ss_discard




Figure 65: Discard rates by fleet for petrale sole. fig:Discard




Figure 66: Fit to the Northern winter fishery mean body weights of discarded fish for petrale sole. [fig:nw_bodywt_fit](#)




Figure 67: Fit to the Northern summer fishery mean body weights of discarded fish for petrale sole. [fig:ns_bodywt_fit](#)




Figure 68: Fit to the Southern winter fishery mean body weights of discarded fish for petrale sole. [fig:sw_bodywt_fit](#)




Figure 69: Fit to the Southern summer fishery mean body weights of discarded fish for petrale sole. [fig:ss_bodywt_fit](#)




Figure 70: Length compositions aggregated across time by fleet. Labels ‘retained’ and ‘discard’ indicate retained or discarded samples for each fleet. Panels without this designation represent the whole catch. | fig: length_agg



Figure 71: Pearson residuals, discard, Winter (N) (max=7.19)
 Closed bubbles are positive residuals (observed \geq expected) and open bubbles are negative residuals (observed $<$ expected). [fig:discard_wn_len_pearson](#)




Figure 72: Pearson residuals, discard, Summer (N) (max=6.6)
 Closed bubbles are positive residuals (observed \geq expected) and open bubbles are negative residuals (observed $<$ expected). [fig:discard_sn_len_pearson](#)




Figure 73: Pearson residuals, discard, Winter (S) (max=3.81)

Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:discard_ws_len_pearson](#)




Figure 74: Pearson residuals, discard, Summer (S) (max=5.02)

Closed bubbles are positive residuals ($\text{observed} > \text{expected}$) and open bubbles are negative residuals ($\text{observed} < \text{expected}$). [fig:discard_ss_len_pearson](#)




Figure 75: Pearson residuals, retained, Winter (N) (max=3.93) (plot 4 of 4)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:wn_len_pearson](#)




Figure 76: Pearson residuals, retained, Summer (N) (max=5.15) (plot 4 of 4)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:sn_len_pearson](#)




Figure 77: Pearson residuals, retained, Winter (S) ($\max=5.1$) (plot 3 of 3)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:ws_len_pearson](#)




Figure 78: Pearson residuals, retained, Summer (S) (max=5.63) (plot 4 of 4)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:ss_len_pearson](#)




Figure 79: Pearson residuals, whole catch, Triennial - Early (max=3.16)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:tri_early_len_pearson](#)




Figure 80: Pearson residuals, whole catch, Triennial _ Late (max=3.54)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected). [fig:tri_late_1en_pearson](#)




Figure 81: Pearson residuals, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (max=5.11)

Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected). [fig:nwfsc_combo_len_pearson](#)




Figure 82: McAllister and Ianelli (harmonic mean) weighting for the Winter North fishery length data. [fig:harm_mean_wn](#)




Figure 83: McAllister and Ianelli (harmonic mean) weighting for the Summer North fishery length data. | [fig:harm_mean_sn](#)




Figure 84: McAllister and Ianelli (harmonic mean) weighting for the Winter South fishery length data. [fig:harm_mean_ws](#)




Figure 85: McAllister and Ianelli (harmonic mean) weighting for the Summer South fishery length data. [fig:harm_mean_wn](#)




Figure 86: McAllister and Ianelli (harmonic mean) weighting for the Triennial Survey - early length data. [fig:harm_mean_tri_early](#)




Figure 87: McAllister and Ianelli (harmonic mean) weighting for the Triennial Survey - late length data. | [fig:harm_mean_tri_late](#)




Figure 88: McAllister and Ianelli (harmonic mean) weighting for the NWFSC West Coast Groundfish Bottom Trawl Survey length data. [fig:harm_mean_nwfsc](#)




Figure 89: Age compositions aggregated across time for each fishery fleet. `fig:age_agg`




Figure 90: Pearson residuals, retained, Winter (N) (max=4.57) (plot 5 of 5)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:wn_age_pearson](#)




Figure 91: Pearson residuals, retained, Summer (N) (max=5.9) (plot 5 of 5)
 Closed bubbles are positive residuals ($\text{observed} > \text{expected}$) and open bubbles are negative residuals ($\text{observed} < \text{expected}$). [fig:sn_age_pearson](#)




Figure 92: Pearson residuals, retained, Winter (S) ($\max=7.4$) (plot 3 of 3)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:ws_age_pearson](#)




Figure 93: Pearson residuals, retained, Summer (S) (max=4.31) (plot 3 of 3)
 Closed bubbles are positive residuals (observed $>$ expected) and open bubbles are negative residuals (observed $<$ expected). [fig:ss_age_pearson](#)




Figure 94: Conditional AAL plot, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (plot 1 of 4) These plots show mean age and std. dev. in conditional AAL. Left plots are mean AAL by size_class (obs. and pred.) with 90% CIs based on adding 1.64 SE of mean to the data. Right plots in each pair are SE of mean AAL (obs. and pred.) with 90% CIs based on the chi_square distribution. [fig:nwfsc_combo_andre_1](#)




Figure 95: Conditional AAL plot, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (plot 2 of 4) [fig:nwfsc_combo_andre_2](#)




Figure 96: Conditional AAL plot, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (plot 3 of 4) [fig:nwfsc_combo_andre_3](#)




Figure 97: Conditional AAL plot, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (plot 4 of 4) | [fig:nwfsc_combo_andre_4](#)




Figure 98: Pearson residuals, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (max=7.72) (plot 1 of 2) [fig:nwfsc_combo_pearson_1](#)




Figure 99: Pearson residuals, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey (max=7.72) (plot 1 of 2) (plot 2 of 2) | [fig:nwfsc_combo_pearson_2](#)




Figure 100: McAllister and Ianelli (harmonic mean) weighting for the Winter North fishery age data. [fig:harm_mean_wn_age](#)




Figure 101: McAllister and Ianelli (harmonic mean) weighting for the Summer North fishery age data. [fig:harm_mean_sn_age](#)




Figure 102: McAllister and Ianelli (harmonic mean) weighting for the Winter South fishery age data.




Figure 103: McAllister and Ianelli (harmonic mean) weighting for the Summer South fishery age data.




Figure 104: Estimated time-series of spawning biomass trajectory (circles and line: median; light broken lines: 95% credibility intervals) for petrale sole. fig:ssb




Figure 105: Estimated time-series of total biomass for petrale sole. ^{fig:total_bio}

Spawning depletion with ~95% asymptotic intervals




Figure 106: Estimated time-series of relative spawning biomass (depletion) (circles and line: median; light broken lines: 95% credibility intervals) for petrale sole. [fig:dep1](#)




Figure 107: Estimated recruitment (colored circles) and the assumed stock-recruit relationship (solid black line). The dashed line shows the effect of the bias correction for the lognormal distribution. [fig:stock_recruit_curve](#)




Figure 108: Estimate spawning biomass for the base model and each sensitivity. `fig:sens_ssb`




Figure 109: Estimate relative spawning biomass for the base model and each sensitivity. fig:sens_depl




Figure 110: Retrospective pattern for spawning biomass. [fig:retro_ssbb](#)




Figure 111: Retrospective pattern for relative spawning biomass. [fig:retro_depl](#)




Figure 112: Retrospective pattern for estimated recruitment deviations. `fig:retro_recdev`




Figure 113: Pattern for estimated spawning biomass from each assessment since 2005. fig:historical




Figure 114: Likelihood profile across steepness values. fig:piner_h




Figure 115: Trajectories of relative spawning biomass across values of steepness. `fig:h_trajectory`




Figure 116: Likelihood profile across female natural mortality values. Male natural mortality was estimated. [fig:m_like](#)




Figure 117: Trajectories of relative spawning biomass across values of natural mortality. fig:m_trajectories




Figure 118: Likelihood profile across R_0 values. [fig:piner_R0](#)




Figure 119: Estimated relative spawning potential ratio 1-SPR for the base model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the SPR30% harvest rate. The last year in the time-series is 2018. [fig:SPR_all_fig](#)




Figure 120: Equilibrium yield curve for the base case model. Values are based on the 2018 fishery selectivity and with steepness estimated at 0.84. fig:yield

11 Appendix A. Detailed Fit to Length Composition Data

appendix-a.-detailed-fit-to-length-composition-data




Figure 121: Length comps, retained, Winter (N) (plot 1 of 4). 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Iannelli tuning method.




Figure 122: Length comps, retained, Winter (N) (plot 2 of 4) `fig:length.fits`




Figure 123: Length comps, retained, Winter (N) (plot 3 of 4) `fig:length.fits`




Figure 124: Length comps, retained, Winter (N) (plot 4 of 4) `fig:length.fits`




Figure 125: Length comps, discard, Winter (N). ‘N adj.’ is the input sample size after data_weighting adjustment. N_eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. | [fig:length_fits](#)




Figure 126: Length comps, retained, Summer (N) (plot 1 of 4). 'N adj.' is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:length_fits](#)




Figure 127: Length comps, retained, Summer (N) (plot 2 of 4) `fig:length_fits`




Figure 128: Length comps, retained, Summer (N) (plot 3 of 4) `fig:length_fits`




Figure 129: Length comps, retained, Summer (N) (plot 4 of 4) `fig:length_fits`




Figure 130: Length comps, discard, Summer (N). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Iannelli tuning method. [fig:length_fits](#)




Figure 131: Length comps, retained, Winter (S) (plot 1 of 3). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:length_fits](#)




Figure 132: Length comps, retained, Winter (S) (plot 2 of 3) `fig:length.fits`




Figure 133: Length comps, retained, Winter (S) (plot 3 of 3) `fig:length.fits`




Figure 134: Length comps, discard, Winter (S). ‘N adj.’ is the input sample size after data_weighting adjustment. N_eff_ is the calculated effective sample size used in the McAllister_Iannelli tuning method. | [fig:length_fits](#)




Figure 135: Length comps, retained, Summer (S) (plot 1 of 4). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:length_fits](#)




Figure 136: Length comps, retained, Summer (S) (plot 2 of 4) `fig:length.fits`





Figure 137: Length comps, retained, Summer (S) (plot 3 of 4) `fig:length.fits`

Proportion



Length (cm)

Figure 138: Length comps, retained, Summer (S) (plot 4 of 4) `fig:length_fits`




Figure 139: Length comps, discard, Summer (S). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister Iannelli tuning method. [fig:length_fits](#)




Figure 140: Length comps, whole catch, Triennial _ Early. 'N adj.' is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:length_fits](#)




Figure 141: Length comps, whole catch, Triennial _ Late. 'N adj.' is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister Iannelli tuning method. [fig:length_fits](#)




Figure 142: Length comps, whole catch, NWFSC West Coast Groundfish Bottom Trawl Survey. ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:length_fits](#)

12 Appendix B. Detailed Fit to Age Composition Data

appendix-b.-detailed-fit-to-age-composition-data




Figure 143: Age comps, retained, Winter (N) (plot 1 of 5). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister Iannelli tuning method. [fig:age_fits](#)




Figure 144: Age comps, retained, Winter (N) (plot 2 of 5) `fig:age_fits`




Figure 145: Age comps, retained, Winter (N) (plot 3 of 5) `fig:age_fits`




Figure 146: Age comps, retained, Winter (N) (plot 4 of 5) `fig:age_fits`




Figure 147: Age comps, retained, Winter (N) (plot 5 of 5) `fig:age_fits`




Figure 148: Age comps, retained, Summer (N) (plot 1 of 5). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister Iannelli tuning method. [fig:age_fits](#)




Figure 149: Age comps, retained, Summer (N) (plot 2 of 5) `fig:age.fits`




Figure 150: Age comps, retained, Summer (N) (plot 3 of 5) `fig:age.fits`




Figure 151: Age comps, retained, Summer (N) (plot 4 of 5) `fig:age.fits`




Figure 152: Age comps, retained, Summer (N) (plot 5 of 5) `fig:age.fits`




Figure 153: Age comps, retained, Winter (S) (plot 1 of 3). 'N adj.' is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method. [fig:age_fits](#)




Figure 154: Age comps, retained, Winter (S) (plot 2 of 3) [fig:age_fits](#)




Figure 155: Age comps, retained, Winter (S) (plot 3 of 3) [fig:age_fits](#)




Figure 156: Age comps, retained, Summer (S) (plot 1 of 3). ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Iannelli tuning method. [fig:age_fits](#)




Figure 157: Age comps, retained, Summer (S) (plot 2 of 3) `fig:age.fits`




Figure 158: Age comps, retained, Summer (S) (plot 3 of 3) `fig:age_fits`

13 Appendix C. List of Auxiliary Files Available

[appendix-c.-list-of-auxiliary-files-available](#)

The listed files are also available as auxiliary files to accompany the assessment document:

1. Numbers at age for female and male petrale sole (Petrale natagef.csv and Petrale natagem.csv)
2. The petrale sole Stock Synthesis 3.30.13 model files
 - (a) 2019petrale.dat
 - (b) 2019petrale.ctl
 - (c) forecast.ss
 - (d) starter.ss

->