



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Social Sciences - Papers

Faculty of Social Sciences

2013

Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: A cohort study

Jiehui Yin

University of Sydney

Glenn P. Salkeld

University of Wollongong, gsalkeld@uow.edu.au

Stephen Lambert

University of Queensland

Alexa Dierig

Children's Hospital Westmead

Leon Heron

University of Sydney

See next page for additional authors

Publication Details

Yin, J. K., Salkeld, G., Lambert, S. B., Dierig, A., Heron, L., Leask, J., Yui Kwan Chow, M. & Booy, R. (2013). Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: A cohort study. *Influenza and Other Respiratory Viruses*, 7 (6), 1103-1112.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: A cohort study

Abstract

Background

Influenza and other respiratory infections cause excess winter morbidity in children. This study assessed the economic impact of influenza-like illness (ILI) on families with children attending childcare using a societal perspective.

Methods

We conducted a prospective cohort study in 90 childcare centres and one general practitioner clinics in Sydney, Australia, during 2010. Healthy children aged ≥ 6 months to < 3 years were enrolled. Economic impacts of ILI (temperature $\geq 37.8^{\circ}\text{C}$ or parental report of fever, plus ≥ 1 respiratory symptoms) were collected at 2 and 4 weeks after ILI onset by telephone interview. Parent-collected respiratory specimens were tested for respiratory viruses using real-time PCR (RT-PCR). Costs associated with healthcare visits, medication usage, carer time lost (work or recreation) and home care and/or additional childcare were collected. Influenza-like illness costs were described and further analysed using a Tobit model. Zero-inflated Poisson regression was employed to compare the numbers of healthcare visits for each ILI.

Results

Of 381 children enrolled and analysed, 105 developed 124 ILIs. Specimens were available for 117 ILIs: five were positive by RT-PCR for A(H1N1)pdm09, 39 for adenovirus, 39 for rhinovirus, 15 for a coronavirus and 27 for a polyomavirus. The mean cost of all ILIs was AU\$626 (95% confidence interval: AU\$484–768) per ILI with no significant differences observed between viruses. Carers lost on average 13 hours of work and 3 hours of leisure time per ILI. Independent drivers of ILI costs were having both parents in employed work and longer duration of ILI. In multivariate analyses, four variables were significantly associated with an increased number of healthcare visits per ILI: non-Caucasian child, living in a detached house, both parents in employed work and having an ILI with one or more viruses identified.

Conclusions

For families with a child attending childcare, ILIs cause a substantial economic burden. An ILI in a child with working parents and/or with longer duration appears to cost more in monetary terms. Healthcare visits were increased if the child was non-Caucasian, lived in a detached house, had working parents or had a virus-positive ILI. Our findings on the estimates and determinants of economic impacts from respiratory virus infection highlight the importance and feasibility of an interdisciplinary (epidemiology/health economics) approach to such research.

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Yin, J. K., Salkeld, G., Lambert, S. B., Dierig, A., Heron, L., Leask, J., Yui Kwan Chow, M. & Booy, R. (2013). Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: A cohort study. *Influenza and Other Respiratory Viruses*, 7 (6), 1103-1112.

Authors

Jiehui Yin, Glenn P. Salkeld, Stephen Lambert, Alexa Dierig, Leon Heron, Julie Leask, Maria Yui, and Robert Booy

Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: a cohort study

Jiehui Kevin Yin,^{a,b} Glenn Salkeld,^c Stephen B. Lambert,^d Alexa Dierig,^{a,e} Leon Heron,^{a,b,f} Julie Leask,^{a,c,f} Maria Yui Kwan Chow,^{a,b} Robert Booy^{a,b,f}

^aNational Centre for Immunisation Research and Surveillance, The Children's Hospital at Westmead, Sydney, NSW, Australia. ^bDiscipline of Paediatrics and Child Health, Faculty of Medicine, The University of Sydney, Sydney, NSW, Australia. ^cSydney School of Public Health, Faculty of Medicine, The University of Sydney, Sydney, NSW, Australia. ^dQueensland Children's Medical Research Institute, The University of Queensland and Queensland Children's Health Services, and Queensland Health Immunisation Program, Brisbane, Qld, Australia. ^eChildren's Hospital Klinikum Dritter Orden, Munich, Germany. ^fSydney Emerging Infections and Biosecurity Institute, Sydney, NSW, Australia.

Correspondence: Jiehui Kevin Yin, National Centre for Immunisation Research and Surveillance, The Children's Hospital at Westmead, Sydney, NSW 2145, Australia.

E-mail: kevin.yin@health.nsw.gov.au

Accepted 30 May 2013. Published Online 06 July 2013.

Background Influenza and other respiratory infections cause excess winter morbidity in children. This study assessed the economic impact of influenza-like illness (ILI) on families with children attending childcare using a societal perspective.

Methods We conducted a prospective cohort study in 90 childcare centres and one general practitioner clinic in Sydney, Australia, during 2010. Healthy children aged ≥ 6 months to < 3 years were enrolled. Economic impacts of ILI (temperature $\geq 37.8^{\circ}\text{C}$ or parental report of fever, plus ≥ 1 respiratory symptoms) were collected at 2 and 4 weeks after ILI onset by telephone interview. Parent-collected respiratory specimens were tested for respiratory viruses using real-time PCR (RT-PCR). Costs associated with healthcare visits, medication usage, carer time lost (work or recreation) and home care and/or additional childcare were collected. Influenza-like illness costs were described and further analysed using a Tobit model. Zero-inflated Poisson regression was employed to compare the numbers of healthcare visits for each ILI.

Results Of 381 children enrolled and analysed, 105 developed 124 ILIs. Specimens were available for 117 ILIs: five were positive by RT-PCR for A(H1N1)pdm09, 39 for adenovirus, 39 for rhinovirus, 15

for a coronavirus and 27 for a polyomavirus. The mean cost of all ILIs was AU\$626 (95% confidence interval: AU\$484–768) per ILI with no significant differences observed between viruses. Carers lost on average 13 hours of work and 3 hours of leisure time per ILI. Independent drivers of ILI costs were having both parents in employed work and longer duration of ILI. In multivariate analyses, four variables were significantly associated with an increased number of healthcare visits per ILI: non-Caucasian child, living in a detached house, both parents in employed work and having an ILI with one or more viruses identified.

Conclusions For families with a child attending childcare, ILIs cause a substantial economic burden. An ILI in a child with working parents and/or with longer duration appears to cost more in monetary terms. Healthcare visits were increased if the child was non-Caucasian, lived in a detached house, had working parents or had a virus-positive ILI. Our findings on the estimates and determinants of economic impacts from respiratory virus infection highlight the importance and feasibility of an interdisciplinary (epidemiology/health economics) approach to such research.

Please cite this paper as: Yin *et al.* (2013) Estimates and determinants of economic impacts from influenza-like illnesses caused by respiratory viruses in Australian children attending childcare: a cohort study. *Influenza and Other Respiratory Viruses* 7(6), 1103–1112.

Introduction

Respiratory virus infection causes significant morbidity and mortality in the paediatric population.^{1–3} It also results in substantial economic burden in terms of healthcare visits, including general practitioner (GP) and specialist consultations, medications, work absenteeism (either to care for a

sick child or due to secondary illness in a carer) and hospitalisations.^{4,5} Attending childcare is likely to increase the risk of respiratory infection and costs associated with respiratory viruses: by providing an opportunity for respiratory virus transmission between children and because carers are required to miss work or leisure activities to either care for a sick child or recuperate from their own illness due

to intrahousehold transmission. According to the Australian Bureau of Statistics, 50% of children aged three attend childcare, with an average attendance of 17 hours per week, and 13% of children attending 35 hours or more a week.⁶ According to the National Health and Medical Research Council guidelines in Australia,⁷ children with influenza-like illness (ILI) need to be excluded from childcare until they have recovered.

To provide accurate and timely information about the economic impacts of respiratory virus infections in childcare, we present here the estimates and determinants of virus-specific costs from respiratory infections using a societal perspective and based on the findings from a prospective cohort study.

Methods

The study cohort

We collected ILI impact data in a prospective cohort study in the southern hemisphere winter season in 2010. From 24 January 2010, healthy children aged ≥ 6 months to < 3 years (as at 01 March 2010) were recruited from 90 childcare centres and one paediatric-focussed GP clinic in metropolitan Sydney, Australia. Exclusion criteria included any known allergies to influenza vaccine, history of Guillain-Barré syndrome and haematological disorders. Informed consent for participation was obtained from parents or legal guardians.

Initial contact for recruitment took place through participating childcares, and following receipt of the information and consent document by mail, families interested in participating were screened and enrolled by telephone.

Detailed household demographic data and risk factors were obtained at enrolment, including household income, parents' education and occupations, and the child's medical and medication history.

Ethical approval was given by the Human Research Ethics Committee of the Children's Hospital at Westmead, Sydney, Australia.

Disease surveillance and specimen collection

Following increased reports of ILI activity in the local community, we started active surveillance from 30 July 2010; we ceased surveillance on 31 October 2010. During this period, all participating families received weekly reminders (via email, text message or telephone call) from the study team asking whether their child had developed an episode of ILI (temperature $\geq 37.8^{\circ}\text{C}$ or carer report of child feeling feverish and at least one of the following symptoms: cough, runny nose, nasal congestion or sore throat).^{8,9} When an ILI occurred, parents/guardians were asked to collect nose and/or throat swabs (Copan Italia S. p. A, Brescia, Italy) and mail these to the Queensland Paediatric Infectious Diseases

(QPID) Laboratory for quantitative real-time multiplex PCR (RT-PCR) assays testing of 19 respiratory viruses, as described previously.^{10–12}

Follow-up telephone interview

We conducted telephone interviews with a carer of a child with an ILI 10–16 days after the illness onset (further call at 26–32 days if the ILI still ongoing at 14 days) and collected data about the illness, including the following:

- Healthcare visits
 - GP, pharmacy, emergency department (ED) and specialist room (SR) visits and hospitalisations,
 - Related costs: public transport (including taxi use), distance (in kilometres) travelled using private vehicle, parking fees and other non-medication costs.
- Carer time lost: time was captured in hours and minutes and classified in time away from work or recreation
 - Seeking healthcare for the ILI episode,
 - Looking after the sick child at home (in excess of usual time spent caring for the child),
 - Secondary illness in a carer (occurring after illness onsets of the study child).
- Costs of home care, additional childcare for other healthy children within the family or hospital care
- Medication costs: use of prescribed and/or over-the-counter (OTC) medication.

The duration of ILI was defined as the number of days between the 1st day when the child fulfilled the ILI definition and the day when all the symptoms disappeared, inclusive. If a child still had symptoms on the 2nd follow-up telephone call (at 26–32 days), we calculated the ILI duration as difference between the date of the phone call and the ILI onset date.

Costing methods

Details of the applied costs and their source are provided in Table 1. Resource use and unit costs were in 2010 Australian dollars. Cost estimates were also converted into Euro (€), US dollar (US\$) and British pound (£) in the Results section using the 2010 exchange rate [the purchasing power parity (PPP) conversion factor was taken into account].¹³ For the PPP values, we chose International Monetary Fund as the source data set; accordingly, in 2010, AU\$1.00 was equal to €0.56, US\$0.66 and £0.44, respectively. The reported hourly wage rate was used to calculate the costs of recreation time lost among the household members where income information was collected. In those whose income information was not collected, the costs of recreation time lost were calculated using the mean income of an Australian male and female in 2010.¹⁴ As the costs incurred in this study were within a short follow-up period (4 months), a discount rate was not applied (study period < 12 months).

Table 1. Value of applied cost and source of cost items (all in Australian dollars)

Cost item	Value	Applied cost	Source
Healthcare visit			
General practitioner	\$34.90	Group A1 – General Practitioner Attendances to which No Other Item Applies, Level B, Code 23	Medicare Australia ³⁷
Emergency department	\$40.00	Australian Ambulatory Classes group 23 (other respiratory diseases without procedure)	The Australian Government Department of Health and Ageing ³⁸
Specialist	\$82.30	Group A3 – Specialist Attendances to which No Other Item Applies, Code 104	Medicare Australia ³⁷
Petrol use (having healthcare visit for influenza-like illness using private car)	\$0.74 per kilometre	Ordinary car: 1-601–2.6 l	Australian Taxation Office ³⁹

Cost item	Costing methods
Medications	The prices of medicine were obtained from Pharmaceutical Benefits Scheme (PBS) ⁴⁰ if indexed; the prices of non-PBS medicine were from carers' reports
Complementary medicine practitioner	Carers' reports (there was one visit to Chinese medicine practitioner in one ILI episode, cost = \$30.00)
Time off work	Calculated using carers' annual income code ⁴¹
Time off recreation	Calculating using average rate generated from 'Average Weekly Earnings' in Australia, November 2010 ¹⁴
Additional childcare for healthy siblings due to a ILI child or missed childcare of the ILI child	Carers' reports

Statistics

For the descriptive analysis, we used SPSS Statistics (version 19; IBM, Armonk, NY, USA) to calculate the mean costs, with standard deviations (SD), and median costs, with interquartile ranges (IQR), for ILI.

We conducted multivariate analyses (with STATA/SE 12.0; StataCorp LP, USA) to assess the association between the economic outcome of ILI and (i) demographic characteristics; (ii) ILI duration, and (iii) virological outcome variables. Due to the non-normality (right skewing) of the total cost variable in our study, a Tobit regression model was employed to determine variables that drive the ILI costs.¹⁵ We also used a zero-inflated Poisson regression model¹⁶ for comparing the counts of healthcare visits per ILI (that has an excess of zero counts) with demographic variables, ILI duration and virological outcomes. Subgroup analysis of ILI costs was performed based on the four groups of fever grading: 37.8–<38°C, 38–<39°C, 39–<40°C and ≥40°C, using STATA/SE.

Results

We initially recruited 387 children; six of them withdrew prior to the completion of the study (Figure 1). A total of 381 children (Figure 1 and Table 2) from 358 families were followed up over 13 weeks from 30 July 2010, at which date children were aged 0.92–3.4 years. Of these 381 children, 340 (89%) were recruited through childcare and 41 (11%) were through one GP clinic.

In our study, we were unable to estimate the number of parents of potential participants we approached during recruitment. The dates that we attended particular childcare centres were recorded, but because the total number of children attending a centre differed every day of the week, it was impossible to obtain the exact total number of children of the target age group attended each centre. Almost always we spoke to only a subset of those parents because carers might drop off or pick up their children while we were conversing with a few.




Figure 1. Numbers of children with carers' contacts collected, enrolled, followed and analysed. *Reasons of withdrawals were known in two children: one carer said she could not follow the study instructions of data collection. The other was concerned about the privacy of the demographic questions.

Descriptive analyses

There were 124 ILIs in 105 children (13 had two ILIs, three had three). Swabs were provided for 117 ILIs, from which 103 ILIs (88%) had 175 viruses identified (Table 3). Of these, 52 ILIs had one virus identified, 36 ILIs yielded two viruses, 10 yielded three viruses, four yielded four and one yielded five viruses.

In the week before onset of their ILIs, nine children were exposed to one or more household members with ILIs (total three other children and eight adults). In the week after onset

Table 2. Demographics of enrolled children and their households

	Study participants	Australian population (year)
Enrolled children:	381	1 457 571 (0–4 years old) ⁴²
Gender male, % (n)	55 (208)	51 (2010 data) ⁴³
Mean age of study children at 30 July 2010 (years)	2.3	n/a
Mean age of mothers of study children as of 30 July 2010 (years)	33.1	30.1 ¹⁷
Proportion of mothers of study children living with a partner (married or de facto)	97%	88% ¹⁸
Mean no. of siblings in household	0.9	n/a
Ethnicity, % (n)		
Caucasian	82 (313)	n/a
Asian	10 (37)	
Middle Eastern	2 (8)	
Maori/Pacific Islander	1 (3)	
Others	5 (18)	
Missing	1 (2)	
Childcare attendance, % (n)	89 (340)	72 ⁴⁴
Mean days at childcare/week	2.83	n/a
Child sleeping alone, % (n)	272 (71)	n/a
Children's households (358)		
Accommodation, % (n)		
Detached house	77 (277)	71 ⁴⁵
Others	23 (81)	29 ⁴⁵
Smoker(s) in household, % (n)	14 (51)	19–23 ⁴⁶
Parents employment, % (n)		
Both working	73 (262)	49 ⁴⁷
1 working	26 (92)	45 ⁴⁷
Neither working	1 (3)	6 ⁴⁷
Parents' education* [two-parent households% (n)] [†] (348 households)		
Both university	51 (174)	34% of Australians in 2010 had a bachelor degree or higher ⁴⁸
Both other	20 (68)	
1 university, 1 other	29 (98)	
Parents education [single-parent households% (n)] (10 households)		
University	40 (4)	24 ⁴⁹
Other	60 (6)	76 ⁴⁹
Household income >\$2000 per week	75 (264/351)	n/a

*University versus lower level of education.

†More than 1 child per household were enrolled.

Table 3. Mean cost, standard deviation, median cost and interquartile range of ILI presented by household characteristics and virus type

	Number	%	Mean cost (AU\$)	Standard deviation	Median cost (AU\$)	Interquartile range (AU\$)
<i>Characteristics of household with a child who reported ≥ 1 ILI</i>						
Gender (male)	62	50	637	812	347	62–986
Age [year(s)]						
<1	5	4	909	469	1034	761–1053
1	52	42	569	721	262	25–848
2	51	41	762	967	509	109–1113
3	16	13	292	409	102	0–431
No. of siblings in household						
0	45	36	752	811	559	196–1065
1	51	41	535	837	117	17–810
2	27	22	606	751	428	59–923
3	1	1	175	n/a	1	n/a
Ethnicity (Caucasian)	99	80	635	770	371	57–1032
Childcare attendance	113	91	653	813	380	65–1012
No. of days at childcare for childcare attendants						
1	12	10	367	589	89	10–346
2	49	40	654	849	363	69–867
3	27	22	624	562	569	134–1099
4	11	9	633	828	220	87–858
5	14	11	971	1184	523	155–1498
Child sleeping alone	87	70	633	803	305	60–972
Accommodation (detached house versus others)	103	83	631	844	305	41–961
No. of smoker in household (none versus ≥ 1)	108	87	627	759	376	59–989
Parents' employment (both working versus others)	92	74	726	864	523	95–1060
Parents' education in two-parent households (both university)*	69	56	597	834	216	17–1058
<i>Virus types</i>						
Adenovirus	39	32	648	869	286	39–960
Rhinovirus	39	32	622	1012	210	17–793
Parainfluenza 3	22	18	734	946	139	54–1124
WU (polyomavirus)	22	18	545	990	118	21–587
Human metapneumovirus	12	10	714	459	747	422–1075
Bocavirus	11	9	772	558	636	399–1055
A(H1N1)pdm09	5	4	599	485	380	216–909
Polyomavirus (WU and/or KI)	27	22	570	929	196	52–690
Coronavirus (HKU-1 and/or NL63)	15	12	752	513	636	425–1141
No virus detected	14	11	459	640	124	10–730
No swab	7	6	519	604	220	128–737
Single infection	50	40	580	745	280	59–858
Co-infection	53	43	729	922	559	60–1072
All ILIs	124	100	626	806	321	50–989

of the child's ILI, eight other children and 38 adults in the sick child's household reported ILIs.

Managing ILIs required 134 GP visits for 70 of the ILI episodes (the other 54 ILIs did not involve GP visits), 106 pharmacy visits for 64 ILIs, 5 ED visits and three specialist visits, but no hospitalisations. The proportion of ILI events associated with ≥ 1 healthcare visits was 58% (72/124). Antibiotics were used for 52 ILIs, and 73 involved treatment with analgesic/antipyretics. The median duration of ILIs was 8 days (IQR: 5–12 days), and mean duration was 9 days

(95% CI: 8–10 days); however, 16 ILIs (13%) lasted >28 days.

The mean cost of the 124 ILIs was AU\$626 (equal to €351, US\$413 or £275) with a median cost of AU\$321 (€180, US \$212 or £141). Of the mean ILI cost, AU\$406 (65%; €227, US \$268 or £179) was due to carer time off work, AU\$102 (16%; €57, US\$67 or £45) was generated by healthcare visits and AU\$118 (19%; €66, US\$78, or £52) was due to other items, such as medications and missed childcare attendance (Figure 2).




Figure 2. Distribution of the costs in the 124 documented ILIs by cost category.

The mean time off work per ILI episode was 12.9 hours (median 4 hours). The mean ILI-related time away from recreation and associated cost was 3.1 hours (median 0 hour). The mean and median costs of ILI by virus identification are presented in Table 3. There were no significant differences in mean cost of ILI by virus identified.

Multivariate analyses

The analyses by Tobit regression (Table 4) revealed two independent drivers of ILI costs, namely having both parents in paid employment and a longer ILI duration (Table 4). The

other characteristic factors and virological outcome variables were not significant in the Tobit model (data not shown).

The comparisons for the numbers of healthcare visit per ILI using zero-inflated Poisson regression are also shown in Table 4. There were four variables associated with a higher number of healthcare visits for ILI, which were (i) being non-Caucasian; (ii) living in a detached house; (iii) having both parents in paid employment, and (iv) having an ILI with one or more viruses identified. However, among the virus-positive ILIs, the number of healthcare visit did not significantly increase with increasing number of detected viruses.

Of the 124 ILIs, temperature was measured in 84 (68%) episodes; the other ILIs were reported to have 'fever/feverish' by the carer. Among these 84 ILIs, there were 8 (10%) with a reported temperature being 37.8–38°C, 43 ILIs (51%) for 38–39°C, 23 ILIs (27%) for 39–40°C and 10 ILIs (12%) for 40°C or over. No statistically significant difference was detected in the ILI costs ($P = 0.06$, Tobit regression) and number of healthcare visit per ILI ($P = 0.21$, zero-inflated Poisson regression).

Discussion

In our study population, largely consisting of healthy urban children attending childcare, viral respiratory illnesses were frequent and costly, with excess carer time off work (13 hours per ILI episode) and recreation time lost (3 hours per ILI) contributing most to total costs. Cost of illness did

Table 4. Significant variables (using multivariate analyses) associated (1) with increased ILI* costs or (2) with higher number of healthcare visits

(1) Variables associated with increased ILI costs using Tobit regression

	Coefficient	95% CI	P value
Parents' employment (one or neither working versus both working)	505.84	170.38 – 841.30	0.003
ILI duration (each 1 day increase in duration)	18.21	0.53 – 35.89	0.04

(2) Variables associated with higher number of healthcare visits (to a GP, pharmacy, ED and/or specialist room) using zero-inflated Poisson regression

	(No. of ILI)	No. of healthcare visit per ILI during the follow-up period	P value
Ethnicity	Caucasian (99)	1.63	0.002
	Non-Caucasian (23)	2.74	
Accommodation	Non-detached house (21)	1.10	<0.001
	Detached house (103)	1.99	
Parents employment	One or neither working (31)	0.87	<0.001
	Both working (91)	2.16	
ILI with one or more viruses identified	None (14)	1.00	0.03
	≥1 virus (103)	1.98	

*Influenza-like illness.

not vary by the type of virus identified. It may be that costs of ILIs in children attending childcare were less affected by virus type, due to the almost uniform requirement for at least one parent to have time away from work (if both work) regardless of severity. In our study, having both parents in employed work and having a longer ILI duration were (perhaps predictably) the key cost drivers. Four variables appeared to be associated with increased number of healthcare visits for ILI: being non-Caucasian, living in a detached house, having working parents (i.e. both) and having an ILI that is positive for at least one virus.

Our findings are largely applicable to urban families with a child attending organised childcare. Strengths of our study include regular reminders to carers to identify any ILIs in the study child by weekly contact. In addition, parent-collected swabs and sensitive RT-PCR for virus identification enabled sensitive detection of outcome parameters; we also achieved a high swabbing rate [94% (117/124) of the ILI episodes]. Our study was observational and study staff did not provide medical care for study child, reducing the likelihood of a Hawthorne effect artificially reducing ILI costs. Finally, this study provides ILI cost-of-illness data for urban children with simple to complex ILIs, attending childcare and with busy working parents – a common and growing demographic in Australian society.

Limitations include the fact that study children and families were not representative of the broader Australian population. Most (89%) study children attended childcare, and compared with the general Australian population,¹⁴ study households had higher income (Table 2), leading to an under-representation of lower-income households. Previous Australian research has shown that the highest mean acute respiratory illness (ARI) cost-of-illness values are not necessarily associated with the highest income households, reducing the likelihood our study population may have upwardly biased mean ILI costs. In addition, as shown in Table 2, the mothers in our study were slightly older (33 years compared with 30 years in the general population¹⁷) and they were more likely to live with a partner (married or de facto) (97% versus 88%).¹⁸

This study showed a low prevalence of A(H1N1)pdm09 in 2010. We believe this is partly because we did not start collecting ILI data until 30 July 2010, thus, some winter cases of influenza may have been missed. Surveillance data revealed that the 2010 influenza peak was September in the state where we conducted the study (NSW),¹⁹ but the incidence of influenza in 2010 was relatively low.¹⁹

It is noteworthy that we used a relatively comprehensive approach to collect information on the costs of ILI, and our study reinforced the importance of indirect costs in economic assessment of infectious diseases. Apart from the major cost drivers (i.e. time off work, healthcare visits), we also collected data on other cost items; these included non-

antibiotic prescription medication, OTC medications, and paid childcare for other children (in the same household of the ILI child), whilst carers were looking after the sick child, travel costs of healthcare visits and all other costs that the carers recalled. In addition, our study highlighted that the time off recreational activities should be adequately measured because it covered a not insubstantial 19% (3/16 hours) of total time lost.

Our study identified some novel variables associated with ILI costs and more healthcare visits for the ILI. Having both parents in paid employment increased the ILI costs probably due to the primary carer and the partner taking time off work to look after the sick child. The lost productivity due to time off work plus costs of recreation time lost accounted for the majority of ILI costs in this study that is consistent with the literature.^{20,21} In our study, particularly where both parents work, even simple ILIs without a GP visit can be costly because of carers' taking time off work. Apart from economic impact, qualitative interviews conducted with a subsample of the carers in this study also demonstrated that these parents have larger quality-of-life impact in terms of daily and social life disruption.²²

To our knowledge, ours is the first study that demonstrates the duration of respiratory infection, when managed outside a hospital, is associated with higher economic costs. This finding may suggest that for future efforts in assessing the economic impacts of a respiratory disease, the measurement of infection duration should be taken into account as a proxy for economic burden. Of the four variables that increased the number of healthcare visits with an ILI, we believe that two of them (living in a detached house and having working parents) are associated with potentially higher socio-economic status. In the literature, it is evident that those of higher economic class are more likely to seek medical visits for their health issues.^{23–25} However, our study did not collect related information to prove the potential link between socio-economic status and health-seeking behaviours.

In our study, the children with virus-positive ILIs had more healthcare visits than those with no virus detected; however, among the virus-positive ILIs, the number of healthcare visits did not significantly increase with increasing number of viruses identified. The use of RT-PCR has contributed to ongoing discovery/detection of simultaneous multiple respiratory pathogens from a single specimen in one child,²⁶ but the real impact of these simultaneous detections remains unclear. There are some reports documenting that (i) the single infection of human metapneumovirus (HMPV), and (ii) the co-infection of HMPV plus respiratory syncytial virus increase hospitalisation rates,^{27,28} while other studies did not detect an association between a multiple viral infections and illness severity.^{29–31} Future research should endeavour to better correlate clinical/economic disease severity with viral quantity.

In valuing the recreation time lost, we used sex-weighted mean income in Australia. There is some controversy on the valuation of recreation time.^{32,33} An economic analysis using a societal perspective requires that all costs resulting from an illness should be taken into account, regardless of who experiences the costs³⁴; thus, we believe measuring and valuing all time spent on an ILI episode are important, including that removed from recreation. The recreation time loss is likely to cause opportunity costs, which needs to be assessed in an economic analysis.

A study by Lambert and colleagues reported the costs of ARIs among 234 preschool children from Victoria, Australia, in 2003.³⁵ The mean and median costs of all ARIs were \$309 and \$156, respectively, which were somewhat lower than our findings. This may be due to inflation, different methods of valuing time and the fact that our study included a higher proportion of families where both parents worked (89% in this study versus 49% in Lambert *et al.*⁸). This study's estimate of the average weekly income of a working person was \$1323, which was around 30% higher than the average income of employed Australians.

A recent French survey of >800 household contacts by Carrat examined the disease burden of influenza in 2010.³⁶ Their results on illness duration and proportion of index cases having a medical visit were similar to our findings. Their estimate on median duration of illness in laboratory-confirmed index cases was 8 days (95% CI: 7–8 days), but their median age was higher (36.7 ± 19.4 years old). Carrat reported that 57% of household members with clinical influenza (not laboratory confirmed) sought medical consultations; in our study, 58% (72/124) of ILIs had at least one medical visit. Carrat's study was conducted in a population sample that was representative of the general population in France.

In summary, this study's finding of a substantial social and economic burden of ILI in children attending childcare highlights the importance of interdisciplinary research in providing primary data for policy making of a possible preventive programme for respiratory infections. To link the disease impact and economic assessment results, interdisciplinary research (epidemiology and health economics) is needed to better assess the desirability and feasibility of preventive or therapeutic interventions.

Disclosure

This study is registered with the Australian New Zealand Clinical Trials Registry (ANZCTR, reference number ACTRN12610000319077) and funded by the Australian Research Council (Linkage Project Scheme LP0884126 with the Kindergarten Union Children's Services and Sanofi Pasteur as partner organisations providing funding or in-kind support).

Conflict of interest

J. Kevin Yin received an educational grant from Sanofi Pasteur for influenza economic research in 2012. Stephen Lambert has received funding from CSL, Sanofi, Glaxo-SmithKline (GSK) and Wyeth to conduct sponsored research, attend/present at scientific meetings and participate in vaccine advisory boards. Leon Heron has received funding from Baxter, CSL, GSK, Merck, Novartis, Pfizer, Roche, and Sanofi Pasteur for the conduct of sponsored research, travel to present at conferences or consultancy work; all funding received is directed to research accounts at the Children's Hospital at Westmead. Robert Booy has received funding from CSL, Roche, Sanofi, GSK, Novartis and Pfizer to conduct sponsored research or attend and present at scientific meetings; any funding received is directed to a research account at the Children's Hospital at Westmead. The other authors declared no conflict of interest.

Acknowledgements

The authors thank the participants and their parents, Australian Research Council, Sanofi Pasteur, KU Children's Services (Sydney), the research nurses (Laura Rost, Camille Lang, Lisa Chalmers, Elizabeth Clarke, Elizabeth Deegan and Edwina Jacobs), Theo P Sloots, Michael D Nissen and other staff members at the Queensland Paediatric Infectious Disease Laboratory.

References

- 1 Chiu C, Dey A, Wang H *et al.* Vaccine preventable diseases in Australia, 2005 to 2007. *Commun Dis Intell* 2010; 34(Suppl):S1–S167.
- 2 Esposito S, Gasparini R, Bosis S *et al.* Clinical and socio-economic impact of influenza and respiratory syncytial virus infection on healthy children and their households. *Clin Microbiol Infect* 2005; 11:933–936.
- 3 Iskander M, Booy R, Lambert S. The burden of influenza in children. *Curr Opin Infect Dis* 2007; 20:259–263.
- 4 Fendrick AM, Monto AS, Nightengale B, Sarnes M. The economic burden of non-influenza-related viral respiratory tract infection in the United States. *Arch Intern Med* 2003; 163:487–494.
- 5 Newall AT, Scuffham PA. Influenza-related disease: the cost to the Australian healthcare system. *Vaccine* 2008; 26:6818–6823.
- 6 Australian Bureau of Statistics. Childhood education and care, Australia, June 2008 (Reissue), Cat. no. 4402.0. 2009; Available at [http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4402.0June%202008%20\(Reissue\)?OpenDocument](http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4402.0June%202008%20(Reissue)?OpenDocument) (Accessed 6 January 2012).
- 7 National Health and Medical Research Council. Staying Healthy in Child Care – preventing infectious diseases in child care. 2007; 4th. Available at http://www.nhmrc.gov.au/_files_nhmrc/publications/attachments/ch43.pdf (Accessed 6 January 2012).
- 8 Lambert SB, Allen KM, Druce JD *et al.* Community epidemiology of human metapneumovirus, human coronavirus NL63, and other respiratory viruses in healthy preschool-aged children using parent-collected specimens. *Pediatrics* 2007; 120:17.

- 9 Yin JK, Lahra MM, Iskander M *et al.* Pilot study of influenza vaccine effectiveness in urban Australian children attending childcare. *J Paediatr Child Health* 2011; 10:1440–1754.
- 10 Lu X, Holloway B, Dare RK *et al.* Real-time reverse transcription-PCR assay for comprehensive detection of human rhinoviruses. *J Clin Microbiol* 2008; 46:533–539.
- 11 O'Grady KA, Torzillo PJ, Rockett RJ *et al.* Successful application of a simple specimen transport method for the conduct of respiratory virus surveillance in remote Indigenous communities in Australia. *Trop Med Int Health* 2011; 16:766–772.
- 12 Bialasiewicz S, Whitley DM, Lambert SB, Nissen MD, Sloots TP. Detection of BK, JC, WU, or KI polyomaviruses in faecal, urine, blood, cerebrospinal fluid and respiratory samples. *J Clin Virol* 2009; 45:249–254.
- 13 Shemilt I, Thomas J, Morciano M. A web-based tool for adjusting costs to a specific target currency and price year. *Evid Policy* 2010; 6:51–59.
- 14 Australian Bureau of Statistics. 6302.0 – average weekly earnings, Australia, Nov 2010 2011; Available at <http://www.abs.gov.au/AUSSTATS/abs@.nsf/0/2BDA3ABC6DB85EA5CA25789400163EB8?opendocument> (Accessed 23 December 2011).
- 15 McDonald JF, Moffitt RA. The uses of tobit analysis. *Rev Econ Stat* 1980; 62:318–321.
- 16 Lambert D. Zero-inflated Poisson regression, with an application to defects in manufacturing. *Technometrics* 1992; 34:1–14.
- 17 Australian Institute of Health and Welfare (AIHW). Perinatal statistics series, Number 24. Australia's mothers and babies 2008. Cat. no. PER 50. 2010; Available at <http://www.aihw.gov.au/WorkArea/DownloadAsset.aspx?id=6442472762> (Accessed 19 April 2013).
- 18 Australian Institute of Health and Welfare (AIHW). A picture of Australia's children 2009. Cat. no. PHE 112. 2009; Available at <http://www.aihw.gov.au/WorkArea/DownloadAsset.aspx?id=6442459928> (Accessed 19 April 2013).
- 19 NSW HEALTH. Influenza monthly epidemiology report, NSW. 2011; Available at http://www0.health.nsw.gov.au/resources/publichealth/infectious/influenza/pdf/20110106_monthly_report.pdf (Accessed 26 January 2013).
- 20 Molinari NA, Ortega-Sanchez IR, Messonnier ML *et al.* The annual impact of seasonal influenza in the US: measuring disease burden and costs. *Vaccine* 2007; 25:5086–5096.
- 21 Simmerman JM, Lertindumrong J, Dowell SF *et al.* The cost of influenza in Thailand. *Vaccine* 2006; 24:4417–4426.
- 22 Chow MY, Morrow A, Booy R, Leask J. The impact of children's influenza-like-illness on parental quality of life: a qualitative study. *J Paediatr Child Health* 2013; doi: 10.1111/jpc.12261.
- 23 Williamson DL. Health behaviours and health: evidence that the relationship is not conditional on income adequacy. *Soc Sci Med* 2000; 51:1741–1754.
- 24 Maclean H, Glynn K, Cao Z, Ansara D. Personal health practices. *BMC Womens Health* 2004; 4(Suppl. 1):S4.
- 25 Pomerleau J, Pederson LL, Ostbye T, Speechley M, Speechley KN. Health behaviours and socio-economic status in Ontario, Canada. *Eur J Epidemiol* 1997; 13:613–622.
- 26 Paranhos-Baccalà G, Komurian-Pradel F, Richard N, Vernet G, Lina B, Floret D. Mixed respiratory virus infections. *J Clin Virol* 2008; 43:407–410.
- 27 Foulongne V, Guyon G, Rodiere M, Segondy M. Human metapneumovirus infection in young children hospitalized with respiratory tract disease. *Pediatr Infect Dis J* 2006; 25:354–359.
- 28 Semple MG, Cowell A, Dove W *et al.* Dual infection of infants by human metapneumovirus and human respiratory syncytial virus is strongly associated with severe bronchiolitis. *J Infect Dis* 2005; 191:382–386.
- 29 Singleton RJ, Bulkow LR, Miernyk K *et al.* Viral respiratory infections in hospitalized and community control children in Alaska. *J Med Virol* 2010; 82:1282–1290.
- 30 Wolf DG, Greenberg D, Kalkstein D *et al.* Comparison of human metapneumovirus, respiratory syncytial virus and influenza A virus lower respiratory tract infections in hospitalized young children. *Pediatr Infect Dis J* 2006; 25:320–324.
- 31 Wilkesmann A, Schildgen O, Eis-Hubinger AM *et al.* Human metapneumovirus infections cause similar symptoms and clinical severity as respiratory syncytial virus infections. *Eur J Pediatr* 2006; 165:467–475.
- 32 Brouwer WB, Koopmanschap MA, Rutten FF. Patient and informal caregiver time in cost-effectiveness analysis. A response to the recommendations of the Washington Panel. *Int J Technol Assess Health Care* 1998; 14:505–513.
- 33 Sculpher M. The role and estimation of productivity costs in economic evaluation; in Drummond MF, McGuire A (eds): *Economic Evaluation in Healthcare: Merging Theory With Practice*. Oxford, UK 94-112: Oxford University Press, 2001.
- 34 Gold MR, Siegel JE, Russel LB, Weinstein MC. *Cost-effectiveness in Health and Medicine*. Oxford, UK: Oxford University Press, 1996.
- 35 Lambert SB, Allen KM, Carter RC, Nolan TM. The cost of community-managed viral respiratory illnesses in a cohort of healthy preschool-aged children. *Respir Res* 2008; 9:11.
- 36 Carrat F, Sahler C, Rogez S *et al.* Influenza burden of illness: estimates from a national prospective survey of household contacts in France. *Arch Intern Med* 2002; 162:1842–1848.
- 37 Australian Government Department of Health and Ageing. Medicare benefits schedule book operating from 01 November 2010; Available at [http://www.mbsonline.gov.au/internet/mbsonline/publishing.nsf/Content/69F5578743C9B654CA257790001AD472/\\$File/201011-MBS.pdf](http://www.mbsonline.gov.au/internet/mbsonline/publishing.nsf/Content/69F5578743C9B654CA257790001AD472/$File/201011-MBS.pdf) (Accessed 6 December 2010).
- 38 Department of Health and Ageing, Australia. Manual of Resource Items and their Associated Costs for use in submissions to the Pharmaceutical Benefits Advisory Committee involving economic evaluation. Available at <http://www.health.gov.au/internet/main/publishing.nsf/Content/health-pbs-general-pubs-manual-appendix3.htm> (Accessed 1 November 2011).
- 39 Australian Taxation Office. Claiming a deduction for car expenses using the cents per kilometre method. 2011; Available at <http://www.ato.gov.au/individuals/content.aspx?doc=/content/33874.htm> (Accessed 05 September 2011).
- 40 Australian Government Department of Health and Ageing. Pharmaceutical Benefits Scheme (PBS). Available at <http://www.pbs.gov.au/browse/medicine-listing> (Accessed 2 October 2011).
- 41 Australian Bureau of Statistics. 2901.0 – Census dictionary, 2006 (Reissue). 2006; Available at <http://www.abs.gov.au/Ausstats/abs@.nsf/0/A17B27BBF0273101CA25729E0008A882?opendocument> (Accessed 22 October 2011).
- 42 Australian Bureau of Statistics. 3235.0 – Population by age and sex, Regions of Australia, 2011. 2012; Available at <http://www.abs.gov.au/ausstats/abs@.nsf/Products/3235.0~2011~Main+Features~Main+Features?OpenDocument#Top> (Accessed 23 December 2012).
- 43 Australian Bureau of Statistics. 3101.0 – Australian demographic statistics, Dec 2010. 2011; Available at <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3101.0Dec%202010?OpenDocument> (Accessed 23 December 2012).
- 44 Australian Bureau of Statistics. Australian Social Trends 4102.0 Dec 2009. Preschool attendance. 2009; Available at [http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/907180B96AFA4596CA257687001D4B06/\\$File/41020_Preschool.pdf](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/907180B96AFA4596CA257687001D4B06/$File/41020_Preschool.pdf) (Accessed 23 December 2012).
- 45 Australian Bureau of Statistics. 3240.1 – Housing Choices, NSW, Oct 2004. 2005; Available at <http://www.abs.gov.au/AUSSTATS/abs@.nsf/>

- ProductsbyTopic/1C87D9876BC11EE8CA256FD5007722A8?OpenDocument (Accessed 23 December 2012).
- 46** Australian Bureau of Statistics. Australian Social Trends 4102.0 Dec 2009. Smoking, risky drinking and obesity. 2009; Available at [http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/0B008A644981DCDDCA257687001D4987/\\$File/41020_riskfactors.pdf](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/0B008A644981DCDDCA257687001D4987/$File/41020_riskfactors.pdf) (Accessed 23 December 2012).
- 47** Australian Bureau of Statistics. 4442.0 – Family characteristics, Australia, 2009–10. 2011; Available at <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4442.02009-10?OpenDocument> (Accessed 23 December 2012).
- 48** Australian Bureau of Statistics. 6227.0 – Education and work, Australia, May 2010. 2010; Available at <http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/6227.0Main+Features1May%202010?OpenDocument> (Accessed 23 December 2012).
- 49** Australian Bureau of Statistics. Education and training experience 6278.0. 2010; Available at [http://www.ausstats.abs.gov.au/Ausstats/subscriber.nsf/0/4EF49EB9552322F4CA2576F500120083/\\$File/62780_2009.pdf](http://www.ausstats.abs.gov.au/Ausstats/subscriber.nsf/0/4EF49EB9552322F4CA2576F500120083/$File/62780_2009.pdf) (Accessed 23 December 2012).