


The exponential formula

We want to find a general construction for certain labelled objects in order to apply a unified formula for all of them.

Def. A **CARD** is a pair (S, p) where p is a "picture" (e.g., a structure with empty labels) and $S \subseteq \mathbb{N}$ is a finite set of labels. We order its **WEIGHT** ordered with the usual \leq .

Example. We can represent an m -cycle as a labelled conf. in $[m]$ to which we can associate a set $S = \{k_1 \leq \dots \leq k_m\}$ of labels. $(1, 3, 5, 2, 4)$ with labels $k_1 \leq \dots \leq k_5$ is $(k_1, k_3, k_5, k_2, k_4)$. (Just like $(1, 3, 5, 2, 4)$)




Def. A card (S, p) is said to be **STANDARD** if $S = [|S|]$.

Def. A **RELABELLING** of a card (S, p) is a card (S', p) with $|S'| = |S|$. We say (S', p) is a **STANDARD RELABELLING** if (S', p) is a standard card.

Def. A **HAND** is a set of cards whose labels form a partition of $[m]$ for some $m \in \mathbb{N}^+$. We call the sum of the weights of its cards its **WEIGHT**.

Example. We can think of a permutation $\sigma \in S_m$ as a hand of cycles, which behave as its cards.



Def. A **DECK** is a finite set of standard cards with the same weight; we call such weight the **WEIGHT** of the deck.

→ a deck contains the "prototype of the cards", i.e., "their configurations".

Def An **EXPONENTIAL FAMILY** F is a collection of decks $\{D_i\}_{i \in \mathbb{N}}$ s.t. D_i has weight i and $D_0 \triangleq 0$.

If we set $d_i \triangleq |D_i|$, we can consider the e.g.f. of $\{d_i\}$:

$$D(x) \triangleq \sum_{m \geq 0} \frac{d_m}{m!} x^m,$$

which is called The **DECK ENUMERATOR** of F .

Example (permutations) As we saw, permutations are hands made of cycles. Then, a suitable family for permutations is $\{D_i\}_{i \in \mathbb{N}}$ s.t.

$$D_i \triangleq \{i\text{-cycles in } S_i\} \implies d_i = \frac{1}{i} = (i-1)! \quad (i \neq 0)$$

$$\text{Hence } D(x) = \sum_{i \geq 1} \frac{(i-1)!}{i!} x^i = \sum_{i \geq 1} \frac{x^i}{i} = -\log(1-x) = \log\left(\frac{1}{1-x}\right).$$

Def For each $m, k \geq 0$ we let $h(m, k)$ be the **number of hands of weight m that consist of k cards** s.t. each card is a relabelling of some card in some deck of F . By convention $h(m, 0) = \delta_{m,0}$.

We call $H(x, y) \triangleq \sum_{m, k \geq 0} \frac{h(m, k)}{m!} x^m y^k$ the **HAND NUMERATOR** of F .

Example (permutations) $h(m, k)$ counts in F_a the number of permutations $\in S_a$ with k cycles.

\leadsto by setting $y=1$ in $H(x, y)$ one gets the e.g.f. of $\{h(x)\}$ where:

$$h(m) \triangleq \#\{\text{hands of weight } m\}.$$

Our goal is to find a relation between $D(x)$ and $H(x, y)$.

Before that we introduce such a relation, we explore the direct sum operation:

Def. Let $F = \{D_i\}_{i \in \mathbb{N}}$ and $F' = \{D'_i\}_{i \in \mathbb{N}}$ be two exponential families s.t. $D_i \cap D'_i = \emptyset \quad \forall i$. Then we define their **MERGE** as: all pictures are diff.

$$F'' = F \oplus F' = \{D''_i = D_i \cup D'_i\}_{i \in \mathbb{N}}$$


$$\leadsto D_{F \oplus F'}(x) = D_F(x) + D_{F'}(x)$$

Prop. (fundamental lemma of labelled counting) If $F'' = F \oplus F'$, then:

$$H_{F''}(x, y) = H_F(x, y) \cdot H_{F'}(x, y)$$

We observe that a hand of weight n made of k cards can be split in two hands: one coming from F and one coming from F' s.t. their weights sum to n and the numbers of their cards sum to k .

" of weight $3+5+6 = 14$
with $2+1 = 3$ cards in F''



Hence the following holds:

$$H_{F''}(m, k) = \sum_{m'=0}^m \sum_{k'=0}^k \binom{m}{m'} H_F(m', k') H_{F'}(m-m', k-k')$$

we choose the labels we pick in F

Since the RHS is exactly $\left[\frac{x^m}{m!} y^k \right] H_F(m, k) H_{F'}(m, k)$, the Theorem is proved. \square

Theorem (Exponential formula) The following formula holds:

$$H(x, y) = e^{y \Delta(x)}$$

Let F_r be an exponential family s.t. $D_i = \emptyset \iff i \neq r$ and D_r consists of only one card.

Then $D_{F_r}(x) = \frac{x^r}{r!}$. Let's count now $h(m, k)$: if $m \neq k \cdot r$, $h(m, k)$ will necessarily be zero since there is no way to build a hand of weight m ; if $m = k \cdot r$, we must employ r cards and we can do so in $\frac{1}{k!} \underbrace{(r \cdot r \cdots r)}_{k \text{ times}}$ ways.

To avoid permutations

Let $\binom{m}{m_1 \dots m_k}$ denote the **MULTINOMIAL COEFFICIENT** which counts how many families $\{S_i\}_{i=1 \dots k}$ we can have s.t. $S_i \subseteq [m]$, $|S_i| = m_i$, $S_i \cap S_j = \emptyset \iff i \neq j$, when $m_1 + \dots + m_k = m$.

We observe that $\binom{m}{m_1 \dots m_k} = \binom{m}{m_1} \binom{m-m_1}{m_2} \cdots \binom{m-m_1-\dots-m_{k-1}}{m_k}$, hence:

$$\binom{m}{m_1 \dots m_k} = \frac{m!}{m_1! \cdots m_k!}$$

$$\leadsto \binom{m}{k, m-k} = \binom{m}{k}$$

$$\begin{aligned} \text{Therefore } h_{F_r}(kr, k) &= \frac{(kr)!}{k! r! k} \Rightarrow H_{F_r}(x, y) = \sum_{k \geq 0} \frac{\cancel{(kr)!}}{r! k} \frac{x^{kr}}{\cancel{(kr)!}} y^k / k! = \\ &= \sum_{k \geq 0} \left(\frac{x^r}{r!} y \right)^k / k! = e^{y \frac{x^r}{r!}} \end{aligned}$$

Since each exponential family is the (possibly infinite) merge of families isomorphic to F_r , by applying the previous lemma the proof follows. \square

$$H = \prod_i H_i + e^{x+y} = e^x e^y$$

→ since $H(x, y) = e^{yD(x)}$,
$$h(m, k) = \left[\frac{x^m}{m!} \right] \frac{D(x)^k}{k!}$$

Example. (permutations) Let's count how many permutations in S_m have k cycles:

$$D(x) = \log((1-x)^{-1}) \xRightarrow[\substack{\text{exp. formula} \\ H(x, y) = (1-x)^{-y}}]{\quad} h(m, k) = \left[\frac{x^m}{m!} \right] \frac{\log(1/(1-x))^k}{k!}$$

$h(m, k)$ is denoted as $\begin{bmatrix} m \\ k \end{bmatrix}$ and such numbers are called **The (unsigned) Stirling numbers of first kind.**

$$\begin{aligned} \sum_{k \geq 0} \begin{bmatrix} m \\ k \end{bmatrix} y^k &= \left[\frac{x^m}{m!} \right] (1-x)^{-y} = m! (-1)^m \binom{-y}{m} = (-1)^m (-y)_m = \\ &= (-1)^m (-1)^m (y+m-1)_m = (y+m-1)_m \stackrel{\Delta}{=} y^{(m)} \quad (1) \end{aligned}$$

$(-y)_m = (-y) \cdots (-y-m+1)$
 $= (-1)^m (y+m-1) \cdots y =$
 $= (-1)^m (y+m-1)_m =$
 $= (-1)^m y^{(m)}$

rising factorial

By substituting $y = -x$ in (1) we get:

$$(x)_m = (-1)^m (-x)^{(m)} = \sum_{k \geq 0} (-1)^{m-k} \begin{bmatrix} m \\ k \end{bmatrix} x^k \quad (2)$$

Therefore:

- The coeff's of $x^{(m)}$ are The **(unsigned) Stirling numbers of first kind** $\begin{bmatrix} m \\ k \end{bmatrix}$
- The coeff's of $(x)_m$ are The **(signed) Stirling numbers of first kind** (see (2).)

Moreover (2) gives us The elements of the matrix changing:

$$\underbrace{\{1, (x)_1, (x)_2, \dots\}}_{\text{falling factorial basis}} \quad \text{into} \quad \underbrace{\{1, x, x^2, \dots\}}_{\text{monomial basis}}$$

Example. (partitions) Let's count partitions. Each D_i will have only one card, i.e., $D_i = \{[i]\}$.

Hence $D(x) = \sum_{k \geq 1} \frac{x^k}{k!} = e^x - 1 \Rightarrow H(x, y) = e^y (e^x - 1)$.

Recall that $\left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\} \triangleq h(m, k)$. Then:

$$\sum_m \left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\} \frac{x^m}{m!} = [y^k] H(x, y) = \frac{D(x)^k}{k!} = \frac{(e^x - 1)^k}{k!},$$

which gives us the e.g.f. for the Stirling numbers of second kind.

Moreover — recalling that $b(m) \triangleq \sum_k \left\{ \begin{smallmatrix} m \\ k \end{smallmatrix} \right\}$ —

$$B(x) = H(x, 1) = e^{e^x - 1},$$

which we already knew.

Example. Let $a(m) = \# \{ \sigma \in S_m \mid \sigma^m = 1 \}$, for a fixed $m \in \mathbb{N}$.

Recall that $\sigma^m = 1$ iff every cycle in σ has a length that divides m .

We can consider a family where $d_i = (i-1)!$ if $i \mid m$ and $d_i = 0$ otherwise.

Then $H(x) = H(x, 1) = e^{D(x)}$, where $D(x) = \sum_{d \mid m} \frac{x^d}{d!}$.

$$\sum_m \frac{a(m)}{m!} x^m$$

For example, if $m=2$, $H_2(x) = e^{x + \frac{x^2}{2}}$.

→ such an example can be adapted to find the number of permutations $\in S_m$ with odd cycles.

In all the examples we gave above, we computed $D(x)$ first and then we deduced $H(x, y)$. Sometimes it's better to go in the other direction.

$$H(x, y) = e^{yD(x)} \Rightarrow \frac{\partial}{\partial x} H(x, y) = y D'(x) e^{yD(x)}$$

Recall that deriving an e.g.f. is equivalent to shifting by one its coeff. Thus:

$$\begin{aligned} h(m+1, k) &= \left[x^m / m! \right] \left[y^k \right] D'(x) \cdot y e^{yD(x)} \\ &= \sum_j \binom{m}{j} d_{m+1-j} h(j, k-1) \end{aligned}$$

By summing over all k 's (with $h(m) \triangleq \sum_k h(m, k)$):

$$\begin{aligned} h(m+1) &= \sum_j \binom{m}{j} d_{m+1-j} h(j) \Rightarrow \\ \Rightarrow \left[d_{m+1} &= h(m+1) - \sum_{j \geq 1} d_{m+1-j} h(j) \right] \quad (3) \end{aligned}$$

which gives us a tool to get $D(x)$ from $H(x, 1) = H(x)$.

Example. (connected labelled graphs) We know how many graphs on n vertices there are:

$$h(n) = 2^{\binom{n}{2}} \quad \left. \vphantom{h(n)} \right\} \begin{array}{l} \text{for each edge, we need} \\ \text{to choose whether to} \\ \text{include it or not} \end{array}$$

Hence the number of connected labelled graphs (which make up for our decks) is given by (3):

$$\left[d_{m+1} = 2^{\binom{m+1}{2}} - \sum_{j \geq 1} 2^{\binom{m+1-j}{2}} d_{m+1-j} \right]$$