

What is ANTIPROL™?

ANTIPROL™ is an essential herbal care formulation for prevention and treatment of prolapse and intussusception in birds. ANTIPROL™ acts by strengthening of intestinal muscles, fine-tunes the muscle tone, and balances the conditions of dysperistalsis or hyperperistalsis in birds.

Composition:

ANTIPROL™ is an oral formulation that contains Vitamin A, detoxifying agents, natural antioxidants, shilajit and herbal extracts such as Withania somnifera, Curcuma longa and Ocimum Sanctum.

Key Benefits:

- ❖ Fine tunes body metabolism and helps in the prevention of abnormal weight gain.
- ❖ Corrects hormonal imbalances and there by prevents the issues with early maturity.
- ❖ Acts as an astringent and helps in skin tightening, strengthening of the muscles and ligaments of intestine and lower viscera
- ❖ Acts as an anti-inflammatory agent and helps in smoothening and reduction of inflammation in cloacal /vent region.
- ❖ Reduces the effects of mycotoxins such as fusariotoxins.
- ❖ Prevents subclinical enteritis, respiratory diseases, coccidiosis and mycosis.
- ❖ Acts at receptor level, Blocks any raise in specific receptor for peristalsis & Fine tunes the peristaltic movement of Gastro Intestinal Tract (GIT).
- ❖ Improves cloacal muscle tone, Improves mucus secretion from shell gland & enhances lubrication.

Dosage:

For broiler breeders & Commercial Layers: 10-15 ml /100 birds for 5-7 days.
(OR) as suggested by veterinarian.

Presentation:
5 Liters HDPE containers.



**RR VETERINARY
HEALTH CARE PVT. LTD.**

An ISO 9001:2015 Certified Company

#4-122/1/1, Plot no. 115 & 123, Kuntloor Village, Abdullapurmet Mandal, R R Dist - 501 505, Telangana, INDIA.

For customer complaints, contact: Customer Care Executive at the above address or call +91 94904 10562.

email: feedback@rrveterinary.in, www.rrveterinary.in, www.rrvhc.in

ANTIPROL™

The Only answer to Prolapse & Intussuception




**RR VETERINARY
HEALTH CARE PVT. LTD.**

feedback@rrveterinary.in, www.rrveterinary.in, www.rrvhc.in

What is VENT PROLAPSE in Poultry?

- ♦ VENT PROLAPSE is a traumatic condition of birds, in which inflamed lower part of oviduct turns inside out and protrudes through the vent / cloaca during the egg lay period.
- ♦ Usually, the hen can retract the oviduct after egg laying. Prolapse occurs as the hen cannot retract the oviduct and the part of it remains outside the body. Vent prolapse is a serious sickness that causes low productivity in laying hens.
- ♦ Vent Prolapse is also known as PROLAPSE VENT, PROLAPSE OVIDUCT or PROLAPSE BLOW - OUT or CLOACAL PROLAPSE or PICKOUT
- ♦ Vent Prolapse is often seen in birds and is associated with cannibalism in poultry. Prolapse is also seen in oviducts of egg-laying females that strain excessively to lay due to uterine or egg-related factors.





Causes for Prolapse:

- ❖ Incorrect pre-lay management.
- ❖ Poor skeletal development during rearing, with under developed pelvic girdle.
- ❖ Nutritional changes prior to the start of egg production.
- ❖ Chickens that begin laying are too young and are under weight.
- ❖ Muscular degeneration resulting in muscle weakness of muscles responsible for egg laying.
- ❖ Tendency to lay larger eggs.
- ❖ Intestinal infections.
- ❖ Age at sexual maturity.
- ❖ Early photo-stimulation to attain sexual maturity
- ❖ Older chickens that is obese with too much fat deposition around their reproductive organs.
- ❖ Un balanced feed formulations
- ❖ On-going Calcium deficiency.
- ❖ Holding droppings for a long time, causing stress and stretching of cloaca.
- ❖ Laying of Extra Large or Jumbo or Double yolk eggs.

What is Intussusception?

- ❖ INTUSSUSCEPTION, is a condition where one section of the intestine telescopes into another, leading to intestinal obstruction and potentially death. Intussusception is a serious, life-threatening gastrointestinal emergency in chickens.
- ❖ Intussusception occurs when part of the intestine (most often the middle and distal part of the small intestine) folds or slides into the next, similar to pieces of a telescope.
- ❖ Birds with intussusception characteristically demonstrate their anterior portion of the intestine invaginated with adhesions into the posterior portion of the intestines wherein the affected areas may be congested, necrotic and gangrenous in nature.
- ❖ When Intussusception occurs, it obstructs food and fluids from passing and being digested through the intestine. When the intestine is blocked, the blood supply to the affected part gets cut off and causes damage to the intestine. If left untreated, it can result in internal bleeding and severe abdominal infection leading to death within 2-5 days.
- ❖ Intussusception occurs in thin walled lower intestines that lost muscle tone due to prevailing enteritis. Usually intussusception is predominantly seen in 7 to 10 week-old layers and pullets compared to broilers.



Causes for Intussusception:

Intussusception occurs in thin walled lower intestines that lost muscle tone due to
Prevailing enteritis,
Coccidiosis, or
Helminth worms.

Intussusception often occurs in the jejunum, the lower part of the small intestine, and is typically triggered by

- Increased intestinal motility,
- Stress / hunger
- Other forms of enteritis.

Clinical Signs of Intussusception:

- Constipation
- Not defecating
- Progressive Weight Loss
- Loss of Appetite
- Abdominal Swelling
- Blood and mucous in faeces
- Weakness