



**RR VETERINARY
HEALTH CARE PVT. LTD.**

FORMAKLEEN™

"SPECIALLY FORMULATED FORMALIN FOR POULTRY INDUSTRY"


- ✦ Disinfection with Formaldehyde is a widely used process in poultry industry especially for disinfection of poultry farms, brooder houses, hatcheries and hatchery vehicles.
- ✦ Formaldehyde is effective in reduction of contamination levels caused by bacteria, viruses and molds throughout the production process.
- ✦ Formaldehyde as a disinfectant kills bacterial spores and other key organisms, such as Salmonella, Pseudomonas, Proteus, E. coli, Staphylococcus, Streptococci and Aspergillus.
- ✦ FORMAKLEEN is a specially formulated formalin solution from RR Veterinary Health Care Private Limited, that contains added stabilizers to prevent aerial oxidation and polymerization of formalin and effectively disinfects poultry farm premises.

Composition:

Formalin 37% (wt/wt) with 20% Methanol and added stabilizers.

Key Benefist:

- ✦ FORMAKLEEN acts by cross linking of macromolecules such as proteins, RNA & DNA
- ✦ FORMAKLEEN is a bactericidal, sporicidal & virucidal disinfectant.
- ✦ Unlike chlorine based disinfectants, FORMAKLEEN does not corrode stainless steel equipment used in poultry farming.
- ✦ FORMAKLEEN is an effective disinfectant for poultry farms, hatcheries and hatchery vehicles.
- ✦ FORMAKLEEN is a properly stabilized formulation, hence it can be stored & used for longer periods up to one year without any internal polymerization.



Applications of FORMAKLEEN

- ✦ Hatchery
 - ✦ Used when fogging, spraying within a building, and in hatcheries during hatching process
- ✦ Egg fumigation
 - ✦ Used for egg and equipment fumigation, usually in a fumigation cabinet
- ✦ Breeding farm ✦ Foot bath ✦ Vehicle spray
- ✦ As an equipment disinfectant

Dosage:

For General spray : 10 ml per litre of water

For Foot Bath : 20 ml per litre of water

For Mopping of the floor : 5 ml per litre of water



Directions for Use:

For fumigation of hatching eggs and equipment:

Pre incubation:

- ✦ Mix .6 gram potassium permanganate (KMnO₄) with 1.2 ml FORMAKLEEN for each cubic foot of space
- ✦ Mix both ingredients in an earthenware or heat resistant container having at least ten times the capacity of the ingredients being added.
- ✦ Circulate the gas for 20 minutes at 70 °F

For fumigating eggs in an incubator

- ✦ Mix 0.4 grams of KMnO₄ and 0.8 ml FORMAKLEEN per ft³
- ✦ Mix both ingredients in an earthenware or heat resistant container having at least ten times the capacity of the ingredients being added.
- ✦ Circulate the gas for 20 minutes at 70 °F

Note: Do not fumigate chicken eggs between the 24th and 96th hours of incubation. Other species of birds may need the incubation intervals adjusted to compensate for total incubation time in relationship to the chicken's incubation period. It is best to incubate after the incubator reaches normal operating temperature and humidity.

Presentation: 50 kg HDPE rackets.



RR VETERINARY
HEALTH CARE PVT. LTD.

An ISO 9001:2015 Certified Company

#4-122/1/1, Plot no. 115 & 123, Kuntloor Village, Abdullapurmet Mandal,
R R Dist - 501 505, Telangana, INDIA.

For customer complaints, contact: Customer Care Executive at the above
address or call +91 94904 10562. email: feedback@rrveterinary.in,
www.rrveterinary.in, www.rrvhc.in