



RR VETERINARY
HEALTH CARE PVT. LTD.


HEP-SET-GO

“A Natural Hepatoprotectant & An
Excellent Supportive Supplement
for LITCHI in birds”



What is HPS or IBH-HPS?

- Hydro pericardium syndrome (HPS) (OR) Hydro pericardium–hepatitis syndrome is an economically important endemic disease that causes huge economic loss to poultry sector, owing to high mortality, reduced productivity and immunosuppression.
- The disease is also known as Angara disease (in Pakistan)/Litchi heart disease (in India) or Inclusion Body Hepatitis–Hydropericardium syndrome (IBH–HPS).
- HPS is an emerging immunosuppressive disease of 3–6 week old broilers, with a characteristic sudden onset with spiking mortality up to 60% but more typically 10–30%, with distinctive accumulation of fluid in pericardial sac (HYDROPERICARDIUM).
- The disease is caused by Fowl Adeno Virus (FAdV), a non-enveloped icosahedral virus belonging to the Avian Adenovirus genus of the Adenoviridae family.
- Both primary and secondary lymphoid organs are depressed significantly in case of HPS. Lymphoid organs such as spleen, thymus, bursa of Fabricius and caecal tonsil are preferentially attacked by FAdV resulting in marked immunosuppression




**Typical litchi heart appearance
at 28 hours post-infection with
discoloured and friable liver**

Transmission of the Disease:

- The disease is highly contagious, can spread quickly from one flock to another and from one farm to another by horizontal and mechanical routes of transmission and with contaminated litter.
- The FAdV is prolifically excreted through the feces of infected birds and serve as a major source of infection to young broilers.
- Horizontal transmission of the virus from bird–to–bird occurs in a flock via oro–faecal route followed by further mechanical spread of the disease along with faecal contamination.
- From one country to another a mechanism of spread may be by commercial hatching of eggs.
- Disease also gets transmitted by contact or mechanically through the poultry farm workers, vaccinators, contaminated equipment and fomites or by flies and rodents.

Disease Management:

- It is essentially important to observe good biosecurity practices for preventing HPS infection, including strict cleaning and disinfection procedures at the farm level.
 - cleaning and disinfection of premises.
 - cleaning and disinfection of equipment.
 - restriction to entry of visitors and crews involved in vaccination in the poultry houses significantly play crucial role in preventing the disease.
- Isolation of flocks accompanied by good sanitary procedures helps to prevent the spread of infection.



HEP-SET-GO



“A Natural Hepatoprotectant & An
Excellent Supportive Supplement
for LITCHI in birds”

Prevention & Treatment::

RR Veterinary Health Care Private Limited launches a unique formulation with natural bio actives, HEP-SET-GO that helps in amelioration of HPS condition in birds, controls sudden mortality and improves productivity.

Composition:

Natural Bioactives from Curcuma longa, Punica granatum, Allium sativum, Propolis, CNCQ-1F, L- Carnitine, Vitamin-D, Zinc sulphate, Selenium & Tricholine citrate.

Functional Effects of HEP-SET-GO®:

Bioactives in HEP-SET-GO® provides protection to birds from Litchi disease by various mechanisms including

- Inhibition of viral replication.
- Down regulation of pro inflammatory cytokines and inhibits the nitric oxide synthase expression in macrophages.
- Up regulating the levels of antioxidant NRF-2, and there by facilitating balanced mitochondrial functions.
- Hepatoprotection by improving fatty acid metabolism.

Key Benefits:

Bioactives in HEP-SET-GO® provides protection to birds from Litchi disease by various mechanisms including

- HEP-SET-GO® Acts as a Hepatostimulant and Hepatoprotective agent that restores liver function.
- HEP-SET-GO® helps in the management of viral induced hydropericardium syndrome and hepatopathy.
- HEP-SET-GO® is a best supportive therapy for LITCHI Disease in birds.
- HEP-SET-GO® helps in the regeneration of hepatic cells that are damaged by viral attack.
- HEP-SET-GO® helps in potentiating the Liver's Antioxidant defense mechanisms.

Directions For Use:

Treatment:

25 ml for 100 birds in drinking water for 3-5 days initially and in case of severity use 75-100 ml for 100 birds for 2-3 days and 20 ml for next 2 days thereafter.

Prevention:

15 ml for 100 birds for two days in a week. (Or) As suggested by Veterinarian



RR VETERINARY
HEALTH CARE PVT. LTD.

An ISO 9001:2015 Certified Company

#4-122/1/1, Plot no. 115 & 123, Kuntloor Village, Abdullapurmet Mandal,
R R Dist - 501 505, Telangana, INDIA.

For customer complaints, contact: Customer Care Executive at the above address or call
+91 94904 10562. email: feedback@rrveterinary.in, www.rrveterinary.in, www.rrvhc.in