

INDICATIONS:

- ❖ For the control and prevention of both Gram positive & Gram negative bacteria.
- ❖ Controls wet droppings and vent pasting
- ❖ Controls and prevents necrotic enteritis
- ❖ Maintains the colonization of native and useful bacteria in intestines

KEY BENEFITS:

- ❖ Improves growth performance by controlling infections at subclinical levels
- ❖ Ensures flock uniformity.
- ❖ Improves FCR and weight gain
- ❖ Protects gut health by controlling diarrhea of unknown etiology

DOSAGE:

Breeders:

Regular dosage: 500 gm /Ton of feed/5-7days a month

Treatment dosage: 1-1.5 kg /Ton of feed/5-7days a month

Commercial Layers & Broilers:

Regular dosage: 250 gm-500 gm /Ton of feed/5-7days a month

Treatment dosage: 500 gm-1kg /Ton of feed/5-7days a month

(or) As suggested by the Veterinarian.



Presentation:

10 kg Triple laminated BOPP bags.



RR VETERINARY
HEALTH CARE PVT. LTD.

An ISO 9001:2015 Certified Company

#4-122/1/1, Plot no. 115 & 123, Kuntloor Village, Abdullapurmet Mandal, R R Dist - 501 505, Telangana, INDIA.

For customer complaints, contact: Customer Care Executive at the above address or call +91 94904 10562.

email: feedback@rrveterinary.in, www.rrveterinary.in, www.rrvhc.in



RR VETERINARY
HEALTH CARE PVT. LTD.

LINEODOX™

Assured per Formance Enhancer



LINEODOX™

Assured performance enhancer

- ❖ The normal intestinal transit time of commercial poultry is very fast, and is only about 4-8 hours, thus serious situations can arise very quickly.
- ❖ Moreover, wet litter is associated with numerous concerns including animal welfare issues such as contact or footpad dermatitis and reduced air quality due to increased ammonia concentrations. Wet litter could also present a food safety concern as it provides an ideal environment for bacterial growth.
- ❖ **Diarrhea** can often be a complex issue as it can be caused by a variety of means including both non-pathogenic and pathogenic agents. There are a few bacteria and viruses that can cause chicken diarrhea. More common causes for diarrhea are
 - Colibacillosis,
 - Lymphoid leukosis,
 - Marek's disease.
- ❖ Additional possible causes (although less common) include:
 - Avian intestinal spirochetosis,
 - Avian tuberculosis,
 - Infectious coryza,
 - Coccidiosis (Eimeria)
 - Fowl cholera
- ❖ Thorough investigation is necessary to identify the underlying cause of the issue and to take corrective actions as soon as possible in order to return to optimal gut health.
- ❖ **LINEODOX-FS™** is a unique Anti-diarrheal cum performance enhancer that essentially combines Lincomycin, Neomycin & Doxycycline and selective herbal extracts that provide sustained and balanced growth in birds by controlling infectious diseases at subclinical level.
- ❖ **LINEODOX-FS™** is specifically active against both Gram positive, gram negative and anaerobic bacteria and effectively controls diarrhea of unknown etiology especially during initial days of laying.
- ❖ Fortified with added herbal supplements, **LINEODOX-FS™** protects the birds against mixed infections, enteritis and improves the overall performance of birds.



Composition:

- ❖ Lincomycin Hydrochloride 1 %
- ❖ Neomycin Sulphate.... 2%
- ❖ Doxycycline....2%
- ❖ Fortified with herbal extracts of Aegle marmelos, Ocimum sanctum, Pongamia glabra (SE), Curcuma longa, Emblica officinalis, Terminalia belerica, Terminalia chebula

Feature	Lincomycin Hydrochloride	Neomycin Sulfate	Doxycycline
Spectrum Activity	Gram Positive	Active against Gram-negative and many Gram-positive bacteria as well as some protozoa	Bacteriostatic activity against broad range of Gram Positive & Gram Negative strains
Mode of Action	Inhibits protein synthesis by binding to the 50 S subunits of bacterial ribosomes and prevents peptide bond formation	Inhibits bacterial protein synthesis through irreversible binding to the 30 S ribosomal subunit of susceptible bacteria	Inhibits bacterial protein synthesis by binding to the 30S ribosomal subunit.
Intestinal Absorption	Rapidly absorbed	Readily absorbed	Readily absorbed
Safety	Safe to use	Safe to use	Safe to use

Herbal Extracts	Physiological Role
Aegle marmelos	Antibacterial and Anti-diarrheal
Ocimum sanctum	Flavonoids in this extract acts Anti-diarrhoeal agents
Pongamia glabra (SE)	Active against entero-invasive bacterial strains causing bloody diarrheal episodes
Curcuma longa	Anti-diarrhoeal activity
Emblica officinalis	Reduces GIT motility.
Terminalia belerica	Stimulates reabsorption and/or exerts anti-secretory effect and decreases intestinal motility.
Terminalia chebula	Effective in treatment of diarrhoea connected with a prolapsed rectum.

