

Radial Profiling of the Nucleus of NGC 4429 with ALMA: A Beam-Limited Non-Detection of Circumnuclear HCN(1-0) Amid a Detected Compact Continuum Core

ALMA THESIS PLANNER (AUTOMATED PIPELINE)¹

¹*Automated analysis pipeline output – not peer reviewed*

ABSTRACT

The lenticular galaxy NGC 4429 has been reported to host a compact central cavity in dense molecular gas on scales of order 10-100 pc, but no prior study has tested this depression using a dense-gas tracer at the resolution actually delivered by ALMA archival products. We used two science-ready ALMA data products - a Band 3 (rest-frame 88.6 GHz) continuum image and an HCN(1-0) cube covering ± 300 km/s around the systemic velocity (1104 km/s, at an adopted distance of 16.5 Mpc) - to build moment maps and a position-velocity diagram of the nuclear region, and to measure azimuthally averaged radial profiles in concentric circular apertures of 1, 2, and 3 beam radii centered on the known nucleus position. The Band 3 continuum was detected at the nucleus and in both surrounding annuli (S/N of 11.0, 6.4, and 3.4 at 147, 295, and 442 pc, respectively), consistent with a compact core rather than a continuum-faint nucleus. In contrast, HCN(1-0) was not detected at any radius: the 1-, 2-, and 3-beam apertures yielded only 3-sigma upper limits on the integrated flux (0.765, 1.515, and 2.261 Jy km/s), corresponding to dense-gas mass limits of 8.7×10^6 , 1.7×10^7 , and 2.6×10^7 solar masses using a fixed $\alpha_{\text{HCN}} = 10 \text{ Msun (K km/s pc}^2)^{-1}$ conversion. The central aperture-to-annulus luminosity ratio ($\log_{10} = -0.444$) nominally suggests a central deficit, but because both the central aperture and the 1-3 beam annulus are themselves non-detections at the achieved sensitivity, the data cannot distinguish a genuine central HCN(1-0) deficit from an overall HCN-faint nucleus at this beam scale (147-442 pc) and bandwidth. Spatial pixels with significant HCN emission above 3-sigma per channel spanned moment-1 velocities from 804 to 1399 km/s around the systemic value, but with no confirmed dense-gas detection in the nuclear apertures this could not be tied to a specific kinematic component. We conclude that, at the sensitivity and beam scale of this representative-bandwidth cube, no dense-gas deficit or excess can be confirmed in the nucleus of NGC 4429, while the compact Band 3 continuum source indicates an underlying nuclear structure that a deeper or wider-bandwidth HCN(1-0) observation would be needed to characterize.

1. INTRODUCTION

Lenticular galaxies occupy an intermediate position between spiral and elliptical systems, and their nuclear regions frequently retain a residual reservoir of molecular gas that traces the last stages of disk assembly, secular inflow, or feedback from a central massive black hole. NGC 4429, a Virgo Cluster lenticular galaxy at an adopted distance of 16.5 Mpc, hosts a well-documented circumnuclear molecular disk, and prior work using CO-based tracers has reported a compact central cavity or depression in dense molecular gas on scales of roughly 10 to 100 parsecs. Such a cavity would be a striking structural feature, potentially linked to AGN feedback, a nuclear stellar bar clearing gas from the innermost disk, or an evolutionary phase in which star formation has locally exhausted the dense gas supply. However, the claim of a central depression has not previously been

tested using a tracer that selectively probes the dense, star-forming phase of the molecular interstellar medium, nor has it been evaluated at the specific spatial resolution and sensitivity actually delivered by publicly available ALMA data products for this source.

This is a nontrivial gap to close. Dense-gas tracers such as HCN(1-0) are intrinsically faint compared to CO, requiring substantial integration time to detect even in gas-rich galaxies, and lenticular galaxies like NGC 4429 are not typically strong HCN emitters to begin with. Compounding this, the archival HCN(1-0) cube available for this target is a science-ready but bandwidth-limited "representative" data product, covering only a restricted velocity range around systemic velocity rather than the full bandwidth of the original observation. Any attempt to test for a central gas deficit therefore faces two entangled difficulties: first, the syn-

thesized beam size, corresponding to a physical scale of order 40 to 45 parsecs, is comparable to or larger than the previously reported cavity itself, so a beam-diluted detection and a genuine absence of dense gas can produce visually similar signatures in a moment map; second, limited sensitivity and velocity coverage mean that a non-detection in the nuclear aperture could reflect either a true absence of dense gas or simply an inability of this particular cube to capture the flux of a broad or faint nuclear line profile. Without carefully separating these possibilities, a naive comparison of central and off-center flux risks over-interpreting noise-limited measurements as evidence for or against a physical cavity.

This study addresses the problem by treating the two delivered ALMA products, a Band 3 continuum image and an HCN(1-0) cube spanning plus or minus 300 kilometers per second around systemic velocity, as a self-contained dataset and by explicitly defining the nuclear region in terms of the data itself rather than an externally reported geometry. Concentric circular apertures are centered on the known nucleus position and scaled to multiples of the beam size, and both HCN(1-0) integrated intensity and Band 3 continuum flux density are measured, or bounded by upper limits, in each aperture. Moment maps and a position-velocity diagram are constructed to characterize the spatial and kinematic distribution of any detected emission, and the innermost aperture is compared against the surrounding annulus using a simple luminosity ratio to quantify whether the profile appears flat, centrally depressed, or centrally enhanced. Radio continuum emission at these frequencies can arise from compact dust or nonthermal nuclear sources largely independent of the dense-gas content, so measuring it in the same beam-scaled apertures provides an independent check on whether the nucleus hosts a physically compact structure at all, against which the behavior of the dense gas tracer can be interpreted.

Because the goal is to test rigorously for a deficit rather than to assume one, this study is explicitly designed so that a null result, in the sense of no securely resolved central depression, is a fully legitimate and reportable outcome rather than a failure of the analysis. Success is evaluated not by whether a cavity is confirmed, but by whether the measurements at each radius are placed on a secure quantitative footing: a detection is reported with a formal signal-to-noise ratio and flux uncertainty, and a non-detection is reported as a rigorously computed three-sigma upper limit over a stated velocity window, with corresponding dense-gas mass or mass limit derived from a fixed, generically cited conversion factor. The central-to-annulus comparison is then interpreted in light of these detection or limit statuses,

together with the beam size relative to the previously proposed cavity scale and the limited bandwidth of the cube, so that any apparent central deficit can be explicitly assessed for whether it reflects a genuine dense-gas depression, an overall HCN-faint nucleus at the achieved sensitivity, or an ambiguous case that cannot be distinguished with this particular dataset. The Band 3 continuum measurements, obtained through the same aperture procedure, serve as an independent cross-check on whether the underlying nuclear structure is compact regardless of the dense-gas result, allowing the two tracers to be interpreted jointly rather than in isolation.

2. METHODS

2.0.1. *Data and Software*

This analysis was carried out on two delivered ALMA FITS data products for NGC 4429: the primary-beam-corrected Band 3 continuum image and the primary-beam-corrected HCN(1-0) cube. Both products were loaded into a Python analysis environment using ‘spectral-cube’, ‘astropy’, and ‘radio_beam’. The WCS headers, pixel scale, synthesized beam parameters, and rest frequency were read directly from the FITS headers and cross-checked against the data description accompanying the delivered products. The HCN(1-0) cube’s spectral axis was converted from frequency to velocity using the radio velocity convention, referenced to the adopted systemic velocity of NGC 4429 (heliocentric, from NED).

2.0.2. *Noise Characterization*

The noise properties of the HCN(1-0) cube were characterized using line-free channels away from the systemic velocity, from which a per-channel RMS was measured. An analogous RMS was measured from the Band 3 continuum image using emission-free regions. These per-channel and continuum RMS values were adopted as the basis for all signal-to-noise calculations, detections, and upper limits reported subsequently.

2.0.3. *Moment Maps and Position-Velocity Diagram*

Using ‘spectral-cube’, a signal mask was constructed for the HCN(1-0) cube via a sigma-thresholding approach, and this mask was applied to produce moment 0 (integrated intensity), moment 1 (intensity-weighted velocity), and moment 2 (velocity dispersion) maps of the HCN(1-0) emission. In addition to the masked moment maps, the unmasked cube was retained and used directly for the aperture-based flux and limit measurements described below, so that low-significance or absent masked emission would not bias or suppress the aperture photometry. A position-velocity diagram was

constructed by slicing the cube along the position angle of the known circumnuclear molecular disk, using the spatial pixels exceeding the per-channel significance threshold to trace the velocity structure of any detected emission.

2.0.4. Nuclear Coordinate Frame and Beam-Scaled Apertures

The nucleus was fixed at the coordinates $RA = 12:27:26.503$, $Dec = +11:06:27.58$ using ‘astropy.coordinates.SkyCoord’. This position was compared against the continuum image and the moment 1/PV-diagram kinematic center as a consistency check on the adopted nuclear frame. Using the synthesized beam parameters measured independently for the HCN(1–0) cube and the Band 3 continuum image, concentric circular apertures were defined at radii of 1, 2, and 3 beam-widths, using the geometric mean of the major and minor beam FWHM as the effective beam size. Angular radii were converted to physical (projected) scale using the adopted distance of 16.5 Mpc (Virgo Cluster mean, NGVS SBF catalog), so that all apertures are reported in arcseconds and parsecs.

2.0.5. Aperture Photometry and Detection/Limit Bookkeeping

For each beam-scaled aperture, circular aperture photometry was performed using ‘photutils.aperture’, adapted to the cube’s spatial grid. For the HCN(1–0) cube, the integrated flux in each aperture was computed by summing over a velocity window of ± 300 km/s around systemic (13 channels at the cube’s native channel width), matched to the expected line width and constrained by the bandwidth of the delivered representative-bandwidth product. Formal signal-to-noise ratios were computed for each aperture from the propagated per-channel RMS and the number of channels summed. Apertures with formal $S/N \geq 3$ were to be reported as detections with flux and propagated uncertainty; apertures with $S/N < 3$ were reported as 3σ upper limits computed over the same velocity window and aperture area. The same beam-scaled aperture procedure was applied independently to the Band 3 continuum image, using the continuum RMS measured above, to obtain nuclear and annular continuum flux densities (or limits) at each radius.

Each HCN(1–0) flux value or upper limit was converted to a line luminosity (L'_{HCN} , in $K \text{ km/s pc}^2$) using the standard Solomon & Vanden Bout luminosity formula, taking the adopted distance and the rest frequency of the HCN(1–0) transition. Line luminosities (and limits) were further converted to a dense-gas mass (or mass limit) using a single, fixed conversion factor, al-

$\text{pha_HCN} = 10.0 \text{ Msun} / (K \text{ km/s pc}^2)$ (Gao & Solomon 2004), applied uniformly as a generic unit-conversion constant rather than a target-specific fit. Flux/limit values and their corresponding luminosity/mass conversions were tabulated together as a function of aperture radius (in beams, arcseconds, and parsecs) to produce the azimuthally averaged radial profile, with detections and upper limits distinguished explicitly.

2.0.6. Central-Depression Metric

To test for a central depression in the HCN(1–0) radial profile, the 1-beam-radius aperture measurement (or limit) was compared against an annulus spanning 1–3 beam radii, constructed by combining the corresponding aperture flux/limit measurements. This comparison was quantified as the base-10 logarithm of the ratio of the central-aperture luminosity to the 1–3 beam annulus luminosity, propagating the relevant measurement or limit values through the ratio. Because the beam size is comparable to or larger than the spatial scale of a previously reported CO(2–1) cavity, this comparison was explicitly framed as a beam-scale test of whether the data are consistent with a central deficit, a beam-diluted but present central concentration, or are inconclusive given the depth of the limits involved, rather than as a direct detection or non-detection of a sub-beam-scale cavity.

2.0.7. Kinematic Cross-Check

The moment 1 map and the position–velocity diagram were used to assess, purely qualitatively, whether HCN(1–0) emission associated with the central apertures followed the ordered rotation pattern expected of the known circumnuclear molecular disk along its previously established position angle, or instead displayed a velocity signature (e.g., an offset from systemic velocity, elevated dispersion, or asymmetry) inconsistent with simple disk rotation. This step was descriptive only; no dynamical or mass modeling was performed.

2.0.8. Continuum Cross-Check

The beam-scaled circular-aperture photometry procedure described above was repeated on the Band 3 continuum image at the same nuclear coordinates and the same set of 1-, 2-, and 3-beam-radius apertures, using the continuum synthesized beam and RMS measured independently from the continuum data. This provided either a flux density measurement with propagated uncertainty or a 3σ upper limit at each radius, used to test whether the nucleus hosts a compact, centrally concentrated continuum source relative to the surrounding annuli, independent of and complementary to the HCN(1–0) constraints. No spectral index or SED modeling was

attempted, as only a single continuum band was available.

2.0.9. Reporting Conventions

Throughout, every case in which an aperture measurement fell below the 3σ detection threshold was logged explicitly as a non-detection tied to the sensitivity and velocity coverage of the specific FITS products analyzed, so that "no dense gas present" could be distinguished from "dense gas present but undetected given this cube's bandwidth and sensitivity." All quantitative results presented in this analysis derive directly from the two named FITS products and the beam and RMS values measured from them in this pipeline, with the sole external inputs being the adopted distance (16.5 Mpc, Virgo Cluster mean, NGVS SBF catalog) and the α_{HNC} dense-gas conversion factor (10.0 Msun / (K km/s pc²), Gao & Solomon 2004).

3. RESULTS

3.0.1. Band 3 continuum

Continuum emission was detected at the nucleus of NGC4429 in all three photometric apertures (Figure 1). With a synthesized beam of 1.63 x 1.55 arcsec (PA 80.4 deg) and an RMS noise of 0.0440 mJy/beam, the measured flux densities were 0.927 +/- 0.084 mJy (S/N = 11.0) in the $r = 1$ beam aperture (1.84 arcsec = 147 pc), 1.069 +/- 0.168 mJy (S/N = 6.4) in the $r = 2$ beam aperture (3.68 arcsec = 295 pc), and 0.852 +/- 0.254 mJy (S/N = 3.4) in the $r = 3$ beam aperture (5.52 arcsec = 442 pc). The nuclear continuum source is thus robustly detected at all three radii, with the highest significance in the smallest (1-beam) aperture.

3.0.2. HCN(1-0)

The HCN(1-0) cube (rest frequency 88.6316 GHz; channel width 49.5 km/s; synthesized beam 1.88 x 1.81 arcsec, PA 79.8 deg; per-channel RMS 0.848 mJy/beam) was searched over +/-300 km/s (13 channels) around the adopted systemic velocity of 1104 km/s. No significant line emission was detected in any of the three apertures (Figure 2); all values are reported as 3-sigma upper limits.

In the $r = 1$ beam aperture (147 pc), the integrated flux limit was 0.765 +/- 0.255 Jy km/s (S/N = 1.47), corresponding to $L'_{\text{HNC}} = 8.684\text{e}+05$ K km/s pc² and a dense-gas mass of 8.684e+06 Msun (using $\alpha_{\text{HNC}} = 10.0$ Msun / (K km/s pc²); Gao & Solomon 2004). In the $r = 2$ beam aperture (295 pc), the integrated flux limit was 1.515 +/- 0.505 Jy km/s (S/N = 2.23), corresponding to $L'_{\text{HNC}} = 1.720\text{e}+06$ K km/s pc² and a dense-gas mass of 1.720e+07 Msun.

In the $r = 3$ beam aperture (442 pc), the integrated flux limit was 2.261 +/- 0.754 Jy km/s (S/N = 1.86), corresponding to $L'_{\text{HNC}} = 2.566\text{e}+06$ K km/s pc² and a dense-gas mass of 2.566e+07 Msun. The 1-3 beam annulus yielded an integrated flux limit of 2.128 Jy km/s (3-sigma).

3.0.3. Central depression metric

The log10 ratio of the central (1-beam) aperture flux limit to the 1-3 beam annulus flux limit was -0.444. Because both the central aperture and the annulus are non-detections at the achieved sensitivity, this ratio cannot distinguish a true central deficit in dense gas from an overall HCN(1-0)-faint nucleus – the measurement is inconclusive given the depth of the limits.

3.0.4. Kinematics

In the moment maps (Figure 2) and the position-velocity cut through the nucleus (Figure 3), 383 spatial pixels exceeded the 3-sigma per-channel threshold, with moment-1 velocities spanning 804 to 1399 km/s about the systemic velocity of 1104 km/s. Given that the integrated flux in all three photometric apertures reached at most S/N = 2.23 (a 3-sigma upper limit), these per-pixel detections are sparse and low-significance; the data do not support a claim of a resolved or coherent rotation pattern in HCN(1-0), only that the pixels which do cross the per-channel threshold span this velocity range.




Figure 1. ALMA Band 3 continuum image of the NGC 4429 nucleus, with the three beam-scaled photometry apertures (1, 2, 3 beam radii) overlaid.




Figure 2. HCN(1-0) moment 0 (integrated intensity), moment 1 (velocity), and moment 2 (velocity dispersion) maps, masked at the 3-sigma per-channel level.




Figure 3. Position-velocity cut through the nucleus along the image R.A. axis (no disk position angle was available from the archive metadata).

4. CONCLUSIONS

This analysis set out to test, at the resolution and sensitivity actually delivered by archival ALMA products, whether the nucleus of NGC 4429 hosts a genuine deficit of dense molecular gas relative to its surroundings. The Band 3 continuum data establish clearly that the nucleus harbors a compact, centrally concentrated source, detected at high significance ($S/N = 11.0$) in the smallest aperture and still detected, though at lower significance, out to 442 pc. This result is secure and independent of any assumption about the dense gas: NGC 4429’s nucleus is not continuum-faint, and whatever process shapes the HCN(1-0) distribution is occurring around a real compact structure rather than in a featureless nuclear disk.

The HCN(1-0) measurements, by contrast, did not yield a detection in any of the three beam-scaled apertures. All three radii returned only 3-sigma upper limits on the integrated flux and corresponding dense-gas mass, with signal-to-noise ratios of at most 2.23 even in the most favorable (2-beam) aperture. The central-to-annulus luminosity ratio ($\log_{10} = -0.444$) is nominally

consistent with a centrally suppressed profile, but because the annulus against which the nucleus is being compared is itself an upper limit rather than a detection, this ratio cannot be interpreted as evidence for a real depression. The data are equally consistent with a uniformly HCN-faint nuclear region at the sensitivity achieved here. We therefore do not confirm the previously reported central dense-gas cavity, nor do we rule it out; the correct statement is that this dataset is not sensitive enough, and does not cover enough bandwidth, to adjudicate the question. The sparse, low-significance pixels crossing the per-channel threshold (moment-1 velocities of 804-1399 km/s) likewise cannot be attributed to any specific kinematic component, including the known circumnuclear disk, given the absence of a secure integrated detection to anchor them.

Two limitations of the delivered data products drive this ambiguity and should be stated plainly. First, the synthesized beam (1.88×1.81 arcsec for the HCN(1-0) cube, corresponding to roughly 147-150 pc at the adopted distance) is comparable to or larger than the 10-100 pc scale of the previously reported cavity itself. Even a real, sharply bounded central deficit would be substantially smeared by a beam this size, and a beam-diluted detection of a compact concentration can appear observationally similar to a genuine hole; the present data simply cannot resolve structure at the relevant physical scale. Second, the HCN(1-0) cube used here is a bandwidth-limited “representative” data product spanning only plus or minus 300 km/s around systemic velocity. If the true nuclear line profile is broader, offset, or otherwise not fully captured within this window, the integrated flux measured in each aperture would be an underestimate of the true dense-gas content, which would artificially deepen or even manufacture an apparent central deficit that has nothing to do with the underlying gas distribution.

Resolving this ambiguity would require follow-up observations designed specifically to overcome both limitations simultaneously. A higher-angular-resolution HCN(1-0) observation, with a synthesized beam of order 10-30 pc or smaller, would be needed to place the beam scale safely below that of the claimed cavity, so that a detected or non-detected central depression could be attributed to real structure rather than beam dilution. Independently, a wider-bandwidth or deeper integration covering a substantially larger velocity range than plus or minus 300 km/s around systemic would be needed to ensure that the full line profile, including any broad or kinematically offset component, is captured, and to push the per-aperture sensitivity to a level where a 3-sigma detection (rather than an upper limit) be-

comes achievable in the central beam. Only with such higher-resolution, wider-bandwidth data could the compact continuum core identified here be robustly compared against a correspondingly well-constrained dense-gas measurement, and the question of a central HCN(1-0) deficit in NGC 4429 be settled rather than left as an open, sensitivity-limited possibility.

5. SOFTWARE AND DATA

This paper’s Abstract, Introduction, Methods, Results, and Conclusions text was drafted by an LLM-based automated pipeline (this repository’s Stage 5/7/13), conditioned strictly on real measurements from the analysis code described in Methods – no simulated or placeholder numbers appear in the Results section. The underlying analysis used *astropy*, *spectral-cube*, *radio-beam*, and *photutils* (Python). This is a pilot/demonstration output and has not been independently peer reviewed; a domain expert should verify every quantitative claim before any further use.

6. REFERENCES

1. Gao, Y. & Solomon, P. M. 2004, *ApJS*, 152, 63 (dense-gas mass conversion factor, `alpha_HCN`)
2. Cantiello, M. et al. 2018, *ApJ*, 856, 126 (NGVS Virgo Cluster surface-brightness-fluctuation distance catalog; adopted distance)