

```
{
  "pageno": 0,
  "type": "StrikeOut",
  "message": {
    "comment": "delete",
    "responses": [
      "reply",
      "another reply"
    ]
  },
  "selection": "stand the failure of the \"Flat Orbit Conjecture\" in higher curvature regimes<StrikeOut>.</StrikeOut>"
}

```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Caret",
  "message": {
    "comment": ". <period: put foootnote after period>",
    "responses": []
  },
  "selection": "stand the failure of the \"Flat Orbit Conjecture\" in higher curvature regimes<Caret></Caret>1."
}
}
```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes. A B

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. A F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Replace",
  "message": {
    "comment": "Flat orbit conjecture",
    "responses": []
  },
  "selection": "stand the failure of the <Replace>\"Flat Orbit Conjecture\"</Replace> in
higher curvature regimes1."
}
}
```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "StrikeOut",
  "message": {
    "comment": "<close up space>",
    "responses": []
  },
  "selection": "function is trivially constant when X is Euclidean, but varies dramatically in non<StrikeOut>-</StrikeOut>"
}
}
```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```

{
  "pageno": 0,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "StabG(x) = {g ∈ G | g · x = x}<StrikeOut>, </StrikeOut>"
}

```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim_{[H:K]}^{\lambda^2}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\}$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Replace",
  "message": {
    "comment": "765489",
    "responses": []
  },
  "selection": "L. H. was supported by NSF grant No. <Replace>314159357</Replace>. F. E. thanks the Department of Linguistics"
}
}
```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Replace",
  "message": {
    "comment": "The first author",
    "responses": []
  },
  "selection": "<Replace>L. H.</Replace> was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics"
}
}
```

# LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

## 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

## 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Highlight",
  "message": {
    "comment": "please move to Acknowledgments section.",
    "responses": []
  },
  "selection": "<Highlight>L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics</Highlight>"
}
}
```

## LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

### 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

### 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 0,
  "type": "Replace",
  "message": {
    "comment": "p <small cap>",
    "responses": []
  },
  "selection": "2. Background and <Replace>P</Replace>reliminaries"
}
```

# LOW ORBIT FOLIATIONS OF CAT(0)

LEROY HUBBARD AND FRANCIS EULER

ABSTRACT. We set  $\mathcal{G} = \sim \frac{\lambda^2}{[H:K]}$  and investigate the orbits of  $\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  provided  $\lambda \in [1 - \varphi, 1 + \varphi]$ , where  $\varphi$  is the golden ratio. Here we provide a novel method for verifying the characteristics of the orbits of  $\mathfrak{I}$ .

## 1. INTRODUCTION

Ever since 1689 with Fermat's treatise on prime enumeration [1], attempts at understanding  $\frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}}$  have been underway but mostly unsuccessful. Our main objective is to describe the low-orbit foliations induced by  $\mathfrak{I}$  on the pseudo-Euclidean completion of a CAT(0) complex. This perspective arose from the need to understand the failure of the “Flat Orbit Conjecture” in higher curvature regimes<sup>1</sup>.

## 2. BACKGROUND AND PRELIMINARIES

Let  $(X, d)$  be a CAT(0) space in the sense of Gromov. For a fixed  $\lambda > 0$ , define the *low orbit foliation*  $\mathcal{F}_\lambda(X)$  as

$$(1) \quad \mathcal{F}_\lambda(X) = \{x \in X \mid \Delta(x, \lambda x) = \text{const.}\},$$

where  $\Delta(x, \lambda x) = d(x, \lambda x)$  denotes the displacement function under  $\lambda$ -scaling. This function is trivially constant when  $X$  is Euclidean, but varies dramatically in non-flat CAT(0) manifolds.

**2.1. A remark on  $\mathcal{G}$ -stabilizers.** We shall repeatedly use the stabilizer group

$$\text{Stab}_{\mathcal{G}}(x) = \{g \in \mathcal{G} \mid g \cdot x = x\},$$

whose index  $[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]$  determines the *orbit density* at  $x$ . In general, we have

$$(2) \quad \rho(x) = \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} \cdot \exp(-\kappa(x)),$$

where  $\kappa(x)$  denotes the local curvature contribution, computed by a modified Ricci form.

Equation (2) implies that low orbit foliations are sensitive to curvature fluctuations, as illustrated in Figure 1.

---

L. H. was supported by NSF grant No. 314159357. F. E. thanks the Department of Linguistics for the valuable conversations.

<sup>1</sup>Originally conjectured by P. Alexandrov, the Flat Orbit Conjecture proposed that all  $\lambda$ -periodic orbits of a CAT(0) space are isometric to Euclidean circles. This is now known to be false in dimensions  $\geq 3$  due to [2].

```
{
  "pageno": 1,
  "type": "Replace",
  "message": {
    "comment": "; e",
    "responses": [
      "a reply to a replace annotation"
    ]
  },
  "selection": "Figure 2. Low orbit foliations centered at x0<Replace>. E</Replace>ach
ellipse repre-"
}
}
```

2

LEROY HUBBARD AND FRANCIS EULER




FIGURE 1. A schematic of local orbit curvature under  $\lambda$ -perturbation.

### 3. MAIN RESULTS

Our principal theorem relates the orbit structure of  $\mathfrak{I}$  to the golden window of  $\lambda$ :

**Theorem 3.1.** *Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform if and only if*

$$(3) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

*Proof.* We proceed by expanding  $\mathfrak{I}$  as a quotient operator:

$$\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}} = \text{CAT}(0) \otimes \mathcal{G}^{-\lambda k}.$$

Substituting into the geometric mean inequality and integrating over  $X$  yields

$$\int_X \rho(x) d\mu(x) = \int_X \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} e^{-\kappa(x)} d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi},$$

after simplification via the  $\varphi$ -symmetric normalization lemma (see Appendix ??).  $\square$

**Corollary 3.2.** *If  $\lambda = 1$ , then  $\mathcal{F}_1(X)$  coincides with the canonical horospherical foliation of  $X$ .*




FIGURE 2. Low orbit foliations centered at  $x_0$ . Each ellipse represents an orbit of constant  $\Delta(x, \lambda)$ .

```
{
  "pageno": 1,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "sents an orbit of constant  $\Delta(x, \lambda)$  LEROY HUBBARD AND FRANCIS EULER"
}
}
```




FIGURE 1. A schematic of local orbit curvature under  $\lambda$ -perturbation.

### 3. MAIN RESULTS

Our principal theorem relates the orbit structure of  $\mathfrak{I}$  to the golden window of  $\lambda$ :

**Theorem 3.1.** *Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform if and only if*

$$(3) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

*Proof.* We proceed by expanding  $\mathfrak{I}$  as a quotient operator:

$$\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}} = \text{CAT}(0) \otimes \mathcal{G}^{-\lambda k}.$$

Substituting into the geometric mean inequality and integrating over  $X$  yields

$$\int_X \rho(x) d\mu(x) = \int_X \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} e^{-\kappa(x)} d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi},$$

after simplification via the  $\varphi$ -symmetric normalization lemma (see Appendix ??).  $\square$

**Corollary 3.2.** *If  $\lambda = 1$ , then  $\mathcal{F}_1(X)$  coincides with the canonical horospherical foliation of  $X$ .*




FIGURE 2. Low orbit foliations centered at  $x_0$ . Each ellipse represents an orbit of constant  $\Delta(x, \lambda)$ .

```
{
  "pageno": 1,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "Figure 1. A schematic of local orbit curvature under LEROY HUBBARD AND FRANCIS EULER  
-perturbation<StrikeOut>. </StrikeOut>"}
}
```




FIGURE 1. A schematic of local orbit curvature under  $\lambda$ -perturbation.

### 3. MAIN RESULTS

Our principal theorem relates the orbit structure of  $\mathfrak{I}$  to the golden window of  $\lambda$ :

**Theorem 3.1.** *Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform if and only if*

$$(3) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

*Proof.* We proceed by expanding  $\mathfrak{I}$  as a quotient operator:

$$\mathfrak{I} = \frac{\text{CAT}(0)}{\mathcal{G}^{\lambda k}} = \text{CAT}(0) \otimes \mathcal{G}^{-\lambda k}.$$

Substituting into the geometric mean inequality and integrating over  $X$  yields

$$\int_X \rho(x) d\mu(x) = \int_X \frac{1}{[\mathcal{G} : \text{Stab}_{\mathcal{G}}(x)]} e^{-\kappa(x)} d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi},$$

after simplification via the  $\varphi$ -symmetric normalization lemma (see Appendix ??).  $\square$

**Corollary 3.2.** *If  $\lambda = 1$ , then  $\mathcal{F}_1(X)$  coincides with the canonical horospherical foliation of  $X$ .*




FIGURE 2. Low orbit foliations centered at  $x_0$ . Each ellipse represents an orbit of constant  $\Delta(x, \lambda)$ .

```

{
  "pageno": 2,
  "type": "Replace",
  "message": {
    "comment": "if and only if",
    "responses": []
  },
  "selection": "Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform <Replace>if</Replace>3  

  LOW ORBIT FOLIATIONS OF CAT(0)
}

```

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform [P]

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

+The proof is omitted for space reasons; see Appendix B.+

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in [Julia 1.10](#) to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "<StrikeOut>(</StrikeOut>The proof is omitted for space reasons; see  
LOW ORBIT FOLIATIONS OF CAT(0)  
Appendix B.)"
}

```

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

The proof is omitted for space reasons; see Appendix B. ↗

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in [Julia 1.10](#) to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "(The proof is omitted for space reasons; see Appendix B.<StrikeOut>)</StrikeOut>"
}

```

LOW ORBIT FOLIATIONS OF CAT(0)

3

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

(The proof is omitted for space reasons; see Appendix B.)

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in [Julia 1.10](#) to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "Highlight",
  "message": {
    "comment": "COMP: set typewriter font <\\texttt{ }>",
    "responses": []
  },
  "selection": "We implemented a simple prototype in Julia 1.10 to visualize  $\mathcal{F}_\lambda(X)$  for synthetic"
}

```

LOW ORBIT FOLIATIONS OF CAT(0)  
3

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

(The proof is omitted for space reasons; see Appendix B.)

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in **Julia** 1.10 to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "Highlight",
  "message": {
    "comment": "small cap",
    "responses": []
  },
  "selection": "5. Numerical <Highlight>Experiments</Highlight> LOW ORBIT FOLIATIONS OF CAT(0)"
}
```

3

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

(The proof is omitted for space reasons; see Appendix B.)

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in **Julia 1.10** to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "Highlight",
  "message": {
    "comment": "lowercase",
    "responses": []
  },
  "selection": "4.1. Numerical Simulation. Following [3], we can simulate the orbit structure"
}

```

LOW ORBIT FOLIATIONS OF CAT(0)  
3

#### 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical **S**imulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

(The proof is omitted for space reasons; see Appendix B.)

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$

which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

#### 5. NUMERICAL **E**XPERIMENTS

We implemented a simple prototype in **Julia 1.10** to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$

```
{
  "pageno": 2,
  "type": "Highlight",
  "message": {
    "comment": "lowercase",
    "responses": []
  },
  "selection": "4. Applications and Examples"
}
```

3

## 4. APPLICATIONS AND EXAMPLES

Consider  $X = \mathbb{H}^2$ , the hyperbolic plane. The displacement  $\Delta(x, \lambda)$  satisfies

$$\cosh \Delta(x, \lambda) = 1 + \frac{\lambda^2}{2} \|x\|^2.$$

Thus  $\mathcal{F}_\lambda(X)$  forms a family of equidistant hyperbolae, asymptotically orthogonal to geodesic boundaries.

**4.1. Numerical Simulation.** Following [3], we can simulate the orbit structure numerically. Let  $x_0 = (0, 0)$  and iterate

$$x_{n+1} = \lambda R(x_n), \quad R(x) = \frac{x}{1 + \|x\|^2},$$

to approximate the fixed points of  $\mathcal{F}_\lambda$ . Convergence occurs for  $\lambda < \sqrt{\varphi}$ .




FIGURE 3. Stable orbits obtained under  $\lambda$ -iteration.

**Theorem 4.3.** Let  $(X, d)$  be a complete CAT(0) space and  $\lambda \in [1 - \varphi, 1 + \varphi]$ . Then the orbit foliation  $\mathcal{F}_\lambda(X)$  is quasi-uniform iff

$$(4) \quad \int_X \rho(x) d\mu(x) = \frac{\lambda^2}{1 + \lambda\varphi}.$$

(The proof is omitted for space reasons; see Appendix B.)

**4.2. Curvature sensitivity.** A quick computation shows that the variance of  $\rho$  satisfies

$$(5) \quad \text{Var}(\rho) = \int_X (\rho(x) - \bar{\rho})^2 d\mu(x) = \frac{\lambda^3 - 1}{2 + \lambda^2},$$


which vanishes only when  $\lambda = 1$ . This implies that even minor perturbations from the Euclidean limit result in exponential orbit divergence.

## 5. NUMERICAL EXPERIMENTS

We implemented a simple prototype in [Julia 1.10](#) to visualize  $\mathcal{F}_\lambda(X)$  for synthetic CAT(0) surfaces generated by random triangulations. Let  $\lambda = 1.3$  and  $X$  be a simplicial complex with edge weights following a truncated Gaussian distribution  $\mathcal{N}(0.8, 0.05)$ .

After  $N = 10^4$  iterations, the mean displacement converged to

$$\bar{\Delta} = 1.274 \pm 0.006,$$


FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.

FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Replace",
  "message": {
    "comment": "Figure",
    "responses": []
  },
  "selection": "A peculiar observation (Replace Fig. Replace 5) was that for large  $\lambda$ , the orbit clusters exhibited a"
}
LEROY HUBBARD AND FRANCIS EULER
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Highlight",
  "message": {
    "comment": "put footnote after period",
    "responses": []
  },
  "selection": "double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian
systems<Highlight>2.</Highlight>"4 LEROY HUBBARD AND FRANCIS EULER
}
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Replace",
  "message": {
    "comment": "Equation",
    "responses": []
  },
  "selection": "geometric invariant for the foliation type. However, \n4LEROY HUBBARD AND FRANCIS EULER\n<Replace>Eq.</Replace> (7) reveals an unexpected"
}
}
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Replace",
  "message": {
    "comment": "(5) <pls link>",
    "responses": []
  },
  "selection": "geometric invariant for the foliation type. However, Eq. \n4 LEROY HUBBARD AND FRANCIS EULER\n<Replace>(7)</Replace> reveals an unexpected"
}
}
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to flip sign, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Highlight",
  "message": {
    "comment": "<link>",
    "responses": []
  },
  "selection": "We hypothesize (Hypothesis 5.1</Highlight>) that this anomaly corresponds to a hidden"
}
4 LEROY HUBBARD AND FRANCIS EULER
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Highlight",
  "message": {
    "comment": "remove new paragraph: continue the last paragraph with this sentence.",
    "responses": []
  },
  "selection": "4LEROY HUBBARD AND FRANCIS EULER We hypothesize</Highlight> (Hypothesis 5.1) that this anomaly corresponds to a hidden"
}
}
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize [(Hypothesis 5.1)] that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Replace",
  "message": {
    "comment": "; at",
    "responses": []
  },
  "selection": "geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected\nthat point, the curvature-weighted orbit integral" = 2
  .2.618<Replace>iatio\n. At</Replace>n type. However, Eq. (7) reveals an unexpected\nthat point, the curvature-weighted orbit integral"
}

```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Highlight",
  "message": {
    "comment": "does this look right?",
    "responses": []
  },
  "selection": "geometric \nresonance <Highlight>invariant for t\nnear = 2
  LEROY HUBBARD AND FRANCIS EULER
  .</Highlight>he foliation type. However, Eq. (7) reveals an unexpected\n2.618. At that
  point, the curvature-weighted orbit integral"
}

```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to flip sign, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```
{
  "pageno": 3,
  "type": "Highlight",
  "message": {
    "comment": "AU: is this necessary?",
    "responses": []
  },
  "selection": "<Highlight>2A referee pointed out that this might be a discretization artifact, but we were unable to</Highlight>"4
  LEROY HUBBARD AND FRANCIS EULER
}
}
```




FIGURE 4. Variance of orbit density  $\rho$  as a function of  $\lambda$ .

while the empirical curvature parameter  $\kappa$  stabilized near  $-0.218$ . The results are summarized in Table 1.

$\lambda$	$\Delta$	$\kappa$
0.9	0.913	-0.054
1.0	1.000	0.000
1.3	1.274	-0.218
1.6	1.589	-0.403

TABLE 1. Empirical orbit metrics under  $\lambda$ -iteration.

A peculiar observation (Fig. 5) was that for large  $\lambda$ , the orbit clusters exhibited a double-lobed structure reminiscent of quasi-periodic tori in Hamiltonian systems<sup>2</sup>.




FIGURE 5. Scatter of simulated orbit centers for  $\lambda = 1.6$ .

## 6. DISCUSSION AND FURTHER WORK

Our experiments confirm that the function  $\psi(\lambda) = \lambda^2/(1 + \lambda\varphi)$  behaves as a geometric invariant for the foliation type. However, Eq. (7) reveals an unexpected resonance near  $\lambda = \varphi^2 \approx 2.618$ . At that point, the curvature-weighted orbit integral appears to *flip sign*, leading to a chaotic drift that violates the CAT(0) inequality in the discrete setting.

We hypothesize (Hypothesis 5.1) that this anomaly corresponds to a hidden symmetry in the  $\mathcal{G}$ -action:

$$g \mapsto \frac{1}{\lambda} g^{-1} \lambda,$$

---

<sup>2</sup>A referee pointed out that this might be a discretization artifact, but we were unable to reproduce it analytically.

```

{
  "pageno": 4,
  "type": "Highlight",
  "message": {
    "comment": "place footnote after period",
    "responses": []
  },
  "selection": "will be discussed in a forthcoming note by the first
author<Highlight>3.</Highlight>" LOW ORBIT FOLIATIONS OF CAT(0) 5
}

```

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author.<sup>3</sup>

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi),$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

```

"pageno": 4,
"type": "Highlight",
"message": {
    "comment": "place footnote after comma",
    "responses": []
},
"selection": "Although the convergence of this expansion is
questionable<Highlight>4,</Highlight> the leading term suf-"
}

```

5

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author<sup>3</sup>.

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi),$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

```

{
  "pageno": 4,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": " $\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi)$ " LOW ORBIT FOLIATIONS OF CAT(0)
}

```

5

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author<sup>3</sup>.

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi).$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

```

{
  "pageno": 4,
  "type": "Highlight",
  "message": {
    "comment": "<link>",
    "responses": []
  },
  "selection": "fices to justify Theorem 4.3.</Highlight>" LOW ORBIT FOLIATIONS OF CAT(0) 5
}

```

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author<sup>3</sup>.

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi),$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

```

{
  "pageno": 4,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "Laplace-Beltrami operator and applying the  $\varphi$ -orthogonality condition<StrikeOut>,</StrikeOut>" LOW ORBIT FOLIATIONS OF CAT(0) 5
}

```

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author<sup>3</sup>.

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition<sup>4</sup>,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi),$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

```

{
  "pageno": 4,
  "type": "StrikeOut",
  "message": {
    "comment": "",
    "responses": []
  },
  "selection": "Figure 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ <StrikeOut>. </StrikeOut>"  

  "text": "LOW ORBIT FOLIATIONS OF CAT(0)"  

}
}

```

which has order two when  $\lambda = \varphi^2$ . The numerical confirmation of this phenomenon will be discussed in a forthcoming note by the first author<sup>3</sup>.

**6.1. Error analysis and convergence.** While most trajectories converged in under  $10^3$  iterations, approximately 2.7% diverged, displaying quasi-helical wandering. We suspect this results from non-uniform floating point rounding in the  $\mathbb{R}^3$  embedding; correcting to arbitrary precision reduces the effect but does not eliminate it entirely.




FIGURE 6. Convergence of displacement difference  $\|\Delta_n - \Delta_{n-1}\|$ .

## 7. APPENDIX B: PROOF SKETCH OF THEOREM 4.3

The argument proceeds by constructing a pseudo-measure  $\nu$  such that

$$d\nu = e^{-\kappa(x)} d\mu(x),$$

then integrating  $\rho$  against  $\nu$  over  $X$ . By expanding  $\rho$  in the eigenbasis of the Laplace–Beltrami operator and applying the  $\varphi$ -orthogonality condition,

$$\langle f_i, f_j \rangle_\varphi = \delta_{ij}(1 + \lambda\varphi),$$

we recover Eq. (5). The rest follows by applying a truncated version of Jensen’s inequality to the quotient  $\mathfrak{I}$  operator:

$$\text{CAT}(0)/\mathcal{G}^{\lambda k} \approx \text{CAT}(0)(1 - \lambda k + O(k^2)).$$

Although the convergence of this expansion is questionable<sup>4</sup>, the leading term suffices to justify Theorem 4.3.

**Acknowledgements.** The authors thank the anonymous reviewers for their sharp-eyed corrections, especially for pointing out a missing minus sign in Eq. (3), which has since been *mostly* fixed.

## REFERENCES

- [1] P. Fermat, *On prime enumeration and spatial convexity*, Toulouse Notes, 1689.
- [2] L. Hubbard, *Counterexamples to the flat orbit conjecture*, Ann. Quad. Math. (2023), 13–57.
- [3] F. Euler, *Iterative dynamics in nonpositively curved complexes*, Proc. Geom. Dyn. (2024), 211–230.
- [4] B. Zelinsky, *On modular eigenmodes of golden-ratio systems*, J. Nonlin. Struct. (2019), 98–114.

<sup>3</sup>Submitted to the *Journal of Approximate Topologies*, 2025.

<sup>4</sup>We observed divergence for  $|\lambda| > 2.1$ , which we did not pursue.

{  
    "pageno": 5,  
    "type": "Caret",  
    "message": {  
        "comment": ",",  
        "responses": []  
    },  
    "selection": "Department of Quadratics, University of Belarus, 3 Corporal Way,  
    Genevive<Caret></Caret> 06578,"  
}  
DEPARTMENT OF QUADRATICS, UNIVERSITY OF BELARUS, 3 CORPORAL WAY, GENEVIVE 06578  
BELARUS

*Email address:* lhubard@qbel.a.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGETOWN UNIVERSITY, 301 PROSPECT  
CIRCLE, WASHINGTON 12765, USA  
*Email address:* feuler@gtown.edu

```
{  
  "pageno": 5,  
  "type": "Caret",  
  "message": {  
    "comment": "",  
    "responses": []  
  },  
  "selection": "Circle, Washington6</Caret> 12765, USA"  
  LEROY HUBBARD AND FRANCÍS EULER  
}
```

DEPARTMENT OF QUADRATICS, UNIVERSITY OF BELARUS, 3 CORPORAL WAY, GENEVIVE 06578,  
BELARUS

*Email address:* lhubard@qbel.a.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGETOWN UNIVERSITY, 301 PROSPECT  
CIRCLE, WASHINGTON 12765, USA

*Email address:* feuler@gtown.edu

```
{  
    "pageno": 5,  
    "type": "Caret",  
    "message": {  
        "comment": "; and Department of Mathematics, Princeton University, Nassau Street,  
Princeton New Jersey, 02854",  
        "responses": []  
    },  
    "selection": "Circle, Washington 12765, USA<Caret></Caret>"  
}  
  
6 LEROY HUBBARD AND FRANCIS EULER  
DEPARTMENT OF QUADRATICS, UNIVERSITY OF BELARUS, 3 CORPORAL WAY, GENEVIVE 06578,  
BELARUS  
Email address: lhubard@qbel.a.edu  
  
DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGETOWN UNIVERSITY, 301 PROSPECT  
CIRCLE, WASHINGTON 12765, USA  
Email address: feuler@gtown.edu
```