

Date Sets

collection of well defined and distinct objects.

Order does not matter in set.
can be unordered.

$a \in A \longrightarrow$ a is an element of set A .
 $\downarrow$ set name

Descriptive :

Tabular :

set of first five natural number.

$$A = \{1, 2, 3, 4, 5\}$$

Set Builder Notation :

$$A = \{x \in \mathbb{N}, 1 \leq x \leq 5\} / \{x \in \mathbb{N}, 1 \leq x < 6\}$$

$$\mathbb{N} = \{0, 1, 2, \dots\} \text{ (Natural no.)}$$

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, \dots\} \text{ (Integer)}$$

$$\mathbb{Z}^+ = \{1, 2, 3, \dots\} \text{ (+ve Integer)}$$

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p \in \mathbb{Z}, q \in \mathbb{Z}, q \neq 0 \right\} \text{ (rational no.)}$$

$$\mathbb{R} = \text{(real no.)}$$

$$\mathbb{R}^+ = \text{+ve real no.}$$

$$\mathbb{C} = \text{complex no (Real + imaginary.)}$$

Real numbers
can not be
written in
Tabular form.

capable of becoming actual.

$\downarrow$
Potential

In $\mathbb{Q}$, we see potential position.

e.g. $\frac{5}{1}$ } potential.
 actual $\neq$

coll a person whose square is +ve is a real number.

Date

• $\sqrt{2}$ cannot be written in $\mathbb{P}$.

any square root which don't have $\sqrt{\quad}$ complete square

(don't stop) • non terminating
(no pattern) • non recurring! } Irrational.

• Open & Close Intervals.

✓ pattern
X terminate

Equal Set:

$[2, 5]$ closed interval, 2 and 5 are included
 $(2, 5)$ open interval, 2 and 5 are not included

$$A = \{1, 2, 3\}$$

$$B = \{2, 3, 1\}$$

$$\{1, 3, 3, 3, 5, 5, 5\}$$

$$= \{1, 3, 5\}$$

Empty set: $\{\}$

Singleton set: $\{1\}$

Size of set: No. of elements in set

$$|A| = 3 \text{ for } A = \{1, 2, 3\}$$

Subset:

set if every element of A is also an element of B.

is subset of B $A \subseteq B$

$$B = \{1, 2, 3\} \quad A \subseteq B$$

$$A = \{1, 2\}$$

$$A \subseteq B \quad A \subset B$$

$$B \subseteq A$$

- $\emptyset$ is a subset of every set.

Date

Proper Subset, Superset.

Can or can not be equal.

$$A = \{1, 2, 3, 4, 5\}$$

$$B = \{2, 3, 5\}$$

$$C = \{1, 2, 3, 4, 5\}$$

$$B \subset A \leftarrow \boxed{B \text{ is a subset of } A \text{ and } A \text{ is a superset of } B.}$$

$$C \subseteq A$$

$$D = \{4, 9, 10\}$$

$$D \not\subseteq A$$

• $\subseteq$ can be proper & improper subset.

• $\subset$ proper subset.

Proper Subset:

cannot be equal.

$$A = \{1, 2, 3, 4, 5\}$$

$$B = \{3, 2, 1\}$$

$$B \subset A$$

Improper Subset

$$A = \{1, 2, 3, 4, 5\}$$

$$B = \{1, 2, 3, 4, 5\}$$

$$A = B$$

Cartesian Product

$$A = \{1, 2, 3\}$$

$$B = \{a, b, c\}$$

$$A \times B = \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c), (3, a), (3, b), (3, c)\}$$

$$2 \times 2 \times 2 \times 2 = 10,$$

Date

$$2^3 = 8$$

Power Set:

Subsets of given set in a single set.

$$2^3 = 8$$

e.g:

Q1 $A = \{1, 2, 3\}$

$$P(A) = \{ \emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\} / A \}$$

→ Number of elements: $2^n = 2^3 = 8$

Q2 $A = \{1, \{2, 3\}\}$

$$2^2 = 4$$

$$P(A) = \{ \emptyset, \{1\}, \{\{2, 3\}\}, \{1, \{2, 3\}\} \}$$

Q3 $A = \{1, 2, \emptyset, \{\emptyset\}\}$

$$2^4 = 16$$

$$P(A) = \{ \emptyset, \{1\}, \{2\}, \{\emptyset\}, \{\{\emptyset\}\},$$

$$= \{ \emptyset, 1\}, \{ \emptyset, 2\}, \{ \emptyset, \emptyset\}, \{ \emptyset, \{\emptyset\}\},$$

$$\{1, \emptyset\}, \{2, \emptyset\}, \{1, 2\}, \{1, \emptyset\}, \{1, \{\emptyset\}\}, \{2, \emptyset\},$$

$$\{2, 1\}, \{2, \emptyset\}, \{2, \{\emptyset\}\}, \{\emptyset, \emptyset\}, \{\emptyset, 1\}, \{\emptyset, 2\}, \{\emptyset, \{\emptyset\}\},$$

$$\{\{\emptyset\}, \emptyset\}, \{\{\emptyset\}, 1\}, \{\{\emptyset\}, 2\}, \{\{\emptyset\}, \{\emptyset\}\}$$

Exercise 4: $\{21, 23, 24, 28, 7, 8, 9\}$
(sets).



$$P(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{\emptyset\}, \{\{1, 2\}\}, \{1, \emptyset\}, \{1, \{1, 2\}\}, \{2, \emptyset\}, \{2, \{1, 2\}\}, \{\emptyset, \{1, 2\}\}, \{1, 2, \emptyset\}, \{1, 2, \{1, 2\}\}, \{2, \emptyset, \{1, 2\}\}, \{1, \emptyset, \{1, 2\}\}, \{1, 2, \emptyset, \{1, 2\}\}\}$$

Q4 $A = \{\{1, 2, 3, 4\}\}$

$$2^1 = 2$$

$$P(A) = \{\emptyset, \{\{1, 2, 3, 4\}\}\}$$

- Q9
a) No
b) No

- $\in$ member
- $\subset$ proper subset
- $\subseteq$ subset.

Exercise:

Q7

- a) Yes
b) No
c) Yes
d) No
e) No
f) No

$$\{2, \{2\}\}$$

a) $0 \in \emptyset$
 $0 \in \{\}$ (No)

b) $\emptyset \in \{0\}$
No, if subset $\subseteq$ then Yes.

c) $\{0\} \subset \emptyset$
No

d) $\emptyset \subseteq \{0\}$

Q8 by your own

f) $\{0\} \subset \{0\}$

Yes

No. g) $\{0\} \subseteq \{0\}$

Yes

Yes

e) $\{0\} \in \{0\}$

$\{0\} \in \{\{0\}\}$ Yes

Yes No. $0 \in \{0\}$ Yes

A set is always a subset of itself

Union



$$\{x | x \in A \vee x \in B\}$$

u v


Date

- (Q8) a) No
b) No
c) Yes
d) Yes

- e) Yes
f) No

Union:

$$A \cup B, A = \{1, 2, 3, 4\}, B = \{3, 4, 6, 7\}$$
$$A \cup B = \{1, 2, 3, 4, 6, 7\}$$




$$A \cup B = \{x | x \in A \vee x \in B\}$$

Intersection / Overlapping: (common)

$$A \cap B = \{x | x \in A \wedge x \in B\}$$

$$A \cap B = \{3, 4\}$$

AND




$A \cap B$ $\wedge$

Disjoint Sets: (No common)

(intersection empty set)

$$A \cap B = \emptyset$$

$$A = \{1, 2, 3\}, B = \{4, 5, 6\}$$



$$A = \{1, 2, 3\}$$
$$B = \{4, 5, 6\}$$

$$|A \cup B| = |A| + |B| - |A \cap B|$$

cardinality

Size /
no. of elements

$$A = \{1, 2, 3, 4\}, B = \{3, 4, 6, 7\}$$

$$A \cup B = |A| + |B| - |A \cap B|$$

$$6 = 4 + 4 - 2$$

$$6 = 6$$

$$A \cup B = \{1, 2, 3, 4, 6, 7\}$$


complete / مکمل / ۳. $\uparrow$

Date


Difference of Sets $(A - B)$ (Complement of B with respect to A)
 $\rightarrow$ elements of A write
 $\rightarrow$ B elements not written which are in A.

$$A - B = \{x \mid x \in A \wedge x \notin B\}$$

$$A = \{1, 2, 3, 4\}, B = \{3, 4, 6, 7\}$$
$$A - B = \{1, 2\}$$



$$A = \{2, 4, 6\}$$
$$A' = U - A = \{1, 3, 5, 7, 8, 9, 10\}$$



$$\bar{A} = \{x \in U \mid x \notin A\}$$

$\downarrow$
A complement

Set Identities:

a ya b ki koi bhi value put
ku dain to $L.H.S = R.H.S$

Date

$$A \cap U = A$$

$$A \cup \phi = A$$

Identity law

$$A \cup U = U$$

$$A \cap \phi = \phi$$

Domination law

$$A \cup A = A$$

$$A \cap A = A$$

Idempotent law

$$\overline{(\overline{A})} = A$$

Complementation law

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Commutative law

$$A \cup (B \cap C) = (A \cup B) \cap C$$

Associative law

$$A \cap (B \cup C) = (A \cap B) \cup C$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

Distributive law

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

De Morgan's law

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$A \cup (A \cap B) = A$$

Absorption law

$$A \cap (A \cup B) = A$$

$$A \cup \overline{A} = U$$

Complement law

$$A \cap \overline{A} = \phi$$

Date

 $\sim A \cap B$

Example 11:

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

$$\overline{A \cap B} = \{x \mid x \notin A \cap B\}$$

$$= \{x \mid \sim(x \in (A \cap B))\}$$

$$= \{x \mid \sim(x \in A \wedge x \in B)\}$$

$$= \{x \mid \sim(x \in A) \vee \sim(x \in B)\}$$



$$= \{x \mid x \notin A \vee x \notin B\}$$

$$= \{x \mid x \in \overline{A} \vee x \in \overline{B}\}$$

$$= \{x \mid x \in \overline{A} \cup \overline{B}\} = \overline{A \cap B}$$

Generalized Unions & Intersections

more than 2

 $A \cup B$  $A \cap B$

Union of Collection of Sets

$$A_1 \cup A_2 \cup \dots \cup A_n = \bigcup_{i=1}^n A_i \rightarrow \text{will change}$$

n - ending index

Intersection of Collection of Sets

$$A_1 \cap A_2 \cap \dots \cap A_n = \bigcap_{i=1}^n A_i$$

1 - starting index

Golden ratio.


Date 2.4 Sequences & Summations.

Sequence:
→ a function.

1, 2, 3, 5, 8, 13, 21, 34, ...

$a_n \rightarrow$ one element

$[a_n] \rightarrow$ all generated elements



Example:

$[a_n]$, $a_n = \frac{1}{n}$, $a_1, a_2, a_3, a_4, \dots$ starts with where.

~~$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$~~

$$a_n = \frac{1}{1} = 1$$

$$a_n = \frac{1}{2} = \frac{1}{2}$$

$$a_n = \frac{1}{3}$$

$$a_n = \frac{1}{4}$$

!

!

2.4 Exercise.

Q1. $[a_n]$ where $a_n = 2 \cdot (-3)^n + 5^n$

$$\begin{aligned} a) \quad a_0 &= 2 \cdot (-3)^0 + 5^0 \\ &= 2 + 1 = 3 \end{aligned}$$

Q2, Q3, Q4

1.5
2.5
2.10

Q1-8, 9, 10, 11

Date

Q5 a) 2, 5, 8, 11, 14, 17, 20, 23, 26, 29

b) 1, 1, 1, 2, 2, 2, 3, 3, 3, 4, 4, 4, 5, 5, 5, 6, 6, 6, ...

c) 1, 3, 3, 5, 5, 7, 7, 9, 9, ...

recursive function

Q9 a) $a_n = 6a_{n-1}$, $a_0 = 2$. (starting point).
Yahan n ki value aisi put krni hai k a_0 aa jaye.

jo function khd ko call krni.

$$\begin{aligned} a_1 &= 6a_{1-1} \\ a_1 &= 6a_0 \\ a_1 &= 6 \cdot 2 \\ a_1 &= 12 \end{aligned}$$

$$\begin{aligned} a_2 &= 6a_1 \\ a_2 &= 6(12) \\ &= 72 \end{aligned}$$

$$\begin{array}{r} 12 \\ \times 6 \\ \hline 72 \\ \times 6 \\ \hline 432 \\ \times 6 \\ \hline 2592 \end{array}$$

$$\begin{aligned} a_3 &= 6a_2 \\ &= 6(72) \\ &= 432 \end{aligned}$$

$$\begin{aligned} a_4 &= 6a_3 \\ &= 6(432) \\ &= 2592 \end{aligned}$$

$$\begin{aligned} a_5 &= 6a_4 \\ &= 6(2592) \end{aligned}$$

c) $a_n = a_{n-1} + 3a_{n-2}$, $a_0 = 1$, $a_1 = 2$

$$\begin{aligned} a_2 &= a_1 + 3a_0 \\ &= 2 + 3(1) \\ &= 5 \end{aligned}$$

Sequence $\rightarrow$ Arithmetic sequence
 $\rightarrow$ Geometric sequence

Arithmetic Sequence:

d = common difference.

$$a_n = a_1 + (n-1)d$$

$$\begin{aligned} &= 5 + (100-1) \times 4 \\ &= 5 + 99 \times 4 = 401 \end{aligned}$$

e.g: 5, 9, 13, 17, ...

Find 100th term.

Find any value of arithmetic progression.

Date

Sum of first n terms of sequence -

$$S_n = \frac{n}{2} \{ 2a_1 + (n-1)d \}$$

$$= \frac{100}{2} \{ 2 \cdot 5 + (100-1)4 \}$$

$$= 50 (10 + 396) = 50(406) = 20300$$

Geometric Sequence

↓
difference matters.

→ have common ratio b/w two consecutive terms.

2, 6, 18, 54,

$$\frac{6}{2} = \frac{18}{6} = \frac{54}{18}$$

$$a_n = a r^{n-1}$$

(formula to find n^{th} term)

To find 20th value: 20-1

$$a_{20} = 2 \cdot 3$$

$$a_{20} = 2 \cdot 3^{19}$$

$$S_n = \frac{a (r^n - 1)}{r - 1}$$

(to find sum)

$$= \frac{2 (3^{20} - 1)}{3 - 1} = \frac{2}{2} \times (3^{20} - 1) = 3^{20} - 1$$

Date

CHESSE, PERSON, KING

5, 5, 5, 5, ...

this is both A.P & G.P.

1 = 1

1 = 1

1 = 1

8x96x85x

40x25

1 + 2 + 4 + 8 + 64 term

$$S_{64} = \frac{1(2^{64} - 1)}{2 - 1} = 2^{64} - 1$$

$$\frac{1}{2} = \frac{1}{2}$$

Summations:

$$S = \frac{a(r^n - 1)}{r - 1}$$

sigma (sum) $\sum_{i=1}^5$ i - expression

$$\sum_{i=1}^5 i = 1 + 2 + 3 + 4 + 5 = 15$$

$$\sum_{i=1}^3 2i = 2 + 4 + 6 = 12$$

$$\sum_{i=1}^x i = 1 + 2 + 3 + \dots + x \quad (\text{This is A.P}) \text{ so,}$$

$$S_n = \left[\frac{n}{2} 2a_1 + (n-1)d \right]$$

$$\sum_{i=1}^{50} 5 = 5 + 5 + 5 \dots 5$$

$$= 5 \times 50 = 250$$

$$\sum_{i=5}^{15} 1 = 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1$$

$$= 11$$

5 sai 15th

1 add ho

tha.

$$15 - 5 + 1 = 11$$

Date

$$\sum_{i=a}^b 2 = \cancel{a} 2(b-a+1)$$

Double Summation

Example: 21

$$\sum_{i=1}^4 \sum_{j=1}^3 ij$$

$$\sum_{i=1}^4 (i + 2i + 3i)$$

$$\sum_{i=1}^4 (6i) = 6(1+2+3+4) = 60$$

$$\sum_{i=0}^x \sum_{j=0}^y \sum_{k=0}^z 1$$

$$= \sum_{i=0}^x \sum_{j=0}^y (z+1)$$

z is constant w.r.t j .

$$= (z+1) + (z+1) + (z+1) \dots$$

$$= \sum_{i=0}^x (y+1)(z+1)$$

$$= (x+1)(y+1)(z+1)$$

Formula for sum of squares

$$n^2 = \frac{n(n+1)(2n+1)}{6}$$

Sum of cubes

$$n^3 = \frac{n^2(n+1)^2}{4}$$

value of k

$$\frac{n(n+1)}{2}$$

$$\sum_{i \in \{3, 7, 11\}} i$$

$$3+7+11 = 21$$

Date: 100/2

Find $\sum_{k=1}^{100} k^2$

$K=50$ if $K=1$ we use $\frac{n(n+1)(2n+1)}{6}$ so,

$$\sum_{k=1}^{100} k^2 = \sum_{k=1}^{49} k^2 + \sum_{k=50}^{100} k^2$$

$$\sum_{k=1}^{100} k^2 - \sum_{k=1}^{49} k^2 = \sum_{k=50}^{100} k^2$$

$$\frac{100(101)(201)}{6} - \frac{49(50)(99)}{6} = \sum_{k=50}^{100} k^2$$

Exercise:

Q25

IMP

Mathematical Induction (Chp # 5)

proof technique.

e.g.:

Sum of n natural no. : $\frac{n(n+1)}{2}$

Proof that wheather it this formula is correct or not.

Mathematical Induction

To 1) check $n=1$

base case.

2) Put k

3) Put $k+1$

Formula
Date

∴ 0 is not a +ve integer.

★ $\frac{n(n+1)}{2}$

• Put $n=1$

$$1 = \frac{1(1+1)}{2} = \frac{2}{2} \neq$$

$$1 = 1 \quad (\text{True})$$

• Put K

$$\frac{K(K+1)}{2} + K+1 = \frac{K^2+K+2K+2}{2} = \frac{K^2+3K+2}{2}$$

Put next term ↑

$$= \frac{(K+1)(K+2)}{2}$$

• Put $K+1$

$$\frac{(K+1)(K+1+1)}{2}$$

$$= \frac{(K+1)^2 (K+1+1)}{2}$$

★ $n^3 - n$ (divisible by 3 formula). (Mathematical Induction) Without

$$n(n^2-1)$$

Induction

$$n(n+1)(n-1)$$

$$-(n-1)(n)(n+1)$$

here any one should be ÷ble by 3.

Use mathematical induction to prove that

4 5 6
6 7 8
8 9 10

$$1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1 \text{ for all non-ve integers } n.$$

$$2^0 + 2^1 + 2^2 + 2^3 + \dots + 2^n = 2^{n+1} - 1 \quad (0 \text{ is included})$$

$$2^0 = 2^{0+1} - 1$$

$$1 = 2 - 1$$

$$1 = 1$$

$$2^0 + 2^1 = 2^{1+1} - 1$$

$$1 + 2 = 2^2 - 1$$

$$3 = 3$$

Date

Put $n = k$.

$$1 + 2 + 2^2 + 2^3 + \dots + 2^n = 2^{n+1} - 1$$

$$\begin{aligned} 1 + 2 + 2^2 + 2^3 + \dots + 2^k + 2^{k+1} &= 2^{k+1} - 1 + 2^{k+1} \\ &= 2^{k+2} - 1 \\ &= 2^{k+1+1} - 1 \quad \text{--- (i)} \end{aligned}$$

next term

$$\begin{aligned} &2^{n+1} - 1 \\ \text{put } k+1 &\text{ in } n \\ &2^{k+1+1} - 1 \quad \text{--- (ii)} \end{aligned}$$

(i) and (ii) proved.

Ex: 02:

Conjecture a formula for the sum of the first n +ve odd integer. Then prove your conjecture using mathematical induction.

$$1 = 1$$

$$1 + 3 = 4$$

$$1 + 3 + 5 = 9$$

$$1 + 3 + 5 + 7 = 16$$

$$1 + 3 + 5 + 7 + 9 = 25$$

$$1, 3, 5, 9, 16, 25, \dots, 2n-1 = n^2$$

Put $n = 1$

$$1 = 1^2$$

$$1 = 1$$

Put $n = k$

$$1 + 3 + 5 + \dots + 2k-1 = k^2$$

$$1 + 3 + 5 + \dots + 2k-1 + 2 = k^2 + 2k + 1$$

$$1 + 3 + 5 + \dots + 2k+1 = (k+1)^2$$

$$n^2 = (k+1)^2$$

because all numbers are increasing by 2.

$$a_n = a + (n-1)d$$

$$2n-1 = 1 + (n-1)2$$

Date

Show that if $2^k < k!$ for an arbitrary +ve integer k where $k \geq 4$, then $2^{k+1} < (k+1)!$

let $2^k < k!$ let,

$$2^4 < 4! = 16 < 24$$

• Put $n = 4$

$$2^4 < 4!$$

$$16 < 24 \text{ (True)}$$

• Put $n = k$

~~$$2^k < k!$$~~

$$2^k < k!$$

$$2^{k+1} = 2 \cdot 2^k$$

$$2^{k+1} < 2 \cdot k!$$

$$2^{k+1} < (k+1) \cdot k!$$

$$2^{k+1} < (k+1)!$$

because we want $k+1$ in answer. so we write this.

we have to prove this -

$$2^{k+1} < (k+1)!$$

Example: 09

Prove that $7^{n+2} + 8^{2n+1}$ is divisible by 57 for every non-ve integer n .

$$= 7^{n+2} + 8^{2n+1}$$

Put $n = 0$

$$= 7^{0+2} + 8^{0+1} = 49 + 8 = 57$$

Date

Put any -ve integer.

let $n = 3$

$$3+2 \quad 2(3)+1$$

$$7^5 + 8^7 \neq 21397$$

~~Put $n = 3$~~

Put $n = k$

Put $n = k$

$$\begin{aligned} & 7^{k+2} + 8^{2k+1} \\ \text{Put } k+1 & 7^{k+1+2} + 8^{2(k+1)+1} \\ & 7^{k+3} + 8^{2k+3} \\ & 7 \cdot 7^{k+2} + 8^2 \cdot 8^{2k+1} \\ & 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ & 7 \cdot 7^{k+2} + (7+57) \cdot 8^{2k+1} \\ & 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ & 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \end{aligned}$$

$7^{n+1+2} + 8^{2(k+1)+1}$ } we have to prove this
we have to prove 57 can divide it.

As this side and this side is $\div$ ble by 57 so prove.

Exercise :

Q17 Prove that $3 + 3 \cdot 5^1 + 3 \cdot 5^2 + \dots + 3 \cdot 5^n =$

$$3(5^{n+1} - 1)/4$$

where n is a non - ve integer.

Base case

Put $n = 0$

$$3 \cdot 5^0 + 3 \cdot 5^1 + 3 \cdot 5^2 + \dots + 3 \cdot 5^n = 3(5^{n+1} - 1)/4$$

$$3 \cdot 5^0 = 3(5^1 - 1)/4$$

$$3 = 3(4)/4$$

$$3 = 3$$

Decimal.

Date :

$$3 \cdot 5^0 + 3 \cdot 5^1 = 3(5^{n+1} - 1)/4$$

$$3 + 15 = 3(5^2 - 1)/4$$

$$18 = \frac{3(24)}{4} = 18$$

$$18 = 18$$

$$n = k$$

$$3 \cdot 5^0 + 3 \cdot 5^1 + 3 \cdot 5^2 + \dots + 3 \cdot 5^n = 3(5^{n+1} - 1)/4$$

$$3 \cdot 5^0 + 3 \cdot 5^1 + 3 \cdot 5^2 + \dots + 3 \cdot 5^k = 3(5^{k+1} - 1)/4$$

$$n = k+1$$

$$3 \cdot 5^0 + 3 \cdot 5^1 + 3 \cdot 5^2 + \dots + 3 \cdot 5^k + 3 \cdot 5^{k+1} = \left[3(5^{k+1} - 1)/4 \right] + 3 \cdot 5^{k+1}$$

$$3(5^{k+1} - 1)/4 + 3 \cdot 5^{k+1} = 3(5^{k+1} - 1)/4 + 3 \cdot 5^{k+1}$$

$$\cancel{3(5^{k+1+1} - 1)/4} = \cancel{3(5^{k+1+1} - 1)/4}$$

$$= 3(5^{k+1} - 1)/4 + 3 \cdot 5^{k+1}$$

$$= \frac{3 \cdot 5^{k+1} - 3 + 12 \cdot 5^{k+1}}{4}$$

$$= \frac{5^{k+1}(3 + 12) - 3}{4}$$

$$= \frac{5^{k+1}(15) - 3}{4}$$

$$\frac{3(5^{n+1} - 1)}{4} = \frac{3(5^{k+1+1} - 1)}{4}$$

Date $k+1$

$$\frac{15 \cdot 5 - 3}{4}$$

$$3 \left(\frac{5^{k+1} - 1}{4} \right)$$

$$\frac{3(5^{k+2} - 1)}{4} = \frac{3(5^{k+1+1} - 1)}{4}$$

Q.6 Prove that

$1 \cdot 1! + 2 \cdot 2! + \dots + n \cdot n! = (n+1)! - 1$ whenever n is a true integer: $n \geq 1$.

$n=1$
 $1 \cdot 1! = (1+1)! - 1$
 $1 = 2! - 1$

$n=2$
 $2 \cdot 2! = 4$

$(2+1)! - 1 = 5$

$3 \cdot 3! = 18$

$4! - 1$

$4 \cdot 3 \cdot 2$

$24 - 1$

$1 = 1$

Put $n=1$

$$1 \cdot 1! = (1+1)! - 1$$

$$1! = 2! - 1$$

$$1 = 1$$

Put $n=k$:

$$1 \cdot 1! + 2 \cdot 2! + 3 \cdot 3! + \dots + k \cdot k! = (k+1)! - 1$$

$$1 \cdot 1! + 2 \cdot 2! + \dots + k \cdot k! + (k+1) \cdot (k+1)! = (k+1)! - 1$$

$$+ (k+1) \cdot (k+1)! - 1$$

$$= (k+1)! [1 + k + 1] - 1$$

$$= (k+1)! \cdot (k+2) - 1$$

$$= (k+2)! - 1$$

$$= (k+1+1)! - 1$$

Date Counting

Combinatorics: science of making combinations.
 Enumeration: counting of objects with certain properties

Rules of Counting:

- Product Rule. $(n_1 \cdot n_2)$
- Addition (Sum) Rule. $(n_1 + n_2)$

Product Rule:

Example 01: (Basic Counting Principle)

1 2 3 4 5 6 7 8 9 10 11 12

$$12 \times 11 = 132$$

↓ for first person
 ↓ after 1 person 11 are left.

$$\begin{array}{r} 12 \\ \times 11 \\ \hline 12 \\ 120 \\ \hline 132 \end{array}$$

Example 02.

A1, A2, A100	}	2600
B1, B2, B100		
Z1, Z2, Z100		

Sum Rule:

Example 01.

$$37 + 83 = 120$$

Date Date.

Exercise:

Q7 How many different 3 letter initial can people have?

$$26^3$$

Q8 How many different three letter initials with none of letters repeated can people have?

$$26 \cdot 25 \cdot 24$$

Q9 How many different 3 letter initials are there that begin with an A?

$$26^2$$

A 26 26.
fix

when anything is fix that power finishes.

Q10 How many bit strings are there of length 8?

$$2^8$$

2 2 2 2 2 2 2 2 choices
(0,1) (0,1) ...

Q11 How many bit strings of length 10 both begin & end with 1?

$$2^8$$

Date

Q13 How many bit strings with length not exceeding n , where n is a true integer, consist entirely of 1's not counting empty strings?

Ans = n .

Q16 How many strings are there of four lowercase letters that have letter 'a' in them?

$$26^4 - 25^4$$

algorithms

Q47 In how many ways can a photographer at a wedding arrange six-people in a row including bride & groom if

a) the bride must be next to groom. factorial (!)

B G _ _ _

~~6!~~

$10 \cdot 4!$

B G _ _ _
_ B G _ _ _
_ _ B G _ _

b) bride must not be next to groom.

$6! - 10 \cdot 4!$

B G _ _ _
_ B G _ _ _
_ _ B G _ _

when G is at first
5

$5 + 5 = 10$

Date.

The Pigeonhole Principle

Ex: 6.1
Q1-40

Inclusion

Exclusion (Subtraction Rule)

Example 18.

How many bit strings of length 8 either start with 1 bit or end with two bits 00?

e.g. 01111100 ✓

1111011 ✓

1001100 X

$$\begin{array}{c} \text{fix} \\ 1 \end{array} \text{ --- } = 2^7$$

$$\text{---} \begin{array}{c} \text{fix} \\ 00 \end{array} = 2^6$$

$$1 \text{ --- } 00 = 2^5$$

$$2^7 + 2^6 - 2^5$$

$$128 + 64 - 32 = 160$$

$$|A_1 \cup A_2| = |A_1| + |A_2| - |A_1 \cap A_2|$$

Permutation

&

Combinations

order matters

order not matters.

If n is a +ve integer with $1 \leq r \leq n$,

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1)$$

Total
↓
Chosen

if $n = 10$

$r = 7$

10.9.8.7.6.5.4

$$\rightarrow 10 - 7 + 1 = 4$$

at most: زیاد سے زیادہ $(x \leq 5)$
 at least: کم سے کم $(x \geq 5)$

Date

c) at least 4 1s (range)

$$= \frac{10!}{4!6!}$$

Pigeon Hole Principle

Decimal: 0-9 (10 digits)

Example: 03

0 - 100

101 scores

102 students.

Generalized Pigeon Hole Principle (Multiplicity version)

53 $\rightarrow$ 3

(A) $\rightarrow$ 1 time

B
C
...

Z

A $\rightarrow$ 2nd time

B
C
...

Z

Z

A/B/C/D... $\rightarrow$ 3rd time.


Date

bones	letters	→ 26
objects	words	→ 53

take ceiling.

$$\left\lceil \frac{\text{Object}}{\text{boxes}} \right\rceil = \left\lceil \frac{N}{K} \right\rceil \quad (\text{Generalized Pigeon Hole principle})$$

Trees



Heirarchical data structure having nodes consisting of relation with other nodes.

leaf node : has no child

Siblings : B, C are siblings

Cousins : F, E are cousins.

Binary tree : Tree whose node has maximum two children.


to us

Traversing Binary Trees :


- Pre order traversal
- In order traversal
- Post order traversal.

Date

Pre-Order Traversal:



(Binary tree)



- Root is traversed (print)
- left subtree in pre-order

Call:


1) A, 2) B, 3) D, 4) H, 5) I, 6) E, 7) C, 8) F, 9) G.

- Right subtree in pre-order.

A, B, D, H, I, E, C, F, G

In order Traversal

- left sub-tree
- Root
- Right sub-tree




H, D, I, B, E, A, F, C, G

Date


Post Order Traversal.

- Left
- Right
- Root



H, I, D, E, B, F, G, C, A

Apply all these



Pre-Order.

1, 2, 4, 8, 10, 9, 5, 3, 6, 11, 7

In
~~Post~~-order

10, 8, 9, 4, 5, 2, 11, 6, 7, 3, 1

Post-Order.

10, 8, 9, 4, 5, 2, 11, 6, 7, 3, 1

Date Relations & their Properties

→ sets (order pair)
→ members
→ members are related or not.

→ link of member of sets.

Cartesian Product

$$A = \{1, 2, 3, 4\}, B = \{a, b, c\}$$

$$A \times B = \{ (1, a), (1, b), (1, c), (2, a), (2, b), (2, c), (3, a), (3, b), (3, c), (4, a), (4, b), (4, c), (4, d) \}$$

(pairing)

$$\text{Subset of } A \times B = \text{Relation } (R) = \boxed{R \subseteq A \times B}$$

Exercise:

Q1 List ordered pairs in the relation R from $A = \{0, 2, 1, 3, 4\}$ to $B = \{0, 1, 2, 3\}$ where $(a, b) \in R$ iff

$$b) a + b = 4$$

$$R = \{ (a, b) \mid a + b = 4 \}$$

$$R = \{ (1, 3), (2, 2), (3, 1), (4, 0) \}$$

Date

divides

d) $a \mid b$

$$R = \{(1,0), (2,0), (2,2), (3,0), (3,3), (4,0)\}$$

e) $\text{gcd}(a, b) = 1$
 ← greatest common divisor

$$R = \{(0,1), (1,0), (1,1), (1,2), (1,3), (2,0), \dots\}$$

Properties of Relation

- Reflexive (connected $\sum_{i=1}^n T_{ii}$)
- Symmetric new $\sum_{i=1}^n T_{ji}$ change $\sum_{i=1}^n T_{ij}$ pair
- Antisymmetric
- Transitive ($a, b \in R$ and $b, c \in R$ then $a, c \in R$)

$B = \{0, 1, 2\}$ **every** element should be reflexive.

$$\rightarrow R = \{(0,0), (1,1), (2,2)\} \text{ (reflexive)}$$

$$R = \{(0,0), (2,2)\} \mid (1,1) \text{ missing so not reflexive.}$$

Symmetric:

Any element can be symmetric.

Examp: R_2, R_3

Transitive:

Example: R_4, R_5, R_6
 by default

Date

Antisymmetric : (not symmetric)

R_4, R_5, R_6

$(a, b) \in R$ &
 $(b, a) \in R$ then
 $a = b$.

1 pair attached (not same)
then change place then
the thing come should
not be there, that is
anti-symmetric

Representing Relations Using matrices

Matrix : 2-D array

(0,1) matrix

Sai relation belong kr sktai hain.

i row
j column

m_{ij} : piece wise function.

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$

Example : 01

after taking transpose and it come same
then it is symmetric

If in centre 1 came it is reflexive.


Date

Representing Relation using graphs/directed graph.

- Graph : combination of vertices of edges

plural of vertex

- loop.



$\vec{AB}$
 $\vec{AC}$

B and C are not directed graph.

Example 8

Directed graph of relation

$$R = \{ (1,1), (1,3), (2,1), (2,3), (2,4), (3,1), (3,2), (4,1) \}$$

