


# EVOLUTION OF LYME BORRELIOSIS EPIDEMIOLOGY IN PRIMARY CARE AND HOSPITAL SETTINGS IN FRANCE DURING THE COVID-19 PANDEMIC (2020-2021)



*Charles Nuttens<sup>1</sup>, Stéphanie Duret<sup>2</sup>, Antoine Bessou<sup>2</sup>, Andreas Pilz<sup>3</sup>, Jennifer Moisi<sup>1</sup>, Elisabeth Baux<sup>4</sup>, Jean-François Faucher<sup>5</sup>, James H. Stark<sup>6</sup>*

1. Pfizer, Paris, France ; 2. IQVIA, Courbevoie, France ; 3. Pfizer, Vienna, Austria ; 4. CHU de Nancy, Nancy, France ; 5. CHU de Limoges, Limoges, France 6. Pfizer, Cambridge, USA

## Introduction

'Stay at home' restrictions during the COVID-19 pandemic temporarily modified outdoor recreational activities habits and access to healthcare services was restrained during this period.

This may have impacted the risk of developing a primary or disseminated form of Lyme borreliosis (LB).

This study analysed [incidence rates of LB by age and region in primary care and hospital settings in 2020 and 2021](#) in France and compared them to previous years.

## Methods

Incidence rates of LB in [primary care](#) were estimated using data from [the national sentinel network \(SN\)](#) and [electronic medical records \(EMR\)](#) of a network of general practitioners independent of the SN.

[Hospitalization rates](#) were calculated from the [national hospital discharge database \(PMSI\)](#). Hospitalized cases were identified using an algorithm combining three LB-specific ICD10 codes (A69.2, M01.2 and L90.4) and compatible codes for disseminated forms.


[Average annual rates](#) were calculated to smooth annual variability.

## Results

### Evolution of incidence rate

Incidence rates in primary care decreased slightly between 2017-2019 and 2020-2021, from 83 cases/100,000 population to 81/100,000 for the SN and 75/100,000 to 63/100,000 for the EMR.


The hospitalization rate significantly decreased, from 1.68 to 1.16 hospitalizations/100,000 population in 2017-2019 and 2020-2021 respectively. The decreasing trend started in 2018.



### Average incidence rate by age group

The age distribution in primary care in 2020-2021 was similar to 2017-2019.


Hospitalization rates remained stable in children 0-4, 5-9 and 10-14 years of age (1.23/100,000 to 1.21/100,000 in 0-14 years) while they decreased in older age groups, from 1.78/100,000 in 2017-2019 to 1.15/100,000 in 2020-2021 in those 15 years of age and older.



### Evolution of Google Trends search volume

Google Trends analysis of [search volume for the keyword "Lyme"](#) during the same time period indicated a trend similar to the evolution of [incidence rates in primary care](#) but preceded by approximately one year.


The incidence rates in primary care reached a maximum in 2018 whereas the search volume for "Lyme" peaked in 2017.



### Evolution of incidence rate by region

Between 2017-2019 and 2020-2021, [incidence rates increased for 11 regions](#) according to the SN and [5 according to the EMR](#), on a total of 22 regions.

[Hospitalization rates decreased in all regions](#) between the two study periods, except in Bretagne (\*).



## Conclusion

Incidence rates in primary care and hospital settings decreased in 2020-2021 compared to 2017-2019.

The decreasing trend started before the COVID-19 era and wasn't amplified in 2020-2021. Also, incidence rates in primary care increased in some regions despite similar restrictions nationwide.

Further analyses will be conducted to explain the evolution of LB epidemiology in 2020-2021.

**Charles Nuttens, PhD**

Medical Development, Scientific and Clinical Affairs - Pfizer  
23-25 avenue du Docteur Lannelongue, 75014 Paris, France  
charles.nuttens@pfizer.com - 0000-0001-6288-8023



Scan the QR code  
to download the PDF  
version of the poster

