Unit 1: Calculus

Charles Rambo

UCLA Anderson

2024

Table of Contents I

1. Limits
2. Continuity
3. Derivatives
   - Definition, Examples, and Formulas
   - Tangent Lines
   - Optimization
   - L'Hôpital's Rule
4. Integration
   - Riemann Integration
   - Indefinite Integration

Table of Contents II

5. Ordinary Differential Equations
   - Separable ODEs
   - First Order Linear ODEs
6. Sequences and Series
   - Sequences
   - Series
   - Power Series
7. Time Value of Money

Limits

Definition

Definition

(a) Define $\lim_{x \to a} f(x) = L$ to mean for all $\epsilon > 0$ there exists a $\delta > 0$ such that

we don't care about f at a

$$0 < |x - a| < \delta \qquad \text{implies} \qquad |f(x) - L| < \epsilon.$$

(b) Define $\lim_{x \to \infty} f(x) = L$ to mean for all $\epsilon > 0$ there exists an N such that

$$x \geq N \qquad \text{implies} \qquad |f(x) - L| < \epsilon.$$

(c) Define $\lim_{x \to -\infty} f(x) = L$ to mean for all $\epsilon > 0$ there exists an N such that

$$x \leq N \qquad \text{implies} \qquad |f(x) - L| < \epsilon.$$

$\lim_{x \to a} f(x) = L$

$$\lim_{x \to 4} \sqrt{x} = 2$$

$\sqrt{3.61} = 1.9$

$\sqrt{x} = 1.9$

$\Rightarrow x = 1.9^2$

Pick $\varepsilon = 0.1$

so, $\delta = 0.39$

$$0 < |x - 4| < 0.39 \Rightarrow |\sqrt{x} - 2| < 0.1$$

A smaller δ would also work, e.g. in this case $\delta = 0.01$.

$\lim_{x \to \infty} f(x) = L$

Using the Definition

Example

Prove

$$\lim_{x\to\infty} \frac{1}{x^p} = 0$$

for all $p > 0$.

Sol The definition $\forall \varepsilon > 0 \ \exists N \in \mathbb{N}$ s.t.

$$x \geq N \implies \left| \frac{1}{x^p} - 0 \right| < \varepsilon.$$

Pick $\varepsilon > 0$. Then Consider

$$\left| \frac{1}{x^p} - 0 \right| = \left| \frac{1}{x^p} \right| = \frac{1}{|x|^p} < \varepsilon$$

$$\iff \frac{1}{\varepsilon} < |x|^p \iff \sqrt[p]{1/\varepsilon} < |x|$$

Let $N = \left\lceil \sqrt[p]{\frac{1}{\varepsilon}} \right\rceil$. ✓

So, we've come up with a rule. You give me $\varepsilon > 0$ and I can give you an N s.t. $x \geq N \Rightarrow \left| \frac{1}{x^p} \right| < \varepsilon$

ϵ-δ Limit Definition on YouTube

Watch BlackPenRedPen explain the ϵ-δ limit definition (https://www.youtube.com/watch?v=DdtEQk_DHQs). There's another video where he goes over the $x \to \infty$ case (https://youtu.be/9JMFLzHtljA?si=1WPW-fmaf2DBe3Ph).

Properties of Limits

Theorem

Suppose a is in the interval $[-\infty, \infty]$. *Let*

$$\lim_{x \to a} f(x) = L_1 \qquad \textit{and} \qquad \lim_{x \to a} g(x) = L_2.$$

(a) $\lim_{x \to a} \alpha f(x) + \beta g(x) = \alpha L_1 + \beta L_2$ *for any real constants* α *and* β

(b) $\lim_{x \to a} f(x) \cdot g(x) = L_1 \cdot L_2$

(c) $\lim_{x \to a} \dfrac{1}{f(x)} = \dfrac{1}{L_1}$ *if* $L_1 \neq 0$.

Useful Limits

Theorem

Suppose $p > 0$.

(a) $\lim_{x\to\infty} \frac{1}{x^p} = 0$

(b) $\lim_{x\to\infty} p^{1/x} = 1$

(c) $\lim_{x\to\infty} x^{1/x} = 1$

(d) *If* $a > 0$, $\lim_{x\to 0} \frac{a^x - 1}{x} = \ln a$

(e) *If* $|r| < 1$, *then* $\lim_{x\to\infty} r^x = 0$

Python Example

Example

Define

$$f(x) = \begin{cases} \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0. \end{cases}$$

Graph f in Python to see that $\lim_{x \to 0} f(x)$ does not exist.

Python Example Cont.

```python
# Import modules
import numpy as np
import matplotlib.pyplot as plt

# Use latex
plt.rcParams['text.usetex'] = True

# Use Seaborn style
plt.style.use('seaborn')

# Define f
f = lambda x: 0 if x == 0 else np.sin(1/x)

# Let's graph on the interval [-pi, pi]
x_vals = np.arange(-np.pi, np.pi, np.pi/200)

# Calculate the y-values
y_vals = [f(x) for x in x_vals]

# Generate the plot
plt.plot(x_vals, y_vals)

# Label the x-axis
plt.xlabel(r'$x$')

# Label the y-axis
plt.ylabel(r'$y$')

# Give the graph a title
plt.title(r'Graph of $y = f(x)$')

# Save the figure
plt.savefig(path + r'Images/ex1-1.png')

# Display the plot
plt.show()
```

By default in radians

start stop

np.arange(_ , _ , _)

step size

def f(x):
→ If x == 0:
→ return 0
else:
→ return np.sin($\frac{1}{x}$)

Python Example Result

The graph isn't perfect, but it's enough to see that f doesn't approach anything in particular as x approaches 0.

$\lim_{x \to 0} f(x) = DNE$ ← does not exist

Continuity

Continuity

Definition

(a) A function f is continuous at a if $\lim_{x \to a} f(x) = f(a)$.

(b) A function f is continuous on the set A if

$$a \in A \qquad \text{implies} \qquad \lim_{x \to a} f(x) = f(a).$$

Continuity Idea

Continuous functions have no breaks, i.e. if you were to draw them you would never need to lift your pencil.

https://machinelearningmastery.com/continuous-functions/

Useful Theorem

Theorem

If f is continuous at b and $\lim_{x \to a} g(x) = b$, then

$$\lim_{x \to a} f\Big(g(x)\Big) = f\left(\lim_{x \to a} g(x)\right) = f(b).$$

$$\lim_{x \to a} f(g(x)) = f\left(\lim_{x \to a} g(x)\right)$$

Example

Example

Suppose $p > 0$. Compute $\lim_{x \to \infty} p^{1/x}$.

Sol First, note

Note: For $p > 0$, p^x is cont. everywhere. So, ...

$$\lim_{x \to \infty} p^{1/x} = p^{\lim_{x \to \infty} 1/x} = p^0 = 1.$$

Derivatives

Derivatives

Definition

(a) The **derivative of a function f** at a number a is

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}.$$

(b) The function f is differentiable on a set A if $f'(a)$ exists for all a in A.

$$\frac{f(a+h) - f(a)}{h}$$

Notation

Leibniz notation is frequently used:

$$\frac{df}{dx} = f'(x) \qquad \text{and} \qquad \left.\frac{df}{dx}\right|_{x=a} = f'(a).$$

The respective second and third derivatives are written

$$\frac{d^2 f}{dx^2} = f''(x) \qquad \text{and} \qquad \frac{d^3 f}{dx^3} = f'''(x)$$

For the k-th derivatives, where $k > 3$, we use the notation

$$\frac{d^k f}{dx^k} = f^{(k)}(x).$$

Derivatives Example

Example

Let

$$f(x) = \begin{cases} xe^{-x^2-x^{-2}}, & x \neq 0 \\ 0, & x = 0. \end{cases}$$

Compute $f'(0)$.

Sol

$$f'(0) = \lim_{h \to 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0} \frac{\cancel{h} e^{-h^2 - h^{-2}} - 0}{\cancel{h}}$$

$$= \lim_{h \to 0} e^{-h^2 - h^{-2}} = e^{-h^2} \cdot e^{-h^{-2}}$$

$$= \lim_{h \to 0} e^{\cancel{-h^2} \to -0^2} \cdot \lim_{h \to 0} e^{\cancel{-h^{-2}} \to -\infty}$$

$$= 1 \cdot e^{\lim_{h \to 0} \cancel{-h^{-2}} \to -\infty}$$

$$= 1 \cdot 0 = \boxed{0}$$

$$\lim_{h\to 0} e^{-h^2} \overset{?}{=} 1$$

$$\lim_{h\to 0} e^{-h^2} = e^{\lim_{h\to 0} -h^2}$$
$$= e^0$$
$$= 1$$

Numerical Approximation

It is often helpful to numerically approximate f'. This can be done by choosing a small value of h and calculating

$$\frac{f(x+h)-f(x)}{h}.$$

The value h can be positive or negative. Often, a better numerical approach is to consider positive and negative values of h at the same time and take the average:

Tends to converge to f' a little faster

$$\frac{1}{2}\cdot\frac{f(x+h)-f(x)}{h}+\frac{1}{2}\cdot\frac{f(x-h)-f(x)}{-h}=\frac{f(x+h)-f(x-h)}{2h}.$$

where $h>0$.

Note

$$\frac{f(x+h) - f(x-h)}{2h}, \quad h > 0$$

tends to go to f' faster when f' exists. But the ratio something odd if f' doesn't exist.

Example $\quad f(x) = |x|$

$f'(0)$ DNE

But

$$\frac{f(0+h) - f(0-h)}{2h}$$

$$= \frac{|h| - |-h|}{2h}$$

$$= \frac{h - h}{2h}$$

$$= 0$$

Since $f'(0)$ doesn't exist, we bad answer

Python Example

Example

Use Python to graph f and f' on the interval $[-2, 2]$, where

$$f(x) = \begin{cases} xe^{-x^2-x^{-2}}, & x \neq 0 \\ 0, & x = 0. \end{cases}$$

Use a numerical approximation for f' with $h = 0.001$.

Python Example Cont.

```python
# Import modules
import numpy as np
import matplotlib.pyplot as plt

# Use latex
plt.rcParams['text.usetex'] = True

# Use Seaborn style
plt.style.use('seaborn')

# Define f
f = lambda x: x * np.exp(-x**2 - x**-2) if
    x != 0 else 0

# Define h
h = 0.001

# Use numerical approximation
f_prime = lambda x: (f(x + h) - f(x - h))
    /(2 * h)

# Get the x-values
x_vals = np.linspace(-2, 2, 100)

# Get the two sets of y-values
y1_vals = [f(x) for x in x_vals]
y2_vals = [f_prime(x) for x in x_vals]
```

means not equal

```python
# Generate the plot for f
plt.plot(x_vals, y1_vals, label = r"$f$")

# Generate the plot for f'
plt.plot(x_vals, y2_vals, label = r"$f'$")

# Label the x-axis
plt.xlabel(r'$x$')

# Label the y-axis
plt.ylabel(r'$y$')

# Give the graph a title
plt.title(r"Graph of $y = f(x)$ and $y = f
    '(x)$")

# Create a legend
plt.legend()

# Save the figure
plt.savefig(path + r'Images/ex1-2.png')

# Display the plot
plt.show()
```

Python Example Result

Derivative Properties

Theorem

Suppose α and β are constants and f' and g' exist.

(a) $\dfrac{d}{dx}(\alpha f + \beta g) = \alpha f' + \beta g'$.

(b) $\dfrac{d}{dx}(f \cdot g) = g \cdot f' + f \cdot g'$ ← Product rule

(c) $\dfrac{d}{dx}\left(\dfrac{f}{g}\right) = \dfrac{f' \cdot g - f \cdot g'}{g^2}$ ← Quotient rule

(d) $\dfrac{d}{dx}(f \circ g) = (f' \circ g) \cdot g'$ ← Chain rule

$\frac{d}{dx}(f(u)) = f'(u) \cdot \frac{du}{dx}$

Useful Derivative Formulas

Suppose $a > 0$.

- $\frac{d}{dx}(x^n) = nx^{n-1}$
- $\frac{d}{dx}(e^x) = e^x$
- $\frac{d}{dx}(a^x) = a^x \ln a$
- $\frac{d}{dx}(\ln|x|) = \frac{1}{x},\ x \neq 0$
- $\frac{d}{dx}(\log_a|x|) = \frac{1}{x \ln a},\ x \neq 0$
- $\frac{d}{dx}(\sin x) = \cos x$
- $\frac{d}{dx}(\cos x) = -\sin x$
- $\frac{d}{dx}(\tan x) = \sec^2 x$
- $\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$

Derivative Example

Example

$$\frac{d}{dx}\left(e^{1/x}\sin x\right) = \frac{d}{dx}\left(e^{x^{-1}}\sin x\right)$$

We will need

- Product rule
- Chain rule
- Power rule

$$\frac{d}{dx}\left(e^{x^{-1}}\right) = e^{+x^{-1}}\cdot\left(-1\cdot x^{-1-1}\right) = e^{-x}\cdot\left(-x^{-2}\right)$$

Sol

$$\frac{d}{dx}\left(e^{1/x}\sin x\right) = \frac{d}{dx}\left(e^{1/x}\right)\cdot\sin x + e^{1/x}\cdot\frac{d}{dx}\left(\overbrace{\sin x}^{\cos x}\right)$$

$$= -\frac{1}{x^2}e^{1/x}\sin x + e^{1/x}\cos x$$

$$\boxed{= \left(\cos x - \frac{\sin x}{x^2}\right)e^{1/x}}$$

Tangent Lines

The tangent line of the graph of $y = f(x)$ at (x_0, y_0) is

$$y = y_0 + f'(x_0)(x - x_0).$$

Tangent Line Example

Example

Approximate $\sqrt{3.9}$.

- Calculate the tangent line at $x = 4$
- Plug in 3.9

This should give us a value close to $\sqrt{3.9}$

Sol $(x_0, y_0) = (4, 2)$; $f(x) = \sqrt{x} = x^{1/2} \Rightarrow f'(x) = \frac{1}{2} x^{1/2 - 1}$

$= \frac{1}{2} x^{-1/2}$

$= \frac{1}{2\sqrt{x}}$

$$\rightarrow f'(4) = \frac{1}{2\sqrt{4}} = \frac{1}{2\cdot 2} = \frac{1}{4}$$

Hence, our tangent line is

$$y = 2 + \frac{1}{4}(x-4)$$
$$= 2 + \frac{1}{4}x - 1$$
$$= 1 + \frac{1}{4}x. \leftarrow$$

good approx of $\sqrt{x}$ for x close to 4

$$\Rightarrow \quad \sqrt{3.9} \approx 1 + \frac{1}{4}\cdot 3.9 = 1 + 0.975$$
$$= 1.975$$

$$\begin{array}{r} .975 \\ 4\overline{)3.900} \\ -36 \\ \hline 30 \\ -28 \\ \hline 20 \end{array}$$

Check: $\sqrt{3.9} \approx 1.974841766\ldots$

Local and Global Extrema

Definition

Let f be a function defined on domain D.

(a) The **global maximum** of f is at c if $f(c) \geq f(x)$ for all x in D. The **global minimum** of f is at c if $f(c) \leq f(x)$ for all x in D. The global maximum and global minimum values of f are called the **global extrema** of f.

(b) A **local maximum** of f is at c if there is an interval (a, b) such that $f(c) \geq f(x)$ for all x in (a, b) and $a < c < b$. Similarly, a **local minimum** of f is at c there is an interval (a, b) such that $f(c) \leq f(x)$ for all x in (a, b) and $a < c < b$. The local maximum and local minimum values of f are called the **local extrema** of f.

Local and Global Extrema Example

y
Local max
Global max
nothing special
Global min
x

Extreme Value Theorem

Theorem (Extreme Value Theorem)

If f is continuous on a closed interval $[a, b]$, then f attains a global maximum value $f(c)$ and a global minimum value $f(d)$ for some numbers c and d in $[a, b]$.

Increasing and Decreasing Functions

For f differentiable.

- If $f'(x) > 0$, then f is increasing at x.
- If $f'(x) < 0$, then f is decreasing at x.

Increasing means: $f(x+h) - f(x) > 0$ for $h > 0$ small.

$$\Rightarrow \frac{f(x+h) - f(x)}{h} > 0$$

$$\Rightarrow f'(x) \geq 0$$

Decreasing means: $f(x+h) - f(x) < 0$ for $h > 0$ small

$$\Rightarrow \frac{f(x+h) - f(x)}{h} < 0$$

$$\Rightarrow f'(x) \leq 0$$

First Derivative Test

Suppose $x = c$ is a critical number, i.e. c is in the domain of f and $f'(c)$ is 0 or undefined.

(a) If f' changes from positive to negative at $x = c$, then f has a local maximum at $x = c$.

(b) If f' changes from negative to positive at $x = c$, then f has a local minimum at $x = c$.

(c) If f' does not change sign at $x = c$, then f has no local extremum at $x = c$.

Local max @ $x = c$ since f' changed from positive to negative at $x = c$.

Same idea for a local min.

Optimization Example

Example

Find the local and global extrema of the function $f(x) = x^3(x-2)^2$. Suppose the domain of f is the closed interval $[-1, 3]$.

Observation: Global extrema occur at endpoints or local extrema
To find local extrema, I'll use the first deriv test.

Sol $f(x) = x^3(x-2)^2 \Rightarrow f'(x) = 3x^2(x-2)^2 + x^3 \cdot 2(x-2)^1 \cdot 1$

$$= 3x^2(x-2)^2 + 2x^3(x-2)$$
$$= x^2\big(3(x-2) + 2x\big)(x-2)$$
$$= x^2(5x-6)(x-2)$$

Critical numbers: $x = 0,\ 6/5,\ 2$

$f(x) = x^3(x-2)^2$

$f(-1) = (-1)^3(-3)^2 = -1 \cdot 9 = -9$

$f(6/5) = (6/5)^3(\overset{-4/5}{6/5 - 2})^2 = \frac{216}{125} \cdot \frac{16}{25} \approx 1.10592$

$f(2) = 2^3 \cdot (2-2)^2 = 0$

$f(3) = 3^3 \cdot (3-2)^2 = 27$

Global min of -9 @ $x = -1$
Global max of 27 @ $x = 3$
Local max of ~ 1.10592 @ $x = 6/5$
Local min of 0 @ $x = 2$

Second Derivative Test

Suppose $f'(c) = 0$.

(a) If $f''(c) > 0$, then f has a local minimum at $x = c$.

(b) If $f''(c) < 0$, then f has a local maximum at $x = c$.

(c) If $f''(c) = 0$ or is undefined, then the test fails.

Concave up
$\Rightarrow f'' > 0$

Concave down
$f'' < 0$

The test fails in some obvious cases.

$f(x) = x^4$

$f'(x) = 4x^3$ ← easy to see crit. at $x = 0$

$\Rightarrow f''(x) = 12x^2$

$\Rightarrow f''(0) = 0$ ← test fails :(

Local Extrema Example

Example

Find all local extrema of $g(x) = x^4 - 4x^3$.

Sol $g(x) = x^4 - 4x^3 \Rightarrow g'(x) = 4x^3 - 12x^2 = 4x^2(x-3)$

The crit. values are when $g'(x) = 0$. So, when

$$4x^3 - 12x^2 = 0$$
$$\Rightarrow 4x^2(x-3) = 0$$
$$\Rightarrow x = 0 \quad \text{or} \quad x = 3$$

Diff. again to find whether these are local mins or maxs:

$g''(x) = 12x^2 - 24x$

$g''(0) = 0$

The test failed, let's use first deriv. test

$g''(3) = 12 \cdot (3)^2 - 24(3) = 108 - 72 = 36$

Since $g''(3) > 0$, local min at $x = 3$.

Nothing special at $x=0$

Let's plug 3 into g, so we can find y-value of local min

$$g(3) = 3^4 - 4(3)^3$$
$$= 81 - 4 \cdot 27$$
$$= 81 - 108$$
$$= -27$$

Local min of -27 @ $x=3$.
No local max.

L'Hôpital's Rule

Theorem (L'Hôpital's Rule)

Suppose f and g are differentiable on the open interval (a, b) and $g'(x) \neq 0$ for all x in (a, b), where $a \leq \cancel{x} < b \leq \infty$. If

$$\lim_{x \to a} \frac{f'(x)}{g'(x)} = L$$

and either

$$\lim_{x \to a} f(x) = \lim_{x \to a} g(x) = 0 \qquad \text{or} \qquad \lim_{x \to a} f(x) = \lim_{x \to a} g(x) = \infty,$$

then

$$\lim_{x \to a} \frac{f(x)}{g(x)} = L.$$

L'Hôpital's Rule Cont.

Example

Prove $\lim_{x\to\infty} x^{1/x} = 1$.

Sol

$\lim_{x\to\infty} x^{1/x} \to \infty^0 \leftarrow$ not a ratio, no so clear

Note: $e^{\ln x} = x$

$$\lim_{x\to\infty} x^{1/x} = \lim_{x\to\infty} (e^{\ln x})^{1/x} = \lim_{x\to\infty} e^{\ln x/x}$$

$$= e^{\lim_{x\to\infty} \frac{\ln x}{x}} \overset{LH}{=} e^{\lim_{x\to\infty} \frac{1/x}{1}} = e^{\lim_{x\to\infty} 1/x} = e^0 = \boxed{1}$$

$\ln x \to \infty$, $x \to \infty$

Integration

Definite Integration

The motivating problem for the definite integral is finding area under the graph $y = f(x)$ for $a \leq x \leq b$.

Riemann Sum

Definition

Suppose we have a function f defined on the interval $[a, b]$. Consider a **partition pair** P and T; $P = (x_0, x_1, \ldots x_n)$ and $T = (t_1, t_2, \ldots, t_n)$, where

$$a = x_0 \leq t_1 \leq x_1 \leq \ldots \leq x_{n-1} \leq t_n \leq x_n = b.$$

The **Riemann sum** corresponding to the partition pair P and T is defined to be

$$\sum_{k=1}^{n} f(t_k)\Delta x_k.$$

Finer and Finer Partition

The rectangles in the right figure do a better job of approximating the area under the curve. This is because the partition in the second figure is "finer", i.e. uses more x-values, than the first.

Partition Mesh

Definition

The **mesh** of partition P is

$$\|P\| = \max\{\Delta x_1, \Delta x_2, \ldots \Delta x_n\}.$$

As $\|P\| \to 0$, the approximation becomes better and better.

Definition

The **Riemann integral** of f over the interval $[a, b]$ is

$$\int_a^b f(x)\ dx = \lim_{\|P\| \to 0} \sum_{k=1}^{n} f(t_k) \Delta x_k$$

whenever the limit converges.

Integral Theorems

Theorem

A bounded function on an interval $[a, b]$ *is Riemann integrable if it is continuous for all but a finite number of points.*

Theorem

If f *and* g *are Riemann-integrable on* $[a, b]$ *and* α *and* β *are constants, then the following hold.*

(a) $\displaystyle\int_a^b \alpha f(x) + \beta g(x)\ dx = \alpha \int_a^b f(x)\ dx + \beta \int_a^b g(x)\ dx$

(b) $\displaystyle\int_a^b f(x)\ dx = \int_a^c f(x)\ dx + \int_c^b f(x)\ dx$ *for any* c *in* $[a, b]$

Analytic Example

Example

Prove $\int_1^a \frac{dx}{x} = \ln a$ for $a > 1$. Use partition pairs of the form

(n+1)-values n-values left-end points

$P = (1, a^{1/n}, a^{2/n} \ldots, a)$ and $T = (1, a^{1/n}, a^{2/n} \ldots, a^{(n-1)/n})$.

Sol

Let's compute the Riemann sum:

height of rectangle

$$\sum_{k=1}^{n} f(t_k)\,\Delta x_k = \sum_{k=1}^{n} \frac{1}{a^{(k-1)/n}} \underbrace{\left(a^{k/n} - a^{(k-1)/n}\right)}_{\text{width}}$$

$t_k = a^{(k-1)/n}$

$\Delta x_k = a^{k/n} - a^{(k-1)/n}$

$f(x) = \frac{1}{x}$

$$= \sum_{k=1}^{n} \frac{a^{k/n}}{a^{(k-1)/n}} - 1$$

$$= \sum_{k=1}^{n} a^{1/n} - 1$$

no k here, so just summing $a^{1/n} - 1$ n-times

$$= n\left(a^{1/n} - 1\right)$$

$$\Rightarrow \int_1^a \frac{dx}{x} = \lim_{n\to\infty} \underbrace{\overset{\infty}{n}\left(\overset{1-1}{a^{1/n} - 1}\right)}_{\infty \cdot 0} = \lim_{n\to\infty} \frac{a^{1/n} - 1}{1/n} = \lim_{x\to\infty} \frac{\overset{0}{a^{1/x} - 1}}{\underset{0}{1/x}}$$

Recall:

$$\frac{d}{du}(a^u) = a^u \cdot u' \ln a$$

$$\overset{LH}{=} \lim_{x \to \infty} \frac{a^{1/x} \cdot -1/x^2 \cdot \ln a - 0}{-1/x^2}$$

$$= \lim_{x \to \infty} \frac{a^{1/x} \cdot \cancel{-1/x^2} \cdot \ln a}{\cancel{-1/x^2}}$$

$$= \lim_{x \to \infty} \underbrace{a^{1/x}}_{1} \cdot \ln a$$

$$= 1 \cdot \ln a$$

$$\boxed{= \ln a}$$

Selection of T

The most common choices for T are

- Left endpoints: $t_k = x_{k-1}$
- Midpoints: $t_k = \frac{x_{k-1}+x_k}{2}$
- Right endpoints: $t_k = x_k$

Left endpoints Midpoints Right endpoints

Selection of P

The most common choice for P, by far, is the uniform partition

$$x_k = a + k\Delta x \qquad \text{and} \qquad \Delta x = \frac{b-a}{n}.$$

Useful formulas

When working analytic problems with a uniform partition, these formulas come up a lot

$$\sum_{k=1}^{n} k = \frac{n(n+1)}{2}, \qquad \sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6},$$

and

$$\sum_{k=1}^{n} k^3 = \frac{n^2(n+1)^2}{4} = \left[\frac{n(n+1)}{2}\right]^2.$$

Uniform Partition Example

Example

Use uniform partitions and right endpoints to find $\int_0^1 x^2\, dx$.

Supposed to be uniform horizontal length!

Steps:
1. Find $\Delta x_k = \Delta x$
2. Find $x_k = a + k\Delta x$ ← right endpoints
3. Find $f(x_k)\Delta x$
4. Find $\sum_{k=1}^{n} f(x_k)\Delta x$
5. Find $\lim_{n\to\infty} \sum_{k=1}^{n} f(x_k)\Delta x$

Sol

1. $\Delta x = \frac{1-0}{n} = 1/n$

2. $x_k = 0 + k \cdot \Delta x$ ← right end points

$= k/n$

3. $f(x_k)\Delta x = (k/n)^2 \cdot 1/n = k^2/n^3$

4. $\sum_{k=1}^{n} k^2/n^3 = \frac{1}{n^3} \sum_{k=1}^{n} k^2 = \frac{1}{n^3} \cdot \frac{n(n+1)(2n+1)}{6}$

5. $\lim_{n\to\infty} \frac{1}{n^{\cancel{3}\,2}} \cdot \frac{\cancel{n}(n+1)(2n+1)}{6} = \lim_{n\to\infty} \frac{(n+1)(2n+1)}{6n^2}$ $\cdot \frac{1}{n^2}$ $\cdot 1/n$ $\cdot 1/n$ 6

$= \lim_{n\to\infty} \frac{(1+1/n)(2+1/n)}{6}$

$= \frac{(1+0)(2+0)}{6}$

$= \boxed{1/3}$

Python Code

```python
# Define Riemann sum
def riemann_sum(f, P, pts):
    # Sort values
    P = np.sort(P)
    # Calculate Delta x
    dx_vals = np.diff(P)
    # Calculate the number of terms
    # Note: dx_vals has one fewer element than P
    N = len(dx_vals)
    # Define T
    if pts == 'left':
        T = [P[i] for i in range(0, N)]
    elif pts == 'right':
        T = [P[i + 1] for i in range(0, N)]
    elif pts == 'mid':
        T = [0.5 * P[i] + 0.5 * P[i + 1] for i in range(0, N)]
    # Get area of rectangles
    rectangles = [f(T[i]) * dx_vals[i] for i in range(0, N)]
    # Return sum
    return np.sum(rectangles)
```

makes sure values are from smallest to biggest

takes diff. of consecutive terms

0, 1, 2, ..., N-1 ← last value is not included

range(N)

$P = (x_0, x_1, x_2, \ldots, x_{N-1}, x_N)$

$T = \left(\frac{x_0+x_1}{2}, \frac{x_1+x_2}{2}, \ldots, \frac{x_{N-1}+x_N}{2}\right)$

height

width

Take sum of rectangle areas to get Riemann sum

Improper Integral

Definition

(a) If the integrals exists for every $t \geq a$ and for every $s \leq b$, then

$$\int_a^{\infty} f(x)\ dx = \lim_{t \to \infty} \int_a^t f(x)\ dx$$

and

$$\int_{-\infty}^{b} f(x)\ dx = \lim_{s \to -\infty} \int_s^b f(x)\ dx.$$

(b) For any c in $\mathbb{R}$, if both $\int_c^{\infty} f(x)\ dx$ and $\int_{-\infty}^{c} f(x)\ dx$ converge, then

$$\int_{-\infty}^{\infty} f(x)\ dx = \int_{-\infty}^{c} f(x)\ dx + \int_c^{\infty} f(x)\ dx.$$

Python Example

Example

Use `pandas` to create a data frame of Riemann sums with left endpoints, midpoints, and right endpoints, and uniform partitions to approximate $\int_0^\infty e^{-x^2/2}\,dx$. Consider $n = 10, 50, 100, 500$, and 1000.

Python Example Cont.

Solution. Since $e^{-x^2/2}$ goes to 0 rapidly, it's safe to use $b = 10$.

```
# Import pandas
import pandas as pd
```

Assume we've already imported numpy as np

$e^{-10^2/2} = e^{-50} \approx 1.93 \times 10^{-22}$

```
# Define function
f = lambda x: np.e**(-x**2/2)

# Define the n-values
n_vals = [10, 50, 100, 500, 1000]

# Define the data frame
results = pd.DataFrame(index = n_vals, columns = ['left', 'mid', 'right'])

# Loop over values
for n in n_vals:
    # We can use np.linspace for a uniform partition
    partition = np.linspace(0, 10, n + 1)
```

note includes endpoints

```
    # Get left endpoint results
    results.loc[n, 'left'] = riemann_sum(f, partition, 'left')
    # Get midpoint results
    results.loc[n, 'mid'] = riemann_sum(f, partition, 'mid')
    # Get right endpoint results
    results.loc[n, 'right'] = riemann_sum(f, partition, 'right')

# Note: screenshot of output is ex1-3
results
```

Python Example Result

The limit as $\|P\| \to 0$ is $\sqrt{\pi/2} \approx 1.253$.

n	left	mid	right
10	1.753314	1.253314	0.753314
50	1.353314	1.253314	1.153314
100	1.303314	1.253314	1.203314
500	1.263314	1.253314	1.243314
1000	1.258314	1.253314	1.248314

Little big

Little small

Indefinite Integral

Definition

The function F is an **indefinite integral** or **antiderivative** of f if $F'(x) = f(x)$. We write

$$\int f(x)\, dx = F(x)$$

to denote this.

Indefinite integrals are only unique up to a constant. For example, two antiderivatives of $2x$ are $x^2 + 1$ and $x^2 - 4$. To handle all possibilities, we write

$2x + 0 = 2x - 0$

$$\int 2x\, dx = x^2 + C.$$

Antiderivative Theorems

Theorem

Let f and g be continuous functions on some domain and let α and β be real numbers. Then

$$\int \alpha f(x) + \beta g(x)\, dx = \alpha \int f(x)\, dx + \beta \int g(x)\, dx.$$

Useful Antiderivative Formulas

- $\int x^n \, dx = \frac{x^{n+1}}{n+1} + C, \; n \neq -1$
- $\int \frac{dx}{x} = \ln|x| + C$
- $\int \frac{dx}{1+x^2} = \arctan x + C$
- $\int e^x \, dx = e^x + C$
- $\int a^x \, dx = \frac{a^x}{\ln a} + C$
- $\int \sin x \, dx = -\cos x + C$
- $\int cosx \, dx = \sin x + C$
- $\int \tan x \, dx = -\ln|cosx| + C$

Fundamental Theorem of Calculus

Theorem (Fundamental Theorem of Calculus)

Suppose f is continuous on the closed interval $[a, b]$. Then

(a) $\int_a^b f(x)\ dx = F(b) - F(a)$, *where* $F'(x) = f(x)$.

(b) $\frac{d}{dx}\left(\int_a^x f(t)\ tx\right) = f(x)$

Just idea of why true. Not proof

(a) $\int_a^b f(x)\,dx = F(b) - F(a)$, where $F'(x) = f(x)$

$$\frac{F(x_k) - F(x_{k-1})}{\Delta x_k} \approx f(x_k), \qquad P = (\overset{a}{\overset{\shortparallel}{x_0}}, x_1, x_2, \ldots, \overset{b}{\overset{\shortparallel}{x_n}})$$

$$\int_a^b f(x)\,dx = \lim_{n\to\infty} \sum_{k=1}^{n} f(x_k)\Delta x_k$$

$$\approx \lim_{n\to\infty} \sum_{k=1}^{n} \frac{F(x_k) - F(x_{k-1})}{\cancel{\Delta x_k}} \cdot \cancel{\Delta x_k}$$

$$= \lim_{n\to\infty} \sum_{k=1}^{n} \left(F(x_k) - F(x_{k-1})\right)$$

$$= \lim_{n\to\infty} \left(F(\cancel{x_1}) - F(a)\right) + \left(F(x_2) - F(\cancel{x_1})\right) + \ldots + \left(F(b) - F(\cancel{x_{n-1}})\right)$$

$$= \lim_{n\to\infty} F(b) - F(a)$$

$$= F(b) - F(a)$$

(b) $\frac{d}{dx}\left(\int_a^x f(t)\,dt\right) = f(x)$

$$\lim_{h\to 0} \frac{\int_a^{x+h} f(t)\,dt - \int_a^x f(t)\,dt}{h} = \lim_{h\to 0} \frac{\int_x^{x+h} f(t)\,dt}{h}$$

$$= \lim_{h\to 0} \frac{f(x)\cdot \cancel{h}}{\cancel{h}}$$

$$= f(x)$$

Fundamental Theorem of Calculus Example

Example

$$\int_0^2 \max\{x, 1\}\, dx =$$

Sol Notice:

$$\max\{x, 1\} = \begin{cases} 1, & x \le 1 \\ x, & x > 1. \end{cases}$$

So,

$$\int_0^2 \max\{x,1\}\,dx = \int_0^1 \max\{x,1\}\,dx + \int_1^2 \max\{x,1\}\,dx$$

$$= \int_0^1 1\,dx + \int_1^2 x\,dx$$

$$= [x]_0^1 + \left[\tfrac{1}{2}x^2\right]_1^2$$

$$= (1-0) + \left(\tfrac{1}{2}\cdot 2^2 - \tfrac{1}{2}\cdot 1^2\right)$$

Note: $\frac{d}{dx}(x+c) = 1$

$\frac{d}{dx}\left(\frac{1}{2}x^2 + c\right) = x$

$$= 1 + \frac{1}{2} \cdot 4 - \frac{1}{2}$$

$$= 1 + 2 - \frac{1}{2}$$

$$= \frac{6}{2} - \frac{1}{2}$$

$$= 5/2$$

Proof of the Fundamental Theorem on YouTube

Watch Peyam prove part (b) of the Fundamental Theorem of Calculus (https://youtu.be/4DrCKhCECHo).

u-Substitution

Theorem (u-Substitution)

Suppose g' is continuous on the closed interval $[a, b]$ and f is continuous on the range of g. Then

$$\int_a^b (f \circ g)(x) \cdot g'(x)\ dx = \int_{g(a)}^{g(b)} f(u)\ du.$$

$u = g(x)$

Example

Example

$$\int xe^{-x^2/2}\,dx =$$

Sol

$$\int x e^{-x^2/2}\,dx = -\int e^{\overbrace{-x^2/2}^{u}} \cdot \overbrace{(-x\,dx)}^{du}$$

Let $u = -x^2/2$.

$\Rightarrow du = -x\,dx$

$$= -\int e^u\,du$$

$$= -e^u + C$$

$$\boxed{= -e^{-x^2/2} + C}$$

Integration by Parts

Theorem (Integration by parts)

Suppose F and G are differentiable functions, $F'(x) = f(x)$, and $G'(x) = g(x)$, where f and g are continuous. Then

$$\int F(x)g(x)\,dx = F(x)G(x) - \int f(x)G(x)\,dx.$$

$$\int u\,dv = uv - \int v\,du$$

Strategy: Choose u so that it becomes less complex when diff.
E.g.
Polynomials, $\ln x$, inverse trig functions
Choose dv so that it's at least not more complex when you integrate
E.g. e^x, trig functions

Example

Example

$$\int \ln x \, dx =$$

Clear we want $u = \ln x$ because $du = \frac{1}{x}\,dx$ ← way easier to deal with than $\ln x$

we can let $dv = dx$, because $v = x$ ← x isn't much harder to deal with than 1

Sol

$u = \ln x \qquad du = \frac{1}{x}\,dx$

$dv = dx \qquad v = x$

$$\int u\,dv = uv - \int v\,du$$

$$\Rightarrow \int \ln x\,dx = \underbrace{(\ln x)}_{u} \cdot \underbrace{x}_{v} - \int \underbrace{x}_{v} \underbrace{\left(\frac{1}{x}\,dx\right)}_{du}$$

$$= x\ln x - \int 1\,dx$$

$$\boxed{= x\ln x - x + C}$$

Ordinary Differential Equations

Ordinary Differential Equations

Definition

(a) An **ordinary differential equation** (ODE) involves an unknown function of a single variable and some of its derivatives.

(b) The **order** of a differential equation is the order of the highest derivative that appears in the equation.

For example, $xy' = e^{xy}$ is a first order ordinary differential equation, while

$$\frac{d^3x}{dt^3} - 2t\frac{d^2x}{dt^2} + t^2x = \cos t$$

is a third order ordinary differential equation.

Separable ODEs

Definition

An ODE of the form

$$\frac{dy}{dx} = F(x, y)$$

is separable if $F(x, y) = f(x)g(y)$.

It's relatively easy to solve separable differential equations

$$\frac{dy}{dx} = f(x)g(y) \quad \text{implies} \quad \frac{1}{g(y)}\frac{dy}{dx} = f(x).$$

Hence,

$$\int \frac{1}{g(y)}\frac{dy}{dx}\, dx = \int f(x)\, dx \qquad \text{implies} \qquad \int \frac{1}{g(y)}\, dy = \int f(x)\, dx.$$

Separable ODE Example

Example

Solve the differential equation

$$\frac{dy}{dt} = \frac{ty + 3t}{t^2 + 1}$$

subject to the initial condition $y(0) = 2$.

Sol

$$\frac{dy}{dt} = \frac{ty + 3t}{t^2+1} = \frac{(y+3)t}{t^2+1} = (y+3) \cdot \frac{t}{t^2+1}$$

$$\Rightarrow \frac{1}{y+3}\frac{dy}{dt} = \frac{t}{t^2+1} \Rightarrow \int \frac{1}{y+3}\overbrace{\frac{dy}{dt}\,dt}^{dy} = \int \frac{t}{t^2+1}\,dt$$

$$\Rightarrow \int \frac{dy}{y+3} = \frac{1}{2}\int \overbrace{\frac{2t}{t^2+1}\,dt}^{du}$$

$u = t^2 + 1$

$du = 2t\,dt$

$$\Rightarrow \ln|y+3| = \frac{1}{2}\int \frac{du}{u} = \frac{1}{2}\ln|u| + C$$

$$\Rightarrow \ln|y+3| = \frac{1}{2}\ln|t^2+1| + C$$

$$\Rightarrow \cancel{e}^{\cancel{\ln}|y+3|} = e^{\frac{1}{2}\ln|t^2+1| + C}$$

$$\Rightarrow |y+3| = e^{C} \cdot \left(\cancel{e}^{\cancel{\ln}|t^2+1|}\right)^{1/2}$$

$$= e^{C}\,|t^2+1|^{1/2}$$

$$\Rightarrow y+3 = \underbrace{\pm e^{C}}_{A}\sqrt{t^2+1}$$

$$\Rightarrow y = A\sqrt{t^2+1} - 3$$

We have initial condition: $y(0) = 2$.

$$\Rightarrow 2 = A\sqrt{0^2+1} - 3$$

$$\Rightarrow 2 = A - 3$$

$$\Rightarrow A = 5$$

Solution

$$\boxed{y = 5\sqrt{t^2+1} - 3}$$

Linear ODEs

Definition

A first order **linear** differential equation is one that can be written in the form

$$\frac{dy}{dx} + P(x)y = Q(x).$$

The idea behind solving these is to find $\mu = f(x)$ such that

$$\frac{d}{dx}(\mu y) = \mu\frac{dy}{dx} + \overbrace{\mu P(x)}^{d\mu/dx}y.$$

Using the product rule, it becomes clear

$$\frac{d\mu}{dx} = \mu P(x) \qquad \text{implies} \qquad \boxed{\mu = \exp\left(\int P(x)\ dx\right).}$$

Linear ODEs Example

Example

Solve the differential equation

$$x\frac{dy}{dx} + 3x^3 y = 6x^3.$$

Sol Put into standard form:

$$x\frac{dy}{dx} + 3x^3 y = 6x^3 \Rightarrow \frac{dy}{dx} + 3x^2 y = 6x^2$$

$$\Rightarrow \underbrace{\mu\frac{dy}{dx} + 3x^2\mu y}_{\frac{d}{dx}(\mu y)} = 6x^2\mu$$

$$\frac{d\mu}{dx} = 3x^2\mu$$

$$\Rightarrow \int \frac{1}{\mu}\frac{d\mu}{dx}dx = \int 3x^2\,dx$$

$$\Rightarrow \ln|\mu| = x^3 + \cancel{C}^{\,0}$$

$$\Rightarrow \mu = e^{x^3}$$

$$\Rightarrow e^{x^3}\frac{dy}{dx} + 3x^2 e^{x^3} y = 6x^2 e^{x^3}$$

$$\Rightarrow \frac{d}{dx}\left(e^{x^3} y\right) = 6x^2 e^{x^3}$$

$$\Rightarrow \quad e^{x^3} y = \int 6x^2 e^{x^3}\, dx$$

Let $u = x^3$.

$\Rightarrow du = 3x^2 dx$

$$= 2\int e^{\overbrace{x^3}^{u}} \cdot \overbrace{(3x^2\, dx)}^{du}$$

$$= 2\int e^u\, du$$

$$= 2e^u + C$$

$$= 2e^{x^3} + C$$

$$\Rightarrow \quad \boxed{y = 2 + ce^{-x^3}}$$

Sequences and Series

Sequences

Definition

A sequence $(a_n)_{n=1}^{\infty}$ is said to **converge**, if there is a value a in $\mathbb{R}$ which has the property that: For all $\epsilon > 0$, there exists an integer N such that $n \geq N$ implies that $|a_n - a| < \epsilon$. We often write

$$a_n \to a \qquad \text{or} \qquad \lim_{n\to\infty} a_n = a$$

when $(a_n)_{n=1}^{\infty}$ converges to a. If $(a_n)_{n=1}^{\infty}$ does not converge, then it **diverges**.

Pick $\varepsilon > 0$. If the series converges to a, there exists N s.t. $n \geq N$ implies a_n and a are less than ε distance apart, i.e. $|a_n - a| < \varepsilon$.

Sequences Example

Example

Determine which of the following sequences converge/diverge. If the sequence converges, find its limit.

(a) $a_n = \frac{1}{n}$ Converge. $\lim_{n\to\infty} a_n = \lim_{n\to\infty} \frac{1}{n} = 0$

(b) $b_n = \sqrt{n}$ Diverges

(c) $c_n = (-1)^n$ $-1, 1, -1, 1, \ldots$ Since this doesn't approaching anything it diverges.

(d) $d_n = 1 + \frac{(-1)^n}{n}$ $\lim_{n\to\infty} 1 + \frac{(-1)^n}{n} = 1 + 0 = 1$ Converges

0

Python Sequences Example

Example

Use Python to graph the four sequences in the previous example. Graph them on separate subplots, and for the sequences that converge use horizontal lines to show their respective limits.

Python Sequences Example Solution

```python
# Import modules
import numpy as np
import matplotlib.pyplot as plt

# Use LaTeX
plt.rcParams['text.usetex'] = True

# Use Seaborn style
plt.style.use('seaborn')

# Define functions
a = lambda n: 1/n
b = lambda n: np.sqrt(n)
c = lambda n: (-1)**n
d = lambda n: 1 + (-1)**n/n

# Define the limits
a_lim, d_lim = 0, 1

# Get the n-values
n_vals = np.arange(1, 21)

# Get the sequence values
# Functions already vectorized
a_vals = a(n_vals)
b_vals = b(n_vals)
c_vals = c(n_vals)
d_vals = d(n_vals)

# Set up subplots
fig, ax = plt.subplots(2, 2, sharey = True,
    figsize = (10, 6))

# Plot a_n and its limit
ax[0, 0].scatter(n_vals, a_vals)
ax[0, 0].axhline(y = a_lim, color = 'r',
    linestyle = 'dashed')
ax[0, 0].set_xlabel(r'$n$')
ax[0, 0].set_ylabel(r'$a_n$')

# Plot b_n and its limit
ax[0, 1].scatter(n_vals, b_vals, label = r'
    $b_n$')
ax[0, 1].set_xlabel(r'$n$')
ax[0, 1].set_ylabel(r'$b_n$')

# Plot c_n and its limit
ax[1, 0].scatter(n_vals, c_vals)
ax[1, 0].set_xlabel(r'$n$')
ax[1, 0].set_ylabel(r'$c_n$')

# Plot d_n and its limit
ax[1, 1].scatter(n_vals, d_vals)
ax[1, 1].axhline(y = d_lim, color = 'r',
    linestyle = 'dashed')
ax[1, 1].set_xlabel(r'$n$')
ax[1, 1].set_ylabel(r'$d_n$')

plt.suptitle(r'Sequence Plots')

# Save the figure
plt.savefig(path + r'ex1-4.png')

plt.show()
```

Plot for an

define sequences

start

stop

The default step value is 1

Adds title on top of everything

red

Python Sequences Example Result

Triangle Inequality

For any real numbers x, y, and z,

$$|x - y| \leq |x - z| + |z - y|.$$

Sequences

Theorem

(a) *The sequence* $(a_n)_{n=1}^{\infty}$ *converges to* a *in* $\mathbb{R}$ *if and only if for every* $\epsilon > 0$, *we have* a_n *in the interval* $(a - \epsilon, a + \epsilon)$ *for all but finitely many* n. More-or-less definition of convergence

(b) *If* $(a_n)_{n=1}^{\infty}$ *converges to both* a *and* b, *then* $a = b$.

(c) *If* $(a_n)_{n=1}^{\infty}$ *converges, then it is bounded. That is, convergence of* $(a_n)_{n=1}^{\infty}$ *implies there exists a real number* B *such that* $|a_n| \leq B$ *for all* n.

All of terms eventually a finite distance from 0.

(b) $a_n \to a$ and $a_n \to b \overset{?}{\Longrightarrow} a = b$

Pick $\epsilon > 0$. Then $\exists\, N_1$ and N_2 such that

$$|a_n - a| < \varepsilon \quad \text{for} \quad n \geq N_1$$

and

$$|a_n - b| < \varepsilon \quad \text{for} \quad n \geq N_2.$$

Let $N = \max\{N_1, N_2\}$. Then $n \geq N$, implies

$$|a_n - a| < \varepsilon \quad \text{and} \quad |a_n - b| < \varepsilon.$$

From the triangle inequality,

$$\begin{aligned} |a - b| &\leq |a_n - a| + |a_n - b| \\ &< \varepsilon + \varepsilon \\ &= 2\varepsilon. \end{aligned}$$

This must hold for all ε. Since a and b are not functions of n, this means $a = b$.

Sequence Properties

Theorem

Suppose $(a_n)_{n=1}^{\infty}$ *and* $(b_n)_{n=1}^{\infty}$ *are real numbered sequences and*

$$\lim_{n\to\infty} a_n = a \qquad \text{and} \qquad \lim_{n\to\infty} b_n = b.$$

Let α *and* β *be real constants.*

(a) $\lim_{n\to\infty} (\alpha a_n + \beta b_n) = \alpha a + \beta b$

(b) $\lim_{n\to\infty} a_n b_n = ab$

(c) $\lim_{n\to\infty} \frac{1}{a_n} = \frac{1}{a}$ *if* $a_n \neq 0$ *and* $a \neq 0$.

Monotonic Sequences

Definition

- A real sequence $(a_n)_{n=1}^{\infty}$ is **monotonically increasing** if $a_n \leq a_{n+1}$ for all n. monotonically increasing
- A real sequence $(a_n)_{n=1}^{\infty}$ is **monotonically decreasing** if $a_n \geq a_{n+1}$ for all n. monotonically decreasing

Theorem

Suppose that $(a_n)_{n=1}^{\infty}$ is monotonic. Then it converges if and only if it is bounded. Really hard proof!

Series

Definition

Consider a series $S = \sum_{k=1}^{\infty} a_k$. Its ***n*-th partial sum** is $S_n = \sum_{k=1}^{n} a_k$. The series S **converges** if the sequence $(S_n)_{n=1}^{\infty}$ converges, and it **diverges** otherwise.

$S = \sum_{k=1}^{\infty} a_k \longrightarrow$

$S_1 = a_1$

$S_2 = a_1 + a_2$

$S_3 = a_1 + a_2 + a_3$

$\vdots$

$S_n = a_1 + a_2 + \dots + a_n$

$\longrightarrow (S_n)_{n=1}^{\infty}$

The series converges if the induced sequence converges.

Example

Example

For what values of r does the geometric series $\sum_{k=1}^{\infty} r^{k-1}$ converge?

Sol Consider the partial sums.

$$S_1 = 1$$
$$S_2 = 1 + r$$
$$S_3 = 1 + r + r^2$$
$$\vdots$$
$$S_n = 1 + r + r^2 + \dots + r^{n-1}$$

So,

$$\begin{aligned} S_n &= 1 + \cancel{r} + \cancel{r^2} + \dots + \cancel{r^{n-1}} \\ -(rS_n &= \cancel{r} + \cancel{r^2} + \dots + \cancel{r^{n-1}} + r^n) \\ \hline S_n - rS_n &= 1 - r^n \end{aligned}$$

$$\Rightarrow (1-r)S_n = 1 - r^n$$

$$\Rightarrow \quad S_n = \frac{1-r^n}{1-r}$$

Hence, the series converges if and only if

$$\left(\underbrace{\frac{1-r^n}{1-r}}_{S_n}\right)_{n=1}^{\infty}$$

converges. So, consider

$$\lim_{n\to\infty} \frac{1-\cancel{r^n}^{\,0}}{1-r}.$$

When does this converge?

Converge if $|r|<1$

The series $S = \sum_{k=1}^{\infty} r^{k-1}$ converges if and only if $|r|<1$.

If $|r|<1$,

$$\boxed{S = \sum_{k=1}^{\infty} r^{k-1} = \lim_{n\to\infty} \frac{1-r^n}{1-r} = \frac{1}{1-r}.}$$

Geometric Series

The geometric series is extremely important in finance. Remember these formulas.

$$\sum_{k=1}^{n} ar^{k-1} = \frac{a(1-r^n)}{1-r}$$

and

$$\sum_{k=1}^{\infty} ar^{k-1} = \begin{cases} \frac{a}{1-r}, & |r| < 1 \\ DNE, & |r| \geq 1 \end{cases}$$

Make sure you know these well!!

Partial Fraction Decomposition

- $\dfrac{ax+b}{(x-c)(x-d)} = \dfrac{A}{x-c} + \dfrac{B}{x-d}, \quad c \neq d.$
- $\dfrac{ax+b}{(x-c)^2} = \dfrac{A}{x-c} + \dfrac{B}{(x-c)^2}.$

Partial Fraction Decomposition Example

Example

Rewrite $\dfrac{x-2}{x(x-3)}$ using partial fraction decomposition.

Sol

$$\frac{x-2}{x(x-3)} = \frac{A}{x} + \frac{B}{x-3}$$

Holds for all x values

$$\Rightarrow \quad x-2 = A(x-3) + Bx$$

Let $x=0$. $\Rightarrow -2 = -3A + 0$
$\Rightarrow A = 2/3$

Let $x=3$. $\Rightarrow 1 = 0 + 3B$
$\Rightarrow B = 1/3$

So,

$$\boxed{\frac{x-2}{x(x-3)} = \frac{2/3}{x} + \frac{1/3}{x-3}}$$

Convergent Series

Example

Show $\displaystyle\sum_{k=1}^{\infty} \frac{1}{k(k+1)}$ converges.

Sol We want to consider the partial sums.

$$S_n = \sum_{k=1}^{n} \frac{1}{k(k+1)}$$

We want to use partial fraction decomposition to break down $\frac{1}{k(k+1)}$

$$\frac{1}{k(k+1)} = \frac{A}{k} + \frac{B}{k+1} \Rightarrow 1 = A(k+1) + Bk$$

let $k=0$: $1 = A(0+1) + 0 \Rightarrow A = 1$

let $k=-1$: $1 = \overbrace{A(-1+1)}^{0} + B \cdot -1$

$\Rightarrow B = -1$

$$\Rightarrow \frac{1}{k(k+1)} = \frac{1}{k} - \frac{1}{k+1}$$

$$S_n = \sum_{k=1}^{n} \frac{1}{k(k+1)} = \sum_{k=1}^{n} \frac{1}{k} - \frac{1}{k+1}$$

$$S_n = \sum_{k=1}^{n}\left(\frac{1}{k} - \frac{1}{k+1}\right) = \underbrace{\left(\frac{1}{1} - \cancel{\frac{1}{2}}\right)}_{k=1} + \underbrace{\left(\cancel{\frac{1}{2}} - \cancel{\frac{1}{3}}\right)}_{k=2} + \cancel{\cdots} + \underbrace{\left(\cancel{\frac{1}{n-1}} - \cancel{\frac{1}{n}}\right)}_{k=n-1} + \underbrace{\left(\cancel{\frac{1}{n}} - \frac{1}{n+1}\right)}_{k=n}$$

$$= 1 - \frac{1}{n+1}$$

To prove $S = \sum_{k=1}^{\infty} \frac{1}{k(k+1)}$ Converges, all we need to do is

show the sequence $S_n = 1 - \frac{1}{n+1}$ Converges.

$$\lim_{n\to\infty} 1 - \cancelto{0}{\frac{1}{n+1}} = 1 - 0 = 1$$

$$\Rightarrow \boxed{S = \sum_{k=1}^{\infty} \frac{1}{k(k+1)} = 1}$$

we proved it converged! we even found what it converged to!

Divergence Test

- If $\lim_{k\to\infty} a_k \neq 0$, then $\sum_{k=1}^{\infty} a_k$ diverges.
- If $\lim_{k\to\infty} a_k = 0$, then $\sum_{k=1}^{\infty} a_k$ may or may not converge.

Property of Series

Theorem

Suppose that $\sum_{k=1}^{\infty} a_k$ *and* $\sum_{k=1}^{\infty} b_k$ *converge. For any real constants* α *and* β

$$\sum_{k=1}^{\infty} (\alpha a_k + \beta b_k) = \alpha \sum_{k=1}^{\infty} a_k + \beta \sum_{k=1}^{\infty} b_k.$$

Dominating Series

Theorem

If a series $\sum_{k=1}^{\infty} b_k$ *dominates a series* $\sum_{k=1}^{\infty} a_k$ *in the sense that for all sufficiently large* k, $|a_k| \leq b_k$, *then convergence of* $\sum_{k=1}^{\infty} b_k$ *implies convergence of* $\sum_{k=1}^{\infty} a_k$.

Dominating Series Example

Example

Show that the series $\sum_{k=1}^{\infty} \frac{\sin k}{2^k}$ converges.

Integral Test

Theorem (Integral Test)

Suppose f is continuous and $f(k) = a_k$.

(a) *If $|a_k| \leq f(x)$ for all sufficiently large k and all x in the interval $(k-1, k]$, then convergence of $\int_1^\infty f(x)\,dx$ implies convergence of $\sum_{k=1}^{\infty} a_k$.*

(b) *If $|f(x)| \leq a_k$ for all sufficiently large k and all x in the interval $[k, k+1)$ then diverges of $\int_1^\infty f(x)\,dx$ implies divergence of $\sum_{k=1}^{\infty} a_k$.*

Example (p-Series)

Example

Prove that $\sum_{k=1}^{\infty} \frac{1}{k^p}$ converges for $p > 1$ and diverges for $p \leq 1$.

Alternating Series Test

Theorem

Suppose the alternating series $\sum_{k=1}^{\infty}(-1)^{k+1}b_k$ *is such that* $0 \leq b_{k+1} \leq b_k$ *for sufficiently large* k*. Then the series converges if* $\lim_{n\to\infty} b_k = 0$*.*

Alternating Series Test Example

Example

Show that the series $\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k}$ converges.

Absolutely and Conditionally Convergent Series

Definition

A series $\sum_{k=1}^{\infty} a_k$ is **absolutely convergent** if $\sum_{k=1}^{\infty} |a_k|$ converges. A series is **conditionally convergent** if $\sum_{k=1}^{\infty} a_k$ converges but $\sum_{k=1}^{\infty} |a_k|$ does not.

For example, the series

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k}$$

is conditionally convergent but not absolutely.

Ratio Test

Theorem

(a) *If* $\lim_{k\to\infty}\left|\frac{a_{k+1}}{a_k}\right| = L < 1$, *then the series* $\sum_{k=1}^{\infty} a_k$ *is absolutely convergent.*

(b) *If* $\lim_{n\to\infty}\left|\frac{a_{k+1}}{a_k}\right| = L > 1$, *then the series* $\sum_{k=1}^{\infty} a_k$ *is divergent.*

(c) *If* $\lim_{n\to\infty}\left|\frac{a_{k+1}}{a_k}\right| = L = 1$, *then the test fails.*

Ratio Test Example

Example

What can be said about the convergence of $\sum_{k=1}^{\infty}(-1)^k \frac{k!}{k^k}$?

Power Series

Definition

A **power series** centered at c is a series of the form

$$\sum_{k=0}^{\infty} a_k (x-c)^k.$$

If the series converges for $|x-c| < R$ and diverges for $|x-c| > R$, then R is the **radius of convergence**. The **interval of convergence** I is the set of all x values where the series converges.

Remark: We assume $0^0 = 1$ within our power series, so the power series always converges at $x = c$.

Power Series Example

Example

Find the radius and interval of convergence of the series
$$\sum_{k=0}^{\infty} \frac{(-3)^k (x+1)^k}{\sqrt{k+1}}.$$

Differentiation and Integration of Power Series

Theorem

If the power series $f(x) = \sum_{k=0}^{\infty} a_k (x-c)^k$ *has radius of convergence* $R > 0$, *then both*

(a) $f'(x) = \sum_{k=1}^{\infty} k a_k (x-c)^{k-1}$ *and*

(b) $\int f(x)\, dx = C + \sum_{k=0}^{\infty} a_k \frac{(x-c)^{k+1}}{k+1}$

have radii of convergence R.

Differentiation of Power Series Example

Example

$$\sum_{k=1}^{\infty} \frac{k}{1.10^k} =$$

Taylor's Theorem

Theorem

Suppose

$$f(x) = \sum_{k=0}^{\infty} a_k (x-c)^k \qquad \textit{for} \qquad |x-c| < R.$$

Then

$$a_k = \frac{f^{(k)}(c)}{k!}.$$

Popular Taylor Series Centered at Zero

- $\dfrac{1}{1-x} = \sum_{k=0}^{\infty} x^k$ for $x \in (-1, 1)$
- $e^x = \sum_{k=0}^{\infty} \dfrac{x^k}{k!}$ for $x \in \mathbb{R}$
- $\sin x = \sum_{k=0}^{\infty} (-1)^k \dfrac{x^{2k+1}}{(2k+1)!}$ for $x \in \mathbb{R}$
- $\cos x = \sum_{k=0}^{\infty} (-1)^k \dfrac{x^{2k}}{(2k)!}$ for $x \in \mathbb{R}$
- $\arctan x = \sum_{k=0}^{\infty} (-1)^k \dfrac{x^{2k+1}}{2k+1}$ for $x \in [-1, 1]$

Taylor's Theorem Example

Example

Prove $\arctan x = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{2k+1}$ for $x \in [-1, 1]$.

Time Value of Money

Time Value of Money

For a time t cash flow C_t discounted at rate r, the *present value* is

$$PV = \frac{C_t}{(1+r)^t}.$$

The time T *future value* is

$$FV_T = PV \cdot (1+r)^T = C_t \cdot (1+r)^{T-t}.$$

Compound Interest

We assumed that interest is compounded once per unit of time. However, if it is compounded n times per unit of time the formulas become

$$PV = \frac{C_t}{\left(1+\frac{r}{n}\right)^{nt}} \qquad \text{and} \qquad FV_T = PV \cdot \left(1+\frac{r}{n}\right)^{nT}.$$

Since

$$\lim_{n\to\infty}\left(1+\frac{r}{n}\right)^{nt} = e^{rt}$$

for continuous compounding (i.e. $n=\infty$) the formulas are

$$PV = C_t e^{-rt} \qquad \text{and} \qquad FV_T = PV \cdot e^{rT} = C_t \cdot e^{r(T-t)}$$

Multiple Cash Flows

Suppose we have a sequence of cash flows $C_0, C_1, C_2, \ldots, C_T$, where the subscript denotes the time of the cash flow, the *net present value* (NPV) of these cash flows discounted at the constant rate r is

$$NPV = C_0 + \frac{C_1}{1+r} + \frac{C_2}{(1+r)^2} + \ldots + \frac{C_n}{(1+r)^T}.$$

Time Value of Money Python Example

Example

Time	0	1	2	3	4
Cash Flow	-100	50	20	70	10

Calculate the net present value given a continuously compounded discount rate of 5%.

Time Value of Money Python Solution

```python
# Import module
import numpy as np

# Record rate
rate = 0.05

# Record time of cash flows
time = np.array([0, 1, 2, 3, 4])

# Record cash flows
cash_flows = np.array([-100, 50, 20, 70, 10])

# Get the NPV
NPV = np.sum(cash_flows * np.exp(-rate * time))

print(f'The NPV of the cash flows is {NPV:.2f}.')
```

$$NPV \approx 34.10$$

numpy_finance Module

There is a numpy_finance module that has a net present value function (https://numpy.org/numpy-financial/latest/npv.html). However, we would need the annual rate to use it in the last example, i.e. we would have to use the rate

$$100\% \times (e^{0.05} - 1) \approx 5.127\%.$$

Time Value of Money Example

Example

Jain borrows \$1,000,000 to purchase a house. The loan is for thirty years and her first payment is one month from when she initially borrows the money. If her annualized rate is 12%, what will be her monthly payments? Ignore fees.

Growing Payments

Suppose 1 is payed at time 1, and payments increase at a rate of g each subsequent period until a final payment of $(1+g)^{n-1}$ is made at time n. If cash flows are discounted at rate r, then the NPV of the cash flows is

$$\frac{1-\left(\frac{1+g}{1+r}\right)^n}{r-g}.$$

Growing Payments Example

Example

Calculate the NPV of the series of end-of-year cash flows. Assume

- \$100 is payed in the first year,
- each subsequent year payments increase by 5%,
- the final payment is made at the end of year ten, and
- the discount rate is 8%.