

**DS-GA 3001.001 Special Topics in Data Science: Modeling Time Series**  
**Homework 1**

**Due date: February 12, by 5 pm**

**Problem 1.** Consider the periodic time series (period determined by parameter  $\omega$ ) constructed as:

$$x_t = U_1 \sin(2\pi\omega t) + U_2 \cos(2\pi\omega t), \quad (1)$$

with  $U_{1,2}$  independent random variables with zero mean and variance  $\sigma^2$ . Show that this series is weakly stationary with autocovariance function  $\gamma(h) = \sigma^2 \cos(2\pi\omega h)$ . Optional: Is this series strictly stationary?

*Solution:*

First we need to show that the mean of the series is time independent:

$$\begin{aligned} \mathbb{E}[x_t] &= \mathbb{E}[U_1 \sin(2\pi\omega t) + U_2 \cos(2\pi\omega t)] \\ &= \mathbb{E}[U_1] \sin(2\pi\omega t) + \mathbb{E}[U_2] \cos(2\pi\omega t) \\ &= 0 \end{aligned}$$

where we have used the fact that the mean of  $U_{1,2}$  is zero.

Second, we need to show that the autocovariance function depends only on the time difference  $h$ :

$$\begin{aligned} \mathbb{E}[x_t x_{t+h}] &= \mathbb{E}[(U_1 \sin(2\pi\omega t) + U_2 \cos(2\pi\omega t))(U_1 \sin(2\pi\omega(t+h)) + U_2 \cos(2\pi\omega(t+h)))] \\ &= \mathbb{E}[U_1^2] \sin(2\pi\omega t) \sin(2\pi\omega(t+h)) \\ &\quad + \mathbb{E}[U_1 U_2] \sin(2\pi\omega t) \cos(2\pi\omega(t+h)) \\ &\quad + \mathbb{E}[U_1 U_2] \sin(2\pi\omega(t+h)) \cos(2\pi\omega t) \\ &\quad + \mathbb{E}[U_2^2] \cos(2\pi\omega t) \cos(2\pi\omega(t+h)) \\ &= \sigma^2 (\cos(2\pi\omega t) \cos(2\pi\omega(t+h)) + \sin(2\pi\omega t) \sin(2\pi\omega(t+h))) \\ &= \sigma^2 \cos(2\pi\omega h) \end{aligned}$$

where we have used the fact that  $U_{1,2}$  are independent (mixed terms are zero), their second moment is  $\mathbb{E}[U_i^2] = \sigma^2 - \mathbb{E}[U_i]^2 = \sigma^2$ , and lastly the trigonometric formula  $\cos(a-b) = \cos(a)\cos(b) + \sin(a)\sin(b)$ .

Note: Since the other moments of the distribution are unspecified, we cannot prove that the series is strictly stationary.

**Problem 2.** For an MA(1),  $x_t = w_t + \theta w_{t-1}$  show that the autocorrelation function  $|\rho_x(1)| \leq 0.5$ , for all  $\theta$ . For which values  $\theta$  is it maximum and minimum?

*Solution:*

We have already computed the autocovariance for an MA(1) in Lecture 2 (but bonus points if you've included the full derivation):

$$R_X(h) = \begin{cases} \sigma^2(1 + \theta^2) & , h = 0 \\ \sigma^2\theta & , |h| = 1 \\ 0, & |h| > 1 \end{cases}$$

After the normalization, the autocorrelation function becomes:

$$\rho_X(h) = \begin{cases} 1 & , h = 0 \\ \frac{\theta}{1+\theta^2} & , |h| = 1 \\ 0, & |h| > 1 \end{cases}$$

To prove that  $|\rho_x(1)| \leq 0.5$ , we multiply both sides by the denominator and rearrange all terms to the right, giving  $\theta^2 - 2|\theta| + 1 \geq 0$  which can be rewritten as  $(|\theta| - 1)^2 \geq 0$  q.e.d. To find the extrema points, we take the first derivative and set it to zero, which gives:  $\frac{1-\theta^2}{(1+\theta^2)^2} = 0$ , with roots  $\theta_{\min/\max} = \pm 1$  and the corresponding optima  $\pm 0.5$ .

**Problem 3.** Identify the following models as ARMA( $p, q$ ):

- $x_t = 0.8x_{t-1} - 0.15x_{t-2} + w_t - 0.3w_{t-1}$
- $x_t = x_{t-1} - 0.5x_{t-2} + w_t - w_{t-1}$

Note: watch out for parameter redundancy!

*Solution:*

We first write the expression in canonical form (AR terms left of the = MA terms to the right),  $x_t - 0.8x_{t-1} + 0.15x_{t-2} = w_t - 0.3w_{t-1}$ , then identify the corresponding P,Q polynomials :

$$\begin{aligned} P(B) &= 1 - 0.8B + 0.15B^2 = (1 - 0.3B)(1 - 0.5B) \\ Q(B) &= 1 - 0.3B \end{aligned}$$

Since the two share a factor, we can simplify the definition of the model to

$$\begin{aligned} P(B) &= 1 - 0.5B \\ Q(B) &= 1 \end{aligned}$$

which translates into  $x_t = 0.5x_{t-1} + w_t$  which is an AR(1) process. Note: P's root is  $2 > 1$ , so the process is causal.

Similarly, the second model  $x_t - x_{t-1} + 0.5x_{t-2} = w_t - w_{t-1}$  has corresponding polynomials

$$\begin{aligned} P(B) &= 1 - B + 0.5B^2 \\ Q(B) &= 1 - B \end{aligned}$$

These don't have common roots so the model is an ARMA(2,1); P has complex roots  $1 \pm i$  which are outside the unit circle, so it's a causal process. Q has root 1 so it's not invertible.

**Problem 4.** Given an AR(1) model  $x_t = \phi x_{t-1} + w_t$ , determine the form of m-step ahead prediction  $x_{t+m}^t = \mathbb{E}[x_{t+m}|x_t]$  and show that the variance of this estimate has the form:

$$\mathbb{E}[(x_{t+m}^t - x_{t+m})^2] = \sigma_w^2 \frac{1 - \phi^{2m}}{1 - \phi^2} \quad (2)$$

*Solution:*

The joint distribution for  $(x_t, x_{t+m})$  is multivariate normal with mean  $[0, 0]$  and covariance  $\begin{bmatrix} R_x(0) & R_x(m) \\ R_x(m) & R_x(0) \end{bmatrix}$

where the AR(1) autocovariance is  $R_x(h) = \sigma^2 \frac{\phi^h}{1 - \phi^2}$  (from lecture or rederived).

The posterior distribution  $x_{t+m}|x_t$  is also gaussian with mean  $\mu_x = \phi^m x_t$  and variance  $R_x(0) - R_x(m)^2/R_x(0)$  (using the multivariate gaussian conditioning formula). After replacing in the expression for the covariance this finally gives the variance we were looking for:

$$\text{Var}(x_{t+m}|x_t) = \frac{\sigma^2}{1 - \phi^2} - \left( \sigma^2 \frac{\phi^m}{1 - \phi^2} \right)^2 \left( \frac{\sigma^2}{1 - \phi^2} \right)^{-1} = \frac{\sigma^2}{1 - \phi^2} (1 - \phi^{2m}) \quad (3)$$

**Problem 5.** Consider the datasets recording weekly *oil* (dollars/barrel) and *gas* (cents/gallon) provided in the data folder.

- plot data on same graph; which of the simulated series in lecture 1 do these series resemble? Do you believe they are stationary? (explain why/ why not)

- argue that the transformation  $y_t = \nabla x_t$  might be useful for the oil data. Plot  $y_t$ , estimate sample ACF and comment on result.
- construct scatterplot of  $y_t$  vs.  $y_{t+h}$  for the two datasets,  $h = \{1, 2, 3\}$ . Are these dependencies linear?

Anything sensible is fine here, but it would be nice to see some attempts to check the statistical structure beyond simple visualisation - e.g. fitting a regression model for the integrated part , or looking at the histogram of residuals to check for gaussianity or any other sensible rationalization of the form "if data were coming from an ARMA model then... "