HOBSONVILLE ULTRASOUND

Monday to Friday: 9:30am - 5:30pm
Saturday: 9:00am - 12:00pm
+64 986 93328
reception@hobus.co.nz
www.hobus.co.nz
124 Hobsonville Road, Hobson Centre,
Level 1 (Upstairs), Hobsonville, Auckland, 0618

Patient Information

Name

Address

NHI Number (Required)

Date of Birth

Contact Phone

ACC Number

Services

General Ultrasound
- ☐ Abdomen
- ☐ Female pelvis
- ☐ Male Pelvis
- ☐ Renal Tract
- ☐ Scrotum
- ☐ Thyroid/Neck
- ☐ Breast
- ☐ Lumps & bumps / Soft Tissue
- ☐ Foreign Body

Musculoskeletal Ultrasound
- ☐ Shoulder
- ☐ Elbow
- ☐ Wrist
- ☐ Knee
- ☐ Ankle
- ☐ Groin / Hernia
- ☐ Abdominal wall / Hernia

Pregnancy Ultrasound
- ☐ 1st Trimester
- ☐ Nuchal Translucency
- ☐ 2nd Trimester / Anatomy
- ☐ 3rd Trimester / Growth
- ☐ Post-partum
- ☐ 4D Pregnancy (non-diagnostic)

Vascular Ultrasound
- ☐ Leg veins DVT
- ☐ Arm veins DVT
- ☐ Carotid Doppler
- ☐ Aorta

Paediatric Ultrasound
- ☐ Abdomen
- ☐ Renal
- ☐ Scrotum

Clinical Information


Tick if urgent ☐

LMP:

EDD:

Map

124 Hobsonville Rd, Hobsonville



Referral Information

Name

Date

Send copies to

Signature

Patient Preparation

Abdominal Ultrasound
Avoid food for 6 hours prior to your scan. You may only drink water. Take medication as usual but consult your doctor if you're diabetic.

First Trimester/Nuchal Translucency/Female & male pelvis/ Renal Ultrasound. A full bladder is required for this scan.
To prepare for the ultrasound, empty your bladder two hours before the exam. Then, drink 4-5 glasses of water 40-60 minutes before the scan. Avoid emptying your bladder after drinking water.

Maternity patients:
After 14 weeks you do not need a full bladder for your scan.