

# STIMULAN®

## POWER TO TRANSFORM OUTCOMES

### CASE STUDY

#### Courtesy of Mr Alan Norrish

Consultant Orthopaedic Surgeon, Mr Dimitrios Giotikas, Specialist Orthopaedic Surgeon, Addenbrooke's Hospital, Cambridge, UK

#### Clinical particulars

35-year-old male presented with recurrent periprosthetic infection after complex hinged total knee replacement (TKR), performed 1 year ago, to treat severe post-traumatic knee osteoarthritis. 6 months after the initial procedure he developed periprosthetic infection. Change of polyethylene, synovectomy, microorganism specific systemic antibiotics and repetitive arthroscopic lavage of the knee failed to control the infection.

Because of the knee stiffness and the compromised soft tissue condition, a revision TKR was not deemed appropriate so the decision was made to treat the infection with knee fusion.

#### Treatment

In order to preserve the length of the leg, a two-stage approach with the induced-membrane-technique was implemented. First stage: the implants were removed, necrotic and infected bone was excised, and the knee was temporarily fused with a long Trigen nail. Normal length was preserved and the bone void filled with a PMMA spacer. Systemic antibiotics were administered. Second stage was conducted 3 months later, after clinical control of infection and normalization of CRP and ESR values. The nail was exchanged to an osteobridge nail and bone graft was harvested with Reamer-Irrigator-Aspirator from the contralateral femur. The bone graft and 10cc of **STIMULAN** beads were inserted to fill the bone void. Vancomycin and gentamicin were used to treat the infection.

#### Outcome

7 months' post-operatively, the infection remained controlled, the bone had consolidated, and the patient had returned to daily activities without significant restrictions or symptoms. 2 years' post-operatively, no recurrence of infection had occurred.



Presentation



Presentation



First stage –  
Post-operative



Second stage –  
Post-operative



7 months

For indications, contraindications, warnings and precautions see Instructions for Use. The treating physician is responsible for deciding the type and quantity of antibiotic used. Concurrent use of locally administered antibiotics may affect setting time. The mixing of antibiotics with the STIMULAN Kit/ STIMULAN Rapid Cure device is considered off-label usage of the medicinal product. To do so is at the professional risk of the surgeon / healthcare professional. This brochure may include the use of STIMULAN or techniques that go beyond the current clearance/ approval granted by the relevant regulatory authority. Please contact your local representative for further information.

©2017, Biocomposites and STIMULAN are registered trademarks of Biocomposites Ltd. All rights reserved. No unauthorized copying, reproduction, distributing or republication is allowed unless prior written permission is granted by the owner, Biocomposites Ltd.

Patents granted: GB2367552, EP 1204599 B1, US 6780391, EP 2594231 B1, US 8883063, CN ZL201210466117.X, GB2496710 / Patents pending: GB1502655.2, EP 16154210.5, US 15/040075, CN 201610089710.5, US 15/288328.

MA0154RI. Canada only. Not for distribution in the USA.