Contrasts

Peter von Rohr

2023-03-27

Contrasts

- ► Linear combination of parameters
- ▶ In R used to determine which estimable functions are shown as factor level effects

Example Dataset

Table 1: Body Weight and Breed of Beef Cattle Animals

Animal	Body Weight	Breed
1	471	Angus
2	463	Angus
4	470	Angus
7	518	Limousin
8	511	Limousin
9	510	Limousin
10	541	Limousin
3	481	Simmental
5	496	Simmental
6	491	Simmental

Contrasts in R

Angus
Limousin
Simmental

```
(mat_ctr <- contrasts(as.factor(tbl_flem_bw_breed$Breed)))
##
        Limousin Simmental</pre>
```

Model Matrix

IVIC	Model Matrix					
	mod	lel.	matrix(lm(`	Body Weight` ~	Breed, data = t	bl_flem_bw_b
	##		(Intercept)	${\tt BreedLimousin}$	${\tt BreedSimmental}$	
	##	1	1	0	0	
	##	2	1	0	0	
	##	3	1	0	0	
	##	4	1	1	0	
	##	5	1	1	0	
	##	6	1	1	0	
	##	7	1	1	0	
	##	8	1	0	1	
	##	9	1	0	1	

10 ## attr(,"assign") ## [1] 0 1 1 ## attr(,"contrasts") ## attr(,"contrasts")\$Breed ## [1] "contr.treatment"

Estimable Functions

extend contrasts matrix by one row of all ones for the intercept

##			Limousin	Simmental
##	Angus	1	0	0
##	Limousin	1	1	0
##	Simmental	1	0	1

Estimable Functions II

Inverse of extended contrasts matrix

##		Angus	Limousin	Simmental
##		1	0	0
##	Limousin	-1	1	0
##	Simmental	-1	0	1

- First row: which group means are used for intercept
- \triangleright Other rows: vectors q^T representing estimable functions

Validation

. . .

Other Contasts

- ► Helmert
- sum

Custom Contrasts