

Interactions

Peter von Rohr

2023-04-03

Definition

- ▶ Effect of given predictor variable depends on level or value of other predictor variable
- ▶ Examples:
 - ▶ Regression of Body Weight on Breast Circumference is different for different breeds
 - ▶ Effect of Breed on Body Weight is different for different male and female animals


Types of Interactions

- ▶ continuous by continuous
- ▶ continuous by categorical
- ▶ categorical by categorical

Continuous by Categorical

- ▶ In a model, expected value of response depends on regression variable plus discrete factor
- ▶ Example: Regression of Body Weight on Breast Circumference plus the factor Breed
- ▶ Interaction is present, if regression of Body Weight on Breast Circumference is different for different breeds

Interaction Plot



Interaction Model

- ▶ Start with model without interactions

$$y_i = b_0 + b_1 \times BC_i + b_2 \times BrLi_i + b_3 \times BrSi_i + e_i$$

- ▶ Assume linear relationship of b_1 with Breed

$$b_1 = a + b_4 \times BrLi + b_5 \times BrSi$$

- ▶ Insert

$$y_i = b_0 + (a + b_4 \times BrLi + b_5 \times BrSi) \times BC_i + b_2 \times BrLi_i + b_3 \times BrSi_i + e_i$$

- ▶ Simplify

$$\begin{aligned} y_i = & b_0 + a \times BC_i + b_2 \times BrLi_i + b_3 \times BrSi_i \\ & + b_4 \times BrLi \times BC_i + b_5 \times BrSi \times BC_i + e_i \end{aligned}$$

Continuous by Continuous

- ▶ Similar to continuous by categorical
- ▶ No interaction

$$y_0 = b_0 + b_1 \times BC_i + b_2 \times HE_i + e_i$$

- ▶ Interaction by dependence of one regression coefficient on other coefficient

$$b_1 = b_3 + b_4 \times HE_i$$

$$y_0 = b_0 + b_2 \times HE_i + b_3 \times BC_i + b_4 \times HE_i \times BC_i + e_i$$

Categorical by Categorical

