

Inverse Numerator Relationship Matrix with Inbreeding

Peter von Rohr

2020-10-30

Inbreeding

- ▶ Elements in matrix D depend on coefficients of inbreeding
- ▶ Recap: From the simple decomposition of a , we derived

$$\begin{aligned} \text{var}(m_i) &= \left(\frac{1}{2} - \frac{1}{4}(F_s + F_d) \right) \sigma_a^2 \\ &= \left(\frac{1}{2} - \frac{1}{4}(A_{ss} - 1 + A_{dd} - 1) \right) \sigma_a^2 \\ &= \left(1 - \frac{1}{4}(A_{ss} + A_{dd}) \right) \sigma_a^2 \\ &= (D)_{ii} \sigma_a^2 \end{aligned}$$

$$\rightarrow (D)_{ii} = \left(\frac{1}{2} - \frac{1}{4}(F_s + F_d) \right) = \left(1 - \frac{1}{4}(A_{ss} + A_{dd}) \right)$$

Computation of Coefficients of Inbreeding

- ▶ Observation: Coefficients of inbreeding F_s and F_d can be read from $(A)_{ss}$ and $(A)_{dd}$ of A
- ▶ Cannot setup A to just get inbreeding coefficients
- ▶ More efficient method required
- ▶ **Cholesky** decomposition of A

$$A = R \cdot R^T$$

where R is a lower triangular matrix

Hint: Function `chol(A)` in R gives matrix R^T

Cholesky Decomposition

- ▶ Diagonal elements $(A)_{ii}$ of A are the sum of the squared elements of one row of R

$$(A)_{ii} = \sum_{j=1}^i (R)_{ij}^2$$

- ▶ Example

$$\begin{bmatrix} (A)_{11} & (A)_{12} & (A)_{13} \\ (A)_{21} & (A)_{22} & (A)_{23} \\ (A)_{31} & (A)_{32} & (A)_{33} \end{bmatrix} = \begin{bmatrix} (R)_{11} & 0 & 0 \\ (R)_{21} & (R)_{22} & 0 \\ (R)_{31} & (R)_{32} & (R)_{33} \end{bmatrix} \cdot \begin{bmatrix} (R)_{11} & (R)_{21} & (R)_{31} \\ 0 & (R)_{22} & (R)_{32} \\ 0 & 0 & (R)_{33} \end{bmatrix}$$

Recursive Computation of R

- ▶ Let us write the matrix R as a product of two matrices L and S :

$$R = L \cdot S$$

where L is the same matrix as in the LDL-decompositon and S is a diagonal matrix.

- ▶ Compute A as

$$A = R \cdot R^T = L \cdot S \cdot S \cdot L^T = L \cdot D \cdot L^T$$

- ▶ Hence

$$D = S \cdot S \quad \rightarrow \quad (S)_{ii} = \sqrt{(D)_{ii}}$$

Example

$$\begin{bmatrix} (R)_{11} & 0 & 0 \\ (R)_{21} & (R)_{22} & 0 \\ (R)_{31} & (R)_{32} & (R)_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ (L)_{21} & 1 & 0 \\ (L)_{31} & (L)_{32} & 1 \end{bmatrix} \cdot \begin{bmatrix} (S)_{11} & 0 & 0 \\ 0 & (S)_{22} & 0 \\ 0 & 0 & (S)_{33} \end{bmatrix}$$

- ▶ Diagonal elements $(R)_{ii} = (S)_{ii}$
- ▶ Because $(S)_{ii} = \sqrt{(D)_{ii}}$, if parents s and d are known diagonal elements $(R)_{ii}$ of matrix R can be computed as

$$(R)_{ii} = (S)_{ii} = \sqrt{(D)_{ii}} = \sqrt{\left(1 - \frac{1}{4}(A_{ss} + A_{dd})\right)}$$

- ▶ A_{ss} and A_{dd} are
 - ▶ 0 if s and d are unknown (NA) or
 - ▶ have been computed before

Recap matrix D

- ▶ Both parents s and d of animal i are known

$$(D)_{ii} = \frac{1}{2} - \frac{1}{4}(F_s + F_d) = \frac{1}{2} - \frac{1}{4}((A)_{ss} - 1 + (A)_{dd} - 1) = 1 - \frac{1}{4}((A)_{ss} + (A)_{dd})$$

- ▶ Parent s of animal i is known

$$(D)_{ii} = \frac{3}{4} - \frac{1}{4}F_s = \frac{3}{4} - \frac{1}{4}((A)_{ss} - 1) = 1 - \frac{1}{4}(A)_{ss}$$

- ▶ Both parents unknown

$$(D)_{ii} = 1$$

Offdiagonal Elements of R

- ▶ Offdiagonal elements $(R)_{ij}$ of R are computed as

$$(R)_{ij} = (L)_{ij} * (S)_{jj}$$

- ▶ Use property of L : $L_{ij} = \frac{1}{2}((L)_{sj} + (L)_{dj})$ if s and d are parents of i

$$\begin{aligned}(R)_{ij} &= (L)_{ij} * (S)_{jj} \\&= \frac{1}{2} [(L)_{sj} + (L)_{dj}] * (S)_{jj} \\&= \frac{1}{2} [(L)_{sj} * (S)_{jj} + (L)_{dj} * (S)_{jj}] \\&= \frac{1}{2} [(R)_{sj} + (R)_{dj}]\end{aligned}$$

Example Pedigree

Calf	Sire	Dam
1	NA	NA
2	NA	NA
3	NA	NA
4	1	2
5	3	2
6	4	5

Computations

- ▶ Compute diagonal elements $(A)_{ii}$ of A to get F_i
- ▶ Prerequisite: Pedigree sorted such that parents before progeny
- ▶ Start with $(A)_{11}$

$$(A)_{11} = (R)_{11}^2 = (D)_{11} = 1$$

- ▶ $(A)_{22} = (R)_{21}^2 + (R)_{22}^2 = 0 + 1 = 1$
- ▶ $(A)_{33} = (R)_{31}^2 + (R)_{32}^2 + (R)_{33}^2 = 0 + 0 + 1 = 1$

Animals With Known Parents

$$\begin{aligned}(A)_{44} &= (R)_{41}^2 + (R)_{42}^2 + (R)_{43}^2 + (R)_{44}^2 \\&= \left(\frac{1}{2}(R_{11} + R_{21})\right)^2 + \left(\frac{1}{2}(R_{12} + R_{22})\right)^2 + \left(\frac{1}{2}(R_{13} + R_{23})\right)^2 \\&\quad + \left(1 - \frac{1}{4}(A_{11} + A_{22})\right) \\&= \frac{1}{4} + \frac{1}{4} + \frac{1}{2} = 1\end{aligned}$$

► $(A)_{55}$

► $(A)_{66}$