

A Study of Active Galactic Nuclei With SDSS DR7

Charlotte Hamilton
University of California, Los Angeles
cmhamilton@g.ucla.edu

April 2026

Abstract

It has long been established that massive galaxies hosting accreting supermassive black holes produce extreme emission lines from their centers. These centers are incredibly small (usually only a bit larger than the solar system), yet produce enough energy to match or outshine the luminosity of the galaxy as a whole. This study will examine the emission line properties of galaxies and determine if they are hosting these active galactic nuclei (AGN), and compare certain properties to those of normal starforming galaxies.

1 An Introduction

1.1 What are Active Galactic Nuclei?

1908 saw the very first discovery of a very peculiar galaxy, as astronomer Edward A. Fath was conducting long exposure photographic spectra studies of possible extragalactic objects¹. He wanted to test whether these objects, now known to be spiral galaxies, would produce stellar absorption lines consistent with a collection of stars, and for the large majority, this was true. But for one particular object, NGC 1068, it showed not only absorption lines but very bright emission lines.

NGC 1068 would later prove to be the first documented case of an active galactic nucleus (AGN), some of the most persistently energetic sources in the universe. The term currently refers to a compact central region of a galaxy that produces extremely luminous and high energy phenomena caused by an accreting supermassive black hole (SMBH). The SMBH is an engine that fuels intense energy production, and results in features such as the relativistic jets, winds, and broad UV/optical emission lines seen in the host galaxies. They have become an area of intense study for astrophysicists, as they provide a unique environment for high energy physics and galactic evolution.

Currently, there are still many unanswered questions in the field. This study will focus on some of the smaller ones and attempt to shed some light on the mysteries of these massive, high energy nuclei.

1.2 Areas of Study

This study will mainly focus on the identification of AGN through emission line ratios. There will also be an examination of the properties of the identified AGN and a comparison of them with proper starforming galaxies.

¹At the time, spiral galaxies were not yet proved to be their own “island universes,” they were still widely believed to be nebulae within the Milky Way. Fath was one of the first to note the true nature of these spiral nebulae.

Starburst galaxies produce extremely strong emission lines of highly ionized species, such as [Fe VIII] or [Ne V], which need a huge amount of energy to lose so many electrons. One of the strongest indicators specifically that a galaxy is hosting an AGN, rather than just being a normal starburst galaxy, is the strength of its Balmer lines. Any starburst galaxy or AGN will intrinsically have a $H\alpha$ to $H\beta$ line ratio of 3, but AGN will have much stronger and broader lines than that of a starburst. However, because $H\beta$ struggles to escape from dust-heavy regions, the extinction of $H\beta$ is much higher than that of $H\alpha$, leading to a much larger observed $H\alpha / H\beta$ ratio, with the $H\alpha$ line sometimes being greater than 5 times stronger than the $H\beta$ line.

In order to counteract the effects of dust, “extinction independent” line ratios are chosen to discriminate against the excitation mechanism of the galaxy. These species experience extinction extremely similarly to $H\alpha$ and $H\beta$. The standard emission lines ratios chosen are $[O III]/H\beta$, $[N II]/H\alpha$, $[S II]/\alpha$, and $[O I]/H\alpha$, with the brackets denoting forbidden lines. This study will focus on the line ratios $[O III]/H\beta$ and $[N II]/H\alpha$, following Kauffman et al. (2003).

There is certain criteria that the emission line ratios of a galaxy must meet in order to be classified as hosting an AGN, the most widely established being that of Kauffman et al. 2003 and Kewley et al. 2001. In order to visualize the data classification, BPT diagrams will be made following the standards in these papers. BPT diagrams refer to the diagrams laid out by Baldwin, Phillips, & Terlevich (1981) that have the emission line ratios on each axis and allow for the easy categorization of galaxies depending on where they lie on the diagram. The specific classification scheme used in this study does not follow the Baldwin, Phillips, & Terlevich paper.

After classification, the two chosen galactic properties that will be considered and compared are the stellar masses and luminosity of the $[O III]$ line, which can track the strength of activity in the nucleus.

2 The Spectroscopic & Photometric Sample

This study needs a large and uniform sample of galaxies with both spectroscopic and photometric abilities, which is why the Sloan Digital Sky Survey (SDSS) public data access catalogs were chosen. SDSS provides very high quality photometric and spectroscopic data of a very large sample of galaxies. In particular, Data Release 7 (DR7), which was in collaboration with MPA/JHU, had a focus on the statistical properties of galaxies and AGN, with value-added catalogs specifically designed for these studies. It has large datasets that cover over 45,000 square degrees of sky and over 1 million galaxies, containing in-depth absorption and emission line spectra, stellar masses, and star formation rates.

2.1 SDSS Telescope

The survey utilized a modified Ritchey-Chretien telescope at the Apache Point Observatory with imaging and spectroscopy abilities – a large-mosaic CCD camera that can image with 5 optical bands and two spectrographs (one with a blue channel and one with a red channel). The CCDs are specifically set up to in six columns of five each, with each row corresponding to the SDSS r, i, u, z, and g filters. The telescope used a drift scan mode that has the telescope moving in circles across the sky that has images of objects being collected at the same rate as the CCD data is being read. This allows for five images of each object (one per filter). The spectroscopic instrument has cartridges containing plates with drilled holes plugged with optical fibers mounted onto the telescope. Exposures are collected on the plates, at least three per object.




Figure 1: Visualization of SDSS Raw Data Processing Activity Flow

2.2 Data Processing

SDSS then did raw data processing, as visualized in Figure (1), and released the raw data in FITS files, which MPA/JHU used to create derived datasets. The derived datasets contain gas-phase metallicity estimates, star formation rates, and stellar masses. Stellar mass calculations follow Kaufmann et al (2003), star formation rates calculations follow Brinchmann et al (2004), and gas phase metallicities follow Tremonti et al (2004).

2.3 The Catalogs

The files used are the `gal_info`, `gal_line`, and `stellarmasses` files as found on the MPA/JHU DR7 Release of Spectrum Measurements website. This website contains all the value added galaxy catalogs as custom galaxy spectral fitting codes of raw data that includes line fluxes, line widths, etc. Both `gal_info` and `gal_line` are part of the raw data FITS files. `Stellarmasses` is a derived data FITS file.

`Gal_info` contains SDSS targeted info from the plugmap and redshift and velocity dispersion, and gives redshifts, velocity dispersions, right ascension, declination. It was chosen due to its redshift columns, which are needed for later luminosity calculations. `Gal_line` contains width and flux measurements of emission lines for 20 different lines in CGS units. `Stellarmasses`, the only file from the derived data, contains log masses calculated from *photometry*, as well as percentile measurements. These files are all integral to calculations that will be done later in this study.

gal_info	gal_line	stellarmasses
RA	OIII_5007_FLUX	MEDIAN
DEC	OIII_5007_FLUX_ERR	P16 ¹
Z	H_BETA_FLUX	P84 ²
Z_WARNING	H_BETA_FLUX_ERR	
SN_MEDIAN	NII_6584_FLUX	
	NII_6584_FLUX_ERR	
	H_ALPHA_FLUX	
	H_ALPHA_FLUX_ERR	

Figure 2: Chosen Columns from `gal_info`, `gal_line`, and `stellarmasses` Files

All data analysis was conducted in Python, using `astropy`, `numpy`, and `matplotlib` libraries. In order to sufficiently analyze the data, specific columns were picked from each file and combined into one singular dataframe using `astropy.table`. Figure (2) depicts which columns were chosen from each file to analyze.

Emission line columns were chosen according to Veilleux & Osterbrock (1986) line ratios of [OIII] $\lambda 5007$ / $H\beta$ and [NII] $\lambda 6583$ / $H\alpha$. Veilleux & Osterbrock follow five criteria for best line ratios for AGN classification, including:

1. Ratios must be made up of strong lines easy to measure in typical spectra.
2. Lines cannot be badly blended with other lines as deblending increases uncertainty.
3. Wavelength separation must be small as to avoid reddening and flux discrepancies.
4. Ratios should contain one element and one HI Balmer line as these are less likely to depend on chemical composition of the galaxy.
5. Lines need to be easily read with present day instrumentation.²

While there are multiple different possible line ratios that fall into these criterion, [OIII] $\lambda 5007$ / $H\beta$ and [NII] $\lambda 6583$ / $H\alpha$ are the strongest candidates, and also follow both Kauffman et al. (2003) and Kewley et al. (2001). These lines specifically are almost completely insensitive to reddening and/or spectrophotometric errors.

The data was then quality cut in order to maximize accuracy. Following Kauffman et al., a signal to noise ratio (S/N) was chosen to be $S/N > 3$ for all four emission lines by $S/N = \frac{flux}{err} > 3$. Redshifts were filtered by the `Z_WARNING` column with bad redshifts having `Z_WARNING` $\neq 0$, which was calculated by MPA/JHU using the Schlegel data model.

Quality cuts also included a chosen redshift range. The redshift range is very important for calculations using emission line fluxes. Higher redshift galaxies have more redshifted emission lines, creating discrepancies in wavelength and which filters can capture the spectrum. Luckily, these

²“Present day instrumentation” here is referring to instrumentation of the year 1986, when Veilleux & Osterbrock was published. By the SDSS DR7 release in 2008, instrumentation had gotten increasingly more sensitive. Nonetheless, the emission lines still follow the criterion as written by Veilleux & Osterbrock to ensure higher accuracy.

catalogs had this issue accounted for by the SDSS data processing team for emission line fluxes and errors, so this would not affect any data analysis in this study. However, a larger issue is that galaxies in different redshift bins are captured in different galactic eras. Much higher redshifted galaxies are still in the early ages of galaxy formation, and therefore AGN would have differing properties. As AGN properties in different redshift bins is not something that is going to be analyzed in this study, a low redshift range of $0.04 < z < 0.2$ was chosen, which follows Kauffman et al.

After quality cuts, 320,759 galaxies were left of the original 927,552 (34.6%).

3 Data Analysis

3.1 Classifications

Classifications were based on Kewley et al. (2001) and Kauffman et al. (2003). Kewley et al. defines a theoretical upper limit for a starburst model in which anything above or to the right of the empirical limit is not a starburst by any current models of metallicity and ionization parameters. Galaxies above this line cannot be explained by any possible combination of parameters, and no normal star-forming HII region can exist above this line, so it must be explained by something else. The limit is defined as

$$\log_{10}([OIII]/H\beta) = \frac{0.61}{\log_{10}([NII]/H\alpha) - 0.47} + 1.19 \quad (1)$$

Kauffman et al., on the other hand, defines a much lower demarcation line between starburst galaxies and AGN. They used very similar, if less conservative, reasoning as Kewley et al., and had an empirical limit defined as

$$\log_{10}([OIII]/H\beta) = \frac{0.61}{\log_{10}([NII]/H\alpha) - 0.05} + 1.3 \quad (2)$$

with galaxies defined to be possible AGN if they satisfy

$$\log_{10}([OIII]/H\beta) > \frac{0.61}{\log_{10}([NII]/H\alpha) - 0.05} + 1.3 \quad (3)$$

For the classification in this study, the Kewley limit was used as an upper limit on starforming galaxies while the Kauffman limit was used as the lower limit for AGN. Galaxies were defined to be starforming if they had

$$\log_{10}([OIII]/H\beta) < \frac{0.61}{\log_{10}([NII]/H\alpha) - 0.05} + 1.3 \quad (\text{The Kauffmann Limit}) \quad (4)$$

and defined to be AGN if they had

$$\log_{10}([OIII]/H\beta) > \frac{0.61}{\log_{10}([NII]/H\alpha) - 0.47} + 1.19 \quad (\text{The Kewley Limit}) \quad (5)$$

For galaxies in between the two lines, or galaxies that were neither defined to be starforming nor AGN, they were classified as a composite group called “questionable”. Questionable galaxies had spectroscopic properties that were classified differently depending on which model was used, and could be either high luminosity starforming or lower luminosity AGN. Figure (3) depicts the resulting BPT diagram. It is also worth noting that though the Kauffman and Kewley equations define certain spectroscopic properties to be the result of AGN only, they could also be the results




Figure 3: BPT Line Ratio Classification Diagram

of shocks. By Kaschinski & Ercolano (2013), shocks are unstable radiation-driven winds that are a result of EUV and X-ray radiation produced from cooling zones. They disrupt spectral energy distributions and produce extra emission lines (though it depends on temperature), resulting in differing photoionization models with shock stellar atmosphere models as an energy source. As a product of this, the AGN classification schema could be affected.

3.2 Properties and Comparisons

In order to study how galactic properties differ between the modeled classes, two particular characteristics were chosen to reveal distribution discrepancies between classified AGN and normal starforming galaxies: stellar mass distribution and [OIII] $\lambda 5007$ emission line luminosity distribution.

The stellar mass **MEDIAN** column was made into an array of $\log_{10}$ masses, and then divided further into each galaxy's classification. Distributions were plotted in a histogram as seen in Figure (4a).

The [OIII] $\lambda 5007$ emission line luminosity distribution needed distance calculations. Distances were derived from redshifts using `astropy.cosmology` and assuming a Flat Lambda CDM cosmological model ($H_0 = 70$, $\Omega_m = 0.3$). Luminosity was then found using the log of the classic equation relating flux, distance, and luminosity, $Flux = \frac{L}{4\pi d^2}$, with L as luminosity and d as the derived distances. Before calculation, fluxes were converted to standard CGS units, as `gal_line` has all flux measurements in $1 \times 10^{-17} \text{ erg/s/cm}^2$. The final calculated luminosities were then made into an array and further divided by each galaxy's classification. A histogram was made depicting the distribution, as seen in Figure (4b).




Figure 4: Galactic Property Comparison Histograms

3.3 Final Measurements by Class

Classification counts and percent fractions of the total sample³ are depicted in Figure (5). Stellar mass distributions by class, including the median, mean, and standard deviation, is depicted in Figure (6). [O III] emission line luminosity distribution by class, including the median, mean, and standard deviation, is depicted in Figure (7).

CLASS	COUNT	FRACTION
Star Forming	202,076	63.0%
Questionable	73,050	22.8%
AGN	45,633	14.2%

Figure 5: Classification Counts and Fractions

CLASS	MEDIAN	MEAN	STD
Star Forming	10.20	10.17	0.50
Questionable	10.74	10.72	0.34
AGN	10.89	10.86	0.33

Figure 6: Stellar Mass Distributions

³The quality cut sample of 320,759 galaxies.

CLASS	MEDIAN	MEAN	STD
Star Forming	39.60	39.66	0.63
Questionable	39.62	39.65	0.47
AGN	39.84	39.96	0.64

Figure 7: [O III] Luminosity Distributions

4 Discussion

The total AGN fraction of the quality cut section of 320,759 galaxies was 14.2%, which is consistent with known fractions of AGN within the local universe. AGN were also more massive on average, peaking past $\log_{10}(\frac{M_*}{M_\odot}) = 11$, and had higher mean [O III] luminosity, with a tail extending past $\log_{10} L_{[OIII]} = 42$.

This result relatively matches what is currently known about AGN. Of course, the dataset was a very limited sample of galaxies because of the very small redshift range, and it would have been very interesting to look at a much larger sample and see the distributions of class, stellar mass, and [O III] luminosity across redshift bins. Because this study only examined within the very local universe, it is limited in significant contributions to the overall understanding of cosmological evolution across time.

It is also important to consider that the "questionable" class was not examined. While these are very likely not AGN because of they do not produce strong enough blamer line emissions, this composite group is still distinct from normal starforming galaxies. Perhaps they are very massive starburst galaxies, or galaxies experiencing shocks, but an examination of this class and a deeper comparison to true AGN should be done in the future.

However, the dataset chosen is one to the strongest for this sort of examination. It is thorough and the value-added catalogs provide very accurate data for all the needed probes. The Kauffman and Kewley equations are also very well-proven to be a good standard of AGN classification. These catalogs and equations have thoroughly increased the accuracy of this study.

Overall, the research done in this paper was a successful investigation into the fraction and properties of active galactic nuclei. These properties are very important for the understanding of black hole evolution and accretion, and how massive galaxies who host accreting supermassive black holes are affected. While the Milky Way itself is not massive enough to host an AGN, at only $\log_{10}(\frac{M_*}{M_\odot}) \cong 10.41$ according to Lian et al. (2025), it still hosts a SMBH, Sagittarius A*, at it's center. Having knowledge of the effects of extreme SMBH accretion in massive galaxies (AGN), allows us to approximate the effects on our own galaxy if Sagittarius A* begins to accrete huge amounts of mass. It is very crucial to emphasize the importance of AGN research in the future of astronomy and astrophysics as these objects truly hold so much information on galactic formation and evolution, and the cosmological web of our universe.

5 Images of Classified AGN Galaxies

Here are some images of randomly chosen AGN classified galaxies. These images are provided by the SDSS DR7 Navigation Tool. Right ascension and declination are included.



Figure 9: $\text{ra} = 209.77393$, $\text{dec} = 55.699425$



Figure 8: $\text{ra} = 145.68114$, $\text{dec} = -0.86722344$



Figure 10: $\text{ra} = 212.0057$, 14.986399



Figure 12: $ra = 196.73401$, $dec = 55.94665$



Figure 11: $ra = 191.28258$, $dec = 57.134197$

6 Acknowledgements

I would like to thank the MPA/JHU and SDSS team for the data processing and value-added catalogs, and for making the DR7 release publicly available.

I would like to thank the team of both Kauffman et al. (2003) and Kewley et al. (2001) for their work in creating the demarcation lines used in this study.

I would like to thank Claire Williams and James Larkin at UCLA for their lectures on extragalactic astronomy, which helped in creating this paper.

References

- [1] Baldwin, J. A., Phillips, M. M., & Terlevich, R. (1981). Classification parameters for the emission-line spectra of extragalactic objects. *Publications of the Astronomical Society of the Pacific*, 93(551), 5. <https://doi.org/10.1086/130766>
- [2] Bansal, A. (2025). Systematic Biases in BPT Diagram Classifications from Emission-line Signal-to-noise Cuts. *Research Notes of the AAS*, 9(8), 221. <https://doi.org/10.3847/2515-5172/adfc53>
- [3] Data products: Value-added catalogs - SDSS DR7. (2026). Sdss.org. https://classic.sdss.org/dr7/products/value_added/index.php

- [4] Juneau, S., et al. (2014). Active galactic nuclei emission line diagnostics and the mass-metallicity relation up to redshift $z \sim 2$. *The Astrophysical Journal*, 788(1), 88. <https://doi.org/10.1088/0004-637x/788/1/88>
- [5] Kaschinski, C. B., & Ercolano, B. (2013). Effects of shocks in stellar atmosphere models on the emission line spectrum of surrounding H II regions. *Monthly Notices of the Royal Astronomical Society*, 431(1), 493–501. <https://doi.org/10.1093/mnras/stt173>
- [6] Kauffmann, G., Heckman, T. M., Tremonti, C., Brinchmann, J., Charlot, S., White, S. D. M., Ridgway, S. E., Brinkmann, J., Fukugita, M., Hall, P. B., Ivezić, Ž., Richards, G. T., & Schneider, D. P. (2003). The host galaxies of active galactic nuclei. *Monthly Notices of the Royal Astronomical Society*, 346(4), 1055–1077. <https://doi.org/10.1111/j.1365-2966.2003.07154.x>
- [7] Kewley, L. J., Dopita, M. A., Sutherland, R., Heisler, C. A., & Trevena, J. (2001). Theoretical modeling of starburst galaxies. *The Astrophysical Journal*, 556(1), 121–140. <https://doi.org/10.1086/321545>
- [8] Robitaille, T. P., Tollerud, E. J., Greenfield, P., et al. (2013). Astropy: A community Python package for astronomy. *Astronomy & Astrophysics*, 558, A33. <https://doi.org/10.1051/0004-6361/201322068>
- [9] SDSS Data Products: `gal_info`. (2026). Mpa-Garching.mpg.de. https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/SDSS_info.html
- [10] SDSS Data Products: `gal_line`. (2026). Mpa-Garching.mpg.de. https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/SDSS_line.html
- [11] Stellar mass estimates. (2026). Mpa-Garching.mpg.de. <https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/Data/stellarmass.html>
- [12] Taylor, M. (2017). TOPCAT: Desktop exploration of tabular data for astronomy and beyond. *Informatics*, 4(3), 18. <https://doi.org/10.3390/informatics4030018>
- [13] The MPA-JHU DR7 release of spectrum measurements. (n.d.). [Wwwmpa.mpa-garching.mpg.de](https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/). <https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/>
- [14] Veilleux, S., & Osterbrock, D. E. (1987). Spectral classification of emission-line galaxies. *NASA Conference Publication*, 2466(63), 737–740. <https://ui.adsabs.harvard.edu/abs/1987NASCP2466..737V/abstract>
- [15] Gaskell, C. M., & Sparke, L. S. (1986). Line variations in quasars and Seyfert galaxies. *The Astrophysical Journal*, 305, 175–186.
- [16] Harrison, C. M., Costa, T., Tadhunter, C. N., Flütsch, A., Kakkad, D., Perna, M., & Vietri, G. (2018). AGN outflows and feedback twenty years on. *Nature Astronomy*, 2(3), 198–205. <https://doi.org/10.1038/s41550-018-0403-6>
- [17] Oke, J. B. (1978). Seyfert galaxies and quasars. *Hale Observatories, California Institute of Technology, Carnegie Institution of Washington*. <https://articles.adsabs.harvard.edu/pdf/1978JRASC..72..1210>

- [18] Perry’s AGN Page. (n.d.). Lweb.cfa.harvard.edu. <https://lweb.cfa.harvard.edu/~pberlind/whipple/agn.html>
- [19] Physical properties for galaxies and active galactic nuclei in the Sloan Digital Sky Survey: Data catalogues from SDSS studies at MPA/JHU. (2026). Mpa-Garching.mpg.de. https://wwwmpa.mpa-garching.mpg.de/SDSS/index_old.html
- [20] Rieke, G. H. (1978). The infrared emission of Seyfert galaxies. *The Astrophysical Journal*, 226, 550–558.
- [21] Shields, G. A. (1999). A brief history of active galactic nuclei. *Publications of the Astronomical Society of the Pacific*, 111(760), 661–678. <https://doi.org/10.1086/316378>
- [22] The MPA-JHU DR7 release of spectrum measurements. (2026). Mpa-Garching.mpg.de. <https://wwwmpa.mpa-garching.mpg.de/SDSS/DR7/#platelist>
- [23] SDSS DR7. (2024). Sdss.org. <https://classic.sdss.org/dr7/>
- [24] Lian, J., et al. 2025, , 990, L37, 10.3847/2041-8213/adfc73