

Measurement of π^0 and η mesons in oxygen-oxygen collisions at
 $\sqrt{s_{\text{NN}}} = 5.36$ TeV using Machine Learning-based photon
identification in ALICE Run 3

Supplemental figures

BDT working point optimisation in oxygen–oxygen collisions at
 $\sqrt{s_{\text{NN}}} = 5.36$ TeV

by Charlize Gohmann

Contents

1	BDT working point optimisation for $P_{\text{primary } \gamma} > 95\%$ and $\varepsilon_{\text{primary } \gamma} > 90\%$	2
----------	---	----------

1 BDT working point optimisation for $P_{\text{primary } \gamma} > 95\%$ and $\varepsilon_{\text{primary } \gamma} > 90\%$




Figure 1: BDT working point optimisation for all p_T bins in the centrality range 0-10 %







Figure 2: BDT working point optimisation for all p_T bins in the centrality range 10-20 %




(a) 0-0.3 GeV/c




(b) 0.3-0.5 GeV/c




(c) 0.5-0.8 GeV/c



(d) 0.8-1.0 GeV/c



(e) 1.0-2.0 GeV/c



(f) 2.0-10000 GeV/c

Figure 3: BDT working point optimisation for all p_T bins in the centrality range 20-40 %




Figure 4: BDT working point optimisation for all p_T bins in the centrality range 40-60 %




Figure 5: BDT working point optimisation for all p_T bins in the centrality range 60-80 %




Figure 6: BDT working point optimisation for all p_T bins in the centrality range 80-100 %