SYNTHETIC DOCUMENT. NOT A REAL PATIENT. FOR DEMO AND TESTING ONLY.

Northbrook General Hospital

Cardiology · Kestrel Ward, Coronary Care (Ward 2)
1 Example Road, Northbrook NB1 0AA (fictional)
Switchboard: 0000 000 000 · Ward: 0000 000 002 (fictional numbers)

DISCHARGE SUMMARY

Our ref: CA/SYN-0004/16-10
Date: Friday 16 October 2026

To: Dr C. Example, Hillside Practice, 2 Hill Road, Northbrook NB5 0EF (fictional) **cc:** Patient

Patient:	Mr Daniel Okafor	**Hospital no.:**	SYN-0004
Date of birth:	5 February 1981 (age 45)	**Address:**	9 Demo Avenue, Northbrook NB5 0RR (fictional)
Admitted:	Tuesday 13 October 2026 (emergency, chest pain)	**Discharged:**	Friday 16 October 2026
Length of stay:	3 days	**Discharged to:**	Home, lives alone
Consultant:	Dr S. Sample, Consultant Cardiologist	**Allergies:**	**Penicillin (rash)**

DIAGNOSIS

Heart attack (NSTEMI). Type 2 diabetes (known).

PROCEDURE

Coronary angiogram through the right wrist on Wednesday 14 October 2026. One narrowed artery was opened and a stent was put in. No complications.

HOSPITAL STAY

No further chest pain after the stent. The heart scan showed mildly reduced pumping. Daniel is walking around the ward and up one flight of stairs without symptoms. He smokes 10 cigarettes a day and has been referred to the stop smoking service.

MEDICINES TO TAKE AT HOME

Medicine	How to take it	Status
1. Aspirin 75 mg dispersible tablets.	Take 1 tablet once a day in the morning with food. Take for life. Do not stop unless your cardiologist tells you.	New
2. Ticagrelor 90 mg tablets.	Take 1 tablet twice a day (morning and evening) for 12 months, until Thursday 14 October 2027. Do not stop early, as this protects your stent.	New
3. Atorvastatin 80 mg tablets.	Take 1 tablet once a day at night.	New
4. Bisoprolol 2.5 mg tablets.	Take 1 tablet once a day in the morning.	New
5. Ramipril 2.5 mg capsules.	Take 1 capsule once a day. Your GP will increase the dose after your blood test.	New
6. Lansoprazole 15 mg capsules.	Take 1 capsule once a day in the morning before breakfast, to protect your stomach.	New
7. GTN spray, 400 micrograms per spray.	Only if you get chest pain: sit down and spray 1 or 2 puffs under your tongue. If the pain has not gone after 5 minutes, take another dose.	New
8. Metformin 500 mg tablets.	Take 1 tablet twice a day, with breakfast and with your evening meal.	Continue

Usual medicines before admission: metformin only. Do not take ibuprofen or other anti-inflammatory painkillers.

DAILY ACTIVITIES

- Walk every day. Start with 10 minutes on flat ground and add a few minutes each day, building up to 30 minutes.
- Check your blood sugar before breakfast and before your evening meal, as you did before.
- Rest in the afternoon if you feel tired. Tiredness is normal for a few weeks.

DIET

- Eat a Mediterranean-style diet: more vegetables, fish, beans and olive oil; less red meat, butter and salt.
- Keep alcohol to no more than 14 units a week, spread over 3 or more days.
- Stop smoking. The stop smoking service will call you this week.

WRIST CARE

- Keep the wrist dressing dry for 24 hours, then you can remove it.
- Do not lift anything heavier than a full kettle with your right hand for 1 week.

All names, places, numbers and details in this letter are invented. It is not medical advice.

SYNTHETIC DOCUMENT. NOT A REAL PATIENT. FOR DEMO AND TESTING ONLY.

RESTRICTIONS

- Do not drive a car for 1 week, until Friday 23 October 2026.
- You must not drive a bus or lorry for at least 6 weeks, and you must tell the DVLA.
- No heavy lifting or hard exercise until your cardiac rehabilitation assessment.
- Return to work: to be discussed at your cardiac rehabilitation assessment.

FOLLOW-UP APPOINTMENTS

- Blood test (kidney function and potassium): Thursday 29 October 2026, 8:40 am, Hillside Practice.
- Cardiac rehabilitation assessment: Tuesday 3 November 2026, 1:30 pm, Heart Centre, Northbrook General Hospital.
- Cardiology clinic in 6 to 8 weeks. You will get a letter with the date and time.

WARNING SIGNS

Call Kestrel Ward on 0000 000 002 (fictional number) or your GP if you have:

- Pain, swelling or a growing lump at the wrist where the tube went in.
- Black, tarry poo, or blood in your urine.
- Feeling dizzy or faint, or a very slow pulse.
- Mild breathlessness. This can happen with ticagrelor. Tell your GP, but do not stop taking it.

Call emergency services straight away if your chest pain is still there 5 minutes after your second dose of GTN spray.
Call emergency services straight away if bleeding from your wrist does not stop after 10 minutes of firm pressure.
Emergency services number: **555-555-0199** (fictional number)

INFORMATION FOR YOUR GP

- NSTEMI treated with a single drug-eluting stent to the left anterior descending artery.
- Dual antiplatelet therapy for 12 months. Please up-titrate ramipril after the blood test on 29 October 2026.
- Group 2 driving licence holder (bus driver). DVLA advice given.

Signed: Dr D. Example, Cardiology Registrar, on behalf of Dr S. Sample, Consultant Cardiologist
Bleep 0002 (fictional) · Friday 16 October 2026, 4:30 pm

Checked by ward pharmacist:
S. Example (fictional), 16 October 2026

All names, places, numbers and details in this letter are invented. It is not medical advice.