

# Diff in Diffs and Synthetic Control

```
# Install packages
if (!require("pacman")) install.packages("pacman")

## Loading required package: pacman
#install.packages("devtools") # download developer tools package
library(devtools)

## Loading required package: usethis
devtools::install_github("ebenmichael/augsynth")

## Skipping install of 'augsynth' from a github remote, the SHA1 (0f141c94) has not changed since last .
##   Use `force = TRUE` to force installation

pacman::p_load(# Tidyverse packages including dplyr and ggplot2
               tidyverse,
               ggthemes,
               augsynth)

# you might need to update the 'rlang' package here if you've had r for a while and haven't updated

set.seed(44)
```

## Introduction

In this lab we will explore difference-in-differences estimates and a newer extension, synthetic control. The basic idea behind both of these methods is simple - assuming two units are similar in a pre-treatment period and one undergoes treatment while the other stays in control, we can estimate a causal effect by taking three differences. First we take the difference between the two in the pre-treatment period, then take another difference in the post-treatment period. Then we take a difference between these two differences (hence the name difference in differences). Let's see how this works in practice!

## Basic DiD

We'll use the kansas dataset that comes from the `augsynth` package. Our goal here is to estimate the effect of the 2012 Kansas tax cuts on state GDP. Let's take a look at our dataset:

```
data(kansas)
summary(kansas)
```

##	fips	year	qtr	state
##	Min. : 1.00	Min. :1990	Min. :1.000	Length:5250
##	1st Qu.:17.00	1st Qu.:1996	1st Qu.:1.000	Class :character
##	Median :29.50	Median :2003	Median :2.000	Mode :character

```

## Mean :29.32 Mean :2003 Mean :2.486
## 3rd Qu.:42.00 3rd Qu.:2009 3rd Qu.:3.000
## Max. :56.00 Max. :2016 Max. :4.000
##
## gdp revenuepop rev_state_total rev_local_total
## Min. : 11509 Min. : 1335 Min. : 1668 Min. : 550
## 1st Qu.: 55151 1st Qu.: 3057 1st Qu.: 7026 1st Qu.: 3268
## Median : 130650 Median : 3628 Median : 13868 Median : 10041
## Mean : 228237 Mean : 3851 Mean : 20813 Mean : 17197
## 3rd Qu.: 276303 3rd Qu.: 4365 3rd Qu.: 24405 3rd Qu.: 18774
## Max. :2568986 Max. :14609 Max. :182530 Max. :143137
## NA's :2250 NA's :2850 NA's :2850
## popestimate qtrly_estabs_count month1_emplvl month2_emplvl
## Min. : 453690 Min. : 15133 Min. : 178737 Min. : 178587
## 1st Qu.: 1652585 1st Qu.: 48170 1st Qu.: 657056 1st Qu.: 663786
## Median : 3997978 Median : 108822 Median : 1675988 Median : 1684341
## Mean : 5767107 Mean : 161021 Mean : 2482331 Mean : 2494933
## 3rd Qu.: 6611215 3rd Qu.: 188730 3rd Qu.: 2990530 3rd Qu.: 2993158
## Max. :39250017 Max. :1448488 Max. :16600851 Max. :16633834
##
## month3_emplvl total_qtrly_wages taxable_qtrly_wages avg_wkly_wage
## Min. : 181521 Min. :8.811e+08 Min. :0.000e+00 Min. : 301.0
## 1st Qu.: 667492 1st Qu.:5.403e+09 1st Qu.:0.000e+00 1st Qu.: 515.2
## Median : 1699044 Median :1.362e+10 Median :1.096e+09 Median : 658.0
## Mean : 2510204 Mean :2.402e+10 Mean :3.776e+09 Mean : 674.8
## 3rd Qu.: 3016494 3rd Qu.:2.973e+10 3rd Qu.:4.177e+09 3rd Qu.: 804.0
## Max. :16606038 Max. :2.753e+11 Max. :7.689e+10 Max. :1792.0
##
## year_qtr treated gdpcapita lngdp
## Min. :1990 Min. :0.000000 Min. :15029 Min. : 9.351
## 1st Qu.:1996 1st Qu.:0.000000 1st Qu.:27989 1st Qu.:10.918
## Median :2003 Median :0.000000 Median :36449 Median :11.780
## Mean :2003 Mean :0.003048 Mean :37808 Mean :11.754
## 3rd Qu.:2010 3rd Qu.:0.000000 3rd Qu.:45531 3rd Qu.:12.529
## Max. :2016 Max. :1.000000 Max. :84382 Max. :14.759
##
## lngdpcapita revstatecapita revlocalcapita emplvl1capita
## Min. : 9.618 Min. : 2021 Min. : 883.6 Min. :0.3249
## 1st Qu.:10.240 1st Qu.: 2903 1st Qu.:2012.4 1st Qu.:0.4113
## Median :10.504 Median : 3380 Median :2428.3 Median :0.4356
## Mean :10.486 Mean : 3742 Mean :2480.2 Mean :0.4368
## 3rd Qu.:10.726 3rd Qu.: 4048 3rd Qu.:2819.4 3rd Qu.:0.4621
## Max. :11.343 Max. :20353 Max. :7160.9 Max. :1.0524
## NA's :2850 NA's :2850
## emplvl2capita emplvl3capita emplvlcapita totalwagescapita
## Min. :0.3251 Min. :0.3289 Min. :0.3269 Min. : 1493
## 1st Qu.:0.4138 1st Qu.:0.4163 1st Qu.:0.4138 1st Qu.: 2941
## Median :0.4378 Median :0.4406 Median :0.4378 Median : 3787
## Mean :0.4390 Mean :0.4420 Mean :0.4393 Mean : 3869
## 3rd Qu.:0.4644 3rd Qu.:0.4676 3rd Qu.:0.4644 3rd Qu.: 4608
## Max. :1.0507 Max. :1.0513 Max. :1.0515 Max. :10275
##
## taxwagescapita avgwklywagecapita estabscapita abb
## Min. : 0.0 Min. : 301.0 Min. :0.01992 Length:5250

```

```
## 1st Qu.: 0.0 1st Qu.: 515.2 1st Qu.:0.02553 Class :character
## Median : 355.7 Median : 658.0 Median :0.02845 Mode :character
## Mean : 728.8 Mean : 674.8 Mean :0.02928
## 3rd Qu.:1224.4 3rd Qu.: 804.0 3rd Qu.:0.03211
## Max. :5254.4 Max. :1792.0 Max. :0.07071
##
```


We have a lot of information here! We have quarterly state GDP from 1990 to 2016 for each U.S. state, as well as some other covariates. Let's begin by adding a treatment indicator to Kansas in Q2 2012 and onward.

```
kansas <- kansas %>%
  select(year, qtr, year_qtr, state, treated, gdp, lngdpcapita, fips) %>%
  mutate(treatment = ifelse(state == "Kansas" & year_qtr >= 2012.50,
                             1,
                             0))
head(kansas)
```

```
## # A tibble: 6 x 9
##   year   qtr year_qtr state   treated   gdp lngdpcapita fips treatment
##   <dbl> <dbl>   <dbl> <chr>   <dbl>   <dbl>   <dbl> <dbl>   <dbl>
## 1  1990     1    1990 Alabama     0 71610     9.78     1         0
## 2  1990     2    1990. Alabama     0 72718.     9.79     1         0
## 3  1990     3    1990. Alabama     0 73826.     9.80     1         0
## 4  1990     4    1991. Alabama     0 74935.     9.82     1         0
## 5  1991     1    1991 Alabama     0 76043     9.83     1         0
## 6  1991     2    1991. Alabama     0 77347.     9.84     1         0
```

One approach might be to compare Kansas to itself pre- and post-treatment. If we plot state GDP over time we get something like this:

```
kansas %>%
  filter(state == 'Kansas') %>%
  ggplot() +
  geom_point(aes(x = year_qtr, y = lngdpcapita)) +
  geom_vline(xintercept = 2012.5) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text()) +
  ggtitle('Kansas State GDP Per Capita Over Time') +
  xlab('Year-Quarter') +
  ylab('State GDP Per Capita')
```




**Question:** Looks like GDP went up after the tax cut! What is the problem with this inference?

**Solution:** It looks like GDP went up after the tax cut, but we have no way of telling whether it went up because of the tax cut or went up because it would have otherwise. In short, we need to compare the treated Kansas to a counterfactual for if taxes weren't cut.

Ideally, we would like to compare treated Kansas to control Kansas. Because of the fundamental problem of causal inference, we will never observe both of these conditions though. The core idea behind DiD is that we could instead use the fact that our treated unit was similar to a control unit, and then measure the differences between them. Perhaps we could choose neighboring Colorado:

```
kansas %>%
  filter(state %in% c("Kansas", "Colorado")) %>%
  filter(year_qtr >= 2012.5 & year_qtr <= 2012.75) %>%
  ggplot() +
  geom_point(aes(x = year_qtr,
                 y = lngdpcapita,
                 color = state)) +
  geom_line(aes(x = year_qtr,
                y = lngdpcapita,
                color = state)) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text()) +
  ggtitle('Colorado and Kansas GDP \n before/after Kansas tax cut') +
  xlab('Year-Quarter') +
  ylab('State GDP')
```



This is basically what Card-Krueger (1994) did measuring unemployment rates among New Jersey and Pennsylvania fast food restaurants.

**Challenge:** Try writing a simple DiD estimate using dplyr/tidyr (use subtraction instead of a regression):

```
# kansas-colorado
kc <- kansas %>%
  filter(state %in% c("Kansas", "Colorado")) %>%
  filter(year_qtr >= 2012.5 & year_qtr <= 2012.75)
```

```
# pre-treatment difference
```

```
pre_diff <- kc %>%
  filter(year_qtr == 2012.5) %>%
  select(state,
         lngdpcapita) %>%
  spread(state,
         lngdpcapita) %>%
  summarise(Colorado - Kansas)
```

```
# post-treatment difference
```

```
post_diff <- kc %>%
  filter(year_qtr == 2012.75) %>%
  select(state,
         lngdpcapita) %>%
  spread(state,
```

```

      lngdpcapita) %>%
    summarise(Colorado - Kansas)

# diff-in-diffs

diff_in_diffs <- post_diff - pre_diff
diff_in_diffs

##    Colorado - Kansas
## 1          0.003193447

```

Looks like our treatment effect is about .003 (in logged thousands dollars per capita). Again this is the basic idea behind Card-Krueger.

**Question:** Why might there still be a problem with this estimate?

**Answer:** We just assumed that Colorado was similar to Kansas because they are neighbors - we don't really have evidence for this idea.


## Parallel Trends Assumptions

One of the core assumptions for difference-in-differences estimation is the “parallel trends” or “constant trends” assumption. Essentially, this assumption requires that the difference between our treatment and control units are constant in the pre-treatment period. Let's see how Kansas and Colorado do on this assumption:

```

kansas %>%
  filter(state %in% c("Kansas", "Colorado")) %>%
  ggplot() +
  geom_point(aes(x = year_qtr,
                 y = lngdpcapita,
                 color = state)) +
  geom_line(aes(x = year_qtr,
                y = lngdpcapita,
                color = state)) +
  geom_vline(aes(xintercept = 2012.5)) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text()) +
  ggtitle('Colorado and Kansas GDP \n before/after Kansas tax cut') +
  xlab('Year-Quarter') +
  ylab('State GDP')


```



The two lines somewhat move together, but the gap does grow and shrink at various points over time. The most concerning part here is that the gap quickly shrinks right before treatment. What do we do if we do not trust the parallel trends assumption? Perhaps we pick a different state.

**Challenge:** Choose another state that you think would be good to try out, and plot it alongside Kansas and Colorado.

```
kansas %>%
  filter(state %in% c("Kansas", "Colorado", "Missouri")) %>%
  ggplot() +
  geom_point(aes(x = year_qtr,
                 y = lngdpcapita,
                 color = state)) +
  geom_line(aes(x = year_qtr,
                y = lngdpcapita,
                color = state)) +
  geom_vline(aes(xintercept = 2012.5)) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text()) +
  ggtitle('Colorado and Kansas GDP \n before/after Kansas tax cut') +
  xlab('Year-Quarter') +
  ylab('State GDP')
```



**Question:** Would you pick Colorado or your choice? be the more plausible control unit in this case? Why?

**Solution:** There is a good argument for both of them (Missouri in this case). However, the gap between Colorado and Kansas closes quickly before the treatment period, and similarly it grows between between Kansas and Missouri at the same point.

Selecting comparative units this way can be hard to justify theoretically, and sometimes we do not have a good candidate. What can we do then? This is where synthetic control comes in.

## Synthetic Control

Synthetic control is motivated by the problem of choosing comparison units for comparative case studies. It aims to create a “synthetic” version of the treatment unit by combining and weighting covariates from other units (“donors”). In this case, we would construct a synthetic Kansas by creating a weighted average of the other 49 U.S. states. Ideally, the synthetic unit would match the treatment unit in the pre-treatment periods.

For constructing a synthetic control, we are going to use the `augsynth` library. The basic syntax for this library is:

```
augsynth(outcome ~ trt, unit, time, t_int, data)
```

```
syn <- augsynth(lngdpcapita ~ treated, state, year_qtr, kansas,
               progfunc = "None", scm = T)
```

```
## One outcome and one treatment time found. Running single_augsynth.
```


```
summary(syn)
```



```
##
## Call:
## single_augsynth(form = form, unit = !!enquo(unit), time = !!enquo(time),
##   t_int = t_int, data = data, progfunc = "None", scm = ..2)
##
## Average ATT Estimate (p Value for Joint Null): -0.029   ( 0.318 )
## L2 Imbalance: 0.083
## Percent improvement from uniform weights: 79.5%
##
## Avg Estimated Bias: NA
##
## Inference type: Conformal inference
##
##   Time Estimate 95% CI Lower Bound 95% CI Upper Bound p Value
## 2012.25   -0.018           -0.045           0.006   0.111
## 2012.50   -0.041           -0.070          -0.015   0.022
## 2012.75   -0.033           -0.062          -0.007   0.044
## 2013.00   -0.019           -0.046           0.005   0.111
## 2013.25   -0.029           -0.053          -0.005   0.044
## 2013.50   -0.046           -0.073          -0.022   0.022
## 2013.75   -0.032           -0.056          -0.010   0.022
## 2014.00   -0.045           -0.074          -0.018   0.022
## 2014.25   -0.043           -0.074          -0.014   0.022
## 2014.50   -0.029           -0.061           0.000   0.044
## 2014.75   -0.018           -0.053           0.011   0.144
## 2015.00   -0.029           -0.066           0.005   0.078
## 2015.25   -0.019           -0.051           0.010   0.122
## 2015.50   -0.022           -0.056           0.007   0.111
## 2015.75   -0.019           -0.055           0.013   0.189
## 2016.00   -0.028           -0.067           0.008   0.100
```


We can use the built in plot function to see how Kansas did relative to synthetic Kansas:

```
plot(syn)
```




We can see which donors contributed the most to the synthetic Kansas:

```
# Convert weights to dataframe
data.frame(syn$weights) %>%
  # change index to a column
  tibble::rownames_to_column('State') %>%
  ggplot() +
  # stat = identity to take the literal value instead of a count for geom_bar()
  geom_bar(aes(x = State,
               y = syn.weights),
            stat = 'identity') +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        axis.text.x = element_text(angle = 90)) +
  ggtitle('Synthetic Control Weights') +
  xlab('State') +
  ylab('Weight')
```



Surprisingly, only a few units ended up contributing! Let's take a closer look at the ones that did:

```
data.frame(syn$weights) %>%
  tibble::rownames_to_column('State') %>%
  filter(syn.weights > 0) %>%
  ggplot() +
  geom_bar(aes(x = State,
               y = syn.weights),
            stat = 'identity') +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        axis.text.x = element_text(angle = 90)) +
  ggtitle('Synthetic Control Weights') +
  xlab('State') +
  ylab('Weight')
```



## Synthetic Control Augmentation

The main advantage of the `asynth` package is that it allows for “augmented synthetic control”. One of the main problems with synthetic control is that if the pre-treatment balance between treatment and control outcomes is poor, the estimate is not valid. Specifically, they advocate for using L2 imbalance, which he first encountered as the penalty that ridge regression uses. L2 uses “squared magnitude” of the coefficient to penalize a particular feature.

```
# Aniket's method for getting the underlying data
syn_sum <- summary(syn)

kansas_synkansas <- kansas %>%
  filter(state == "Kansas") %>%
  bind_cols(difference = syn_sum$att$Estimate) %>%
  mutate(synthetic_kansas = lngdpcapita + difference)


# Plot

kansas_synkansas %>%
  ggplot() +
  geom_point(aes(x = year_qtr,
                 y = lngdpcapita,
                 color = 'Kansas')) +
  geom_line(aes(x = year_qtr,
                y = lngdpcapita,
```

```

    color = 'Kansas')) +
geom_point(aes(x = year_qtr,
               y = synthetic_kansas,
               color = 'Synthetic Kansas')) +
geom_line(aes(x = year_qtr,
              y = synthetic_kansas,
              color = 'Synthetic Kansas')) +
scale_color_manual(values = c('Kansas' = 'red', 'Synthetic Kansas' = 'blue')) +
geom_vline(aes(xintercept = 2012.5)) +
theme_fivethirtyeight() +
theme(axis.title = element_text()) +
ggtitle('Kansas and Synthetic Kansas') +
xlab('Year-Quarter') +
ylab('State GDP Per Capita')

```



**Question:** How does pre-treatment matching between Kansas and Synthetic Kansas look here?

**Answer:** Pretty good! We may not need to augment this synthetic control, though let's try anyway.

```

ridge_syn <- augsynth(lngdpcapita ~ treated, state, year_qtr, kansas,
                     progfunc = "ridge", scm = T)

```

```
## One outcome and one treatment time found. Running single_augsynth.
```

```
summary(ridge_syn)
```


```
##
```

```
## Call:
```

```
## single_augsynth(form = form, unit = !!enquo(unit), time = !!enquo(time),
##   t_int = t_int, data = data, progfunc = "ridge", scm = ..2)
##
## Average ATT Estimate (p Value for Joint Null): -0.040   ( 0.067 )
## L2 Imbalance: 0.062
## Percent improvement from uniform weights: 84.7%
##
## Avg Estimated Bias: 0.011
##
## Inference type: Conformal inference
##
##   Time Estimate 95% CI Lower Bound 95% CI Upper Bound p Value
## 2012.25    -0.022           -0.044           0.003    0.056
## 2012.50    -0.047           -0.076          -0.018    0.022
## 2012.75    -0.043           -0.071          -0.010    0.022
## 2013.00    -0.030           -0.055          -0.004    0.033
## 2013.25    -0.041           -0.067          -0.012    0.022
## 2013.50    -0.059           -0.088          -0.030    0.022
## 2013.75    -0.045           -0.073          -0.019    0.022
## 2014.00    -0.058           -0.090          -0.026    0.022
## 2014.25    -0.055           -0.091          -0.020    0.022
## 2014.50    -0.041           -0.080          -0.006    0.033
## 2014.75    -0.029           -0.068           0.006    0.056
## 2015.00    -0.040           -0.082           0.000    0.056
## 2015.25    -0.030           -0.066           0.002    0.056
## 2015.50    -0.033           -0.072           0.003    0.056
## 2015.75    -0.029           -0.071           0.010    0.056
## 2016.00    -0.038           -0.087           0.004    0.056
```

Let's look at the weights:

```
data.frame(ridge_syn$weights) %>%
  tibble::rownames_to_column('State') %>%
  ggplot() +
  geom_bar(aes(x = State, y = ridge_syn.weights),
    stat = 'identity') +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
    axis.text.x = element_text(angle = 90)) +
  ggtitle('Synthetic Control Weights') +
  xlab('State') +
  ylab('Weight')
```



Notice how with the ridge augmentation, some weights are allowed to be negative now. Now let's go ahead and plot the ridge augmented synthetic Kansas alongside Kansas and synthetic Kansas:

```
ridge_sum <- summary(ridge_syn)


kansas_synkansas_ridgesynkansas <- kansas_synkansas %>%
  bind_cols(ridge_difference = ridge_sum$att$Estimate) %>%
  mutate(ridge_synthetic_kansas = lngdpcapita + ridge_difference)

kansas_synkansas_ridgesynkansas %>%
  ggplot() +
  geom_point(aes(x = year_qtr,
                 y = lngdpcapita,
                 color = 'Kansas')) +
  geom_line(aes(x = year_qtr,
                y = lngdpcapita,
                color = 'Kansas')) +
  geom_point(aes(x = year_qtr,
                 y = synthetic_kansas,
                 color = 'Synthetic Kansas')) +
  geom_line(aes(x = year_qtr,
                y = synthetic_kansas,
                color = 'Synthetic Kansas')) +
  geom_point(aes(x = year_qtr,
                 y = ridge_synthetic_kansas,
                 color = 'Ridge Synthetic Kansas')) +
  geom_line(aes(x = year_qtr,
```

```

    y = ridge_synthetic_kansas,
    color = 'Ridge Synthetic Kansas')) +
scale_color_manual(values = c('Kansas' = 'red',
                              'Synthetic Kansas' = 'blue',
                              'Ridge Synthetic Kansas' = 'green')) +
geom_vline(aes(xintercept = 2012.5)) +
theme_fivethirtyeight() +
theme(axis.title = element_text()) +
ggtitle('Kansas, Synthetic Kansas, Ridge Synthetic Kansas') +
xlab('Year-Quarter') +
ylab('State GDP Per Capita')

```



These all seem pretty good! Like we thought, augmentation did not necessarily improve the matches in this particular dataset. We can check the two L2 imbalances and see that we have reduced the overall imbalance a bit with our ridge model:

```
print(syn$l2_imbalance)
```

```
## [1] 0.08255471
```

```
print(ridge_syn$l2_imbalance)
```

```
## [1] 0.06151525
```

Finally, we can add covariates to our model if we would like:

```
data(kansas)
```



```

covsyn <- augsynth(lngdpcapita ~ treated | lngdpcapita + log(revstatecapita) +
                  log(revlocalcapita) + log(avgwkllywagecapita) +
                  estabscapita + emplvlcapita,
                  fips, year_qtr, kansas,
                  progfunc = "ridge", scm = T)

## One outcome and one treatment time found. Running single_augsynth.
summary(covsyn)

##
## Call:
## single_augsynth(form = form, unit = !!enquo(unit), time = !!enquo(time),
##   t_int = t_int, data = data, progfunc = "ridge", scm = ..2)
##
## Average ATT Estimate (p Value for Joint Null): -0.061 ( 0.118 )
## L2 Imbalance: 0.054
## Percent improvement from uniform weights: 86.6%
##
## Covariate L2 Imbalance: 0.005
## Percent improvement from uniform weights: 97.7%
##
## Avg Estimated Bias: 0.027
##
## Inference type: Conformal inference
##
##      Time Estimate 95% CI Lower Bound 95% CI Upper Bound p Value
## 2012.25    -0.021         -0.044         0.002    0.067
## 2012.50    -0.047         -0.076        -0.014    0.033
## 2012.75    -0.050         -0.083        -0.007    0.033
## 2013.00    -0.045         -0.074        -0.012    0.033
## 2013.25    -0.055         -0.088        -0.022    0.022
## 2013.50    -0.071         -0.105        -0.033    0.022
## 2013.75    -0.058         -0.091        -0.025    0.022
## 2014.00    -0.081         -0.119        -0.037    0.022
## 2014.25    -0.078         -0.121        -0.034    0.022
## 2014.50    -0.065         -0.114        -0.021    0.033
## 2014.75    -0.057         -0.110        -0.008    0.044
## 2015.00    -0.075         -0.124        -0.022    0.033
## 2015.25    -0.063         -0.106        -0.014    0.033
## 2015.50    -0.067         -0.106        -0.019    0.022
## 2015.75    -0.063         -0.101        -0.009    0.022
## 2016.00    -0.078         -0.122        -0.019    0.022

```

## Staggered Adoption

The last technique we'll look at is "staggered adoption" of some policy. In the original Hainmueller paper, states that already had similar cigarette taxes were discarded from the donor pool to create a synthetic California. But what if we were interested in the effect of a policy overall, for every unit that adopted treatment? The problem is, these units all choose to adopt treatment at different times. We could construct different synthetic controls for each one, or we can use a staggered adoption approach.

To explore this question, we'll continue using the `augsynth` package's vignette. This time we will load a dataset that examines the effect of states instituting mandatory collective bargaining agreements.

```
collective_bargaining <- read_delim("https://dataverse.harvard.edu/api/access/datafile/:persistentId?persistentId=doi:10.7927/H73T-6K9Q")

##
## -- Column specification -----
## cols(
##   .default = col_double(),
##   State = col_character()
## )
## i Use `spec()` for the full column specifications.
head(collective_bargaining)

## # A tibble: 6 x 23
##   year State Stateid avgteachsal YearCBrequired CBstatusby1990 ppexpend
##   <dbl> <chr>   <dbl>      <dbl>          <dbl>          <dbl>      <dbl>
## 1  1899 AK         1         NA            1970            2         NA
## 2  1900 AK         1         NA            1970            2         NA
## 3  1904 AK         1         NA            1970            2         NA
## 4  1909 AK         1         NA            1970            2         NA
## 5  1910 AK         1         NA            1970            2         NA
## 6  1912 AK         1         NA            1970            2         NA
## # ... with 16 more variables: avginstrucsal <dbl>, agr <dbl>, perinc <dbl>,
## #   pnwht <dbl>, purban <dbl>, ESWI <dbl>, studteachratio <dbl>,
## #   nonwageppexpend <dbl>, lnppexpend <dbl>, lnavginstrucsal <dbl>,
## #   lnavgteachsal <dbl>, lnnonwageppexpend <dbl>, CBrequired_SY <dbl>,
## #   CBeverrequired <dbl>, South <dbl>, idmap <dbl>
```

The main variables we'll use here are:

The dataset contains several important variables that we'll use:

- **year, State:** The state and year of the measurement
- **YearCBrequired:** The year that the state adopted mandatory collective bargaining
- **lnppexpend:** Log per pupil expenditures in 2010 dollars

Let's do some preprocessing before we estimate some models. We're going to remove DC and Wisconsin from the analysis and cabin our dataset to 1959 - 1997. Finally, we'll add a treatment indicator **cbr** which takes a 1 if the observation was a treated state after it adopted mandatory collective bargaining, or a 0 otherwise:

```
collective_bargaining_clean <- collective_bargaining %>%
  filter(!State %in% c("DC", "WI"),
         year >= 1959,
         year <= 1997) %>%
  mutate(YearCBrequired = ifelse(is.na(YearCBrequired),
                                Inf, YearCBrequired),
         cbr = 1 * (year >= YearCBrequired))
```

We're ready to start estimating a model! To do this, we use the **multisynth()** function that has the following signature:

```
multisynth(outcome ~ treatment, unit, time, nu, data, n_leads)
```

The key parameters here are **nu** and **n\_leads**. Staggered adoption uses multi-synthetic control which essentially pools together similar units and estimates a synthetic control for each pool. **nu** determines how much pooling to do. A value of 0 will fit a separate synthetic control for each model, whereas a value of 1 will pool all units together. Leaving this argument blank will have **augsynth** search for the best value of **nu** that minimizes L2 loss. **n\_leads** determines how many time periods to estimate in the post-treatment period.

```
# with a choice of nu
ppool_syn <- multisynth(lnppexpend ~ cbr, State, year,
                        nu = 0.5, collective_bargaining_clean, n_leads = 10)

# with default nu
ppool_syn <- multisynth(lnppexpend ~ cbr, State, year,
                        collective_bargaining_clean, n_leads = 10)

print(ppool_syn$nu)
```

```
## [1] 0.2618752
```

```
ppool_syn
```

```
##
## Call:
## multisynth(form = lnppexpend ~ cbr, unit = State, time = year,
##   data = collective_bargaining_clean, n_leads = 10)
##
## Average ATT Estimate: -0.010
```

After you've fit a model that you like, use the `summary()` function to get the ATT and balance statistics.


```
ppool_syn_summ <- summary(ppool_syn)
```

Next, plot the estimates for each state as well as the average average treatment effect (so average for all treated states). Try to do this with `ggplot()` instead of the built-in plotting function (hint: how did we get the dataframe with the estimates before?)

```
ppool_syn_summ$att %>%
  ggplot(aes(x = Time, y = Estimate, color = Level)) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 0) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        legend.position = "bottom") +
  ggtitle('Synthetic Controls for State Collective Bargaining') +
  xlab('Time') +
  ylab('Expenditure on Pupil Estimate')
```

```
## Warning: Removed 506 rows containing missing values (geom_point).
```

```
## Warning: Removed 506 row(s) containing missing values (geom_path).
```




```
ppool_syn_summ$att %>%
  ggplot(aes(x = Time, y = Estimate, color = Level)) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 0) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        legend.position = 'None') +
  ggtitle('Synthetic Controls for State Collective Bargaining') +
  xlab('Time') +
  ylab('Expenditure on Pupil Estimate') +
  facet_wrap(~Level)
```

```
## Warning: Removed 506 rows containing missing values (geom_point).
```

```
## Warning: Removed 506 row(s) containing missing values (geom_path).
```

## Synthetic Controls for State Collective Bargaining



We can also combine our observations into “time cohorts” or units that adopted treatment at the same time. Try adding `time_cohort = TRUE` to your `multisynth` function and see if your estimates differ. Plot these results as well.

```
ppool_syn_time <- multisynth(lnppexpend ~ cbr, State, year,
                             collective_bargaining_clean, n_leads = 10, time_cohort = TRUE)

ppool_syn_time_summ <- summary(ppool_syn_time)

ppool_syn_time_summ
```


```
##
## Call:
## multisynth(form = lnppexpend ~ cbr, unit = State, time = year,
##   data = collective_bargaining_clean, n_leads = 10, time_cohort = TRUE)
##
## Average ATT Estimate (Std. Error): -0.016 (0.023)
##
## Global L2 Imbalance: 0.005
## Scaled Global L2 Imbalance: 0.018
## Percent improvement from uniform global weights: 98.2
##
## Individual L2 Imbalance: 0.039
## Scaled Individual L2 Imbalance: 0.058
## Percent improvement from uniform individual weights: 94.2
##
## Time Since Treatment   Level      Estimate  Std.Error lower_bound upper_bound
```

```
##          0 Average  0.0038263026  0.02494062 -0.04462064  0.05367291
##          1 Average -0.0130748834  0.02444229 -0.06224330  0.03521447
##          2 Average  0.0018300044  0.02404155 -0.04419823  0.05035362
##          3 Average  0.0005232868  0.02679212 -0.05356853  0.05481323
##          4 Average -0.0184345032  0.02494437 -0.06676505  0.02833195
##          5 Average -0.0258163688  0.02603120 -0.07479496  0.02772436
##          6 Average -0.0217543090  0.02571936 -0.06989603  0.02927911
##          7 Average -0.0105432314  0.03101850 -0.06969423  0.05017729
##          8 Average -0.0262042318  0.03091288 -0.08398609  0.03114278
##          9 Average -0.0476919393  0.03067941 -0.10681797  0.01305787
```

```
ppool_syn_time_summ$att %>%
  ggplot(aes(x = Time, y = Estimate, color = Level)) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 0) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        legend.position = 'None') +
  ggtitle('Synthetic Controls for State Collective Bargaining') +
  xlab('Time') +
  ylab('Expenditure on Pupil Estimate') +
  facet_wrap(~Level)
```

```
## Warning: Removed 205 rows containing missing values (geom_point).
```

```
## Warning: Removed 205 row(s) containing missing values (geom_path).
```



Finally, we can add in augmentation. Again augmentation essentially adds a regularization penalty to the synthetic control weights. In the multisynth context, you may especially want to do this when the pre-treatment fit is poor for some of your units. There are a couple of different options for augmentation. One is to specify `fixed_effects = TRUE` in the multisynth call, and this will estimate unit fixed effects models after de-meaning each unit. We can also specify a `n_factors =` argument (substituting an integer in) to use the `gsynth` method that uses cross-validation to estimate the weights for multi-synthetic control.

Try creating an augmented synthetic control model. How do your balance and estimates compare?

```
scm_gsyn <- multisynth(lnppexpend ~ cbr, State, year,
                      collective_bargaining_clean, n_leads = 10,
                      fixedeff = T, n_factors = 2)

scm_gsyn_summ <- summary(scm_gsyn)

scm_gsyn_summ$att %>%
  ggplot(aes(x = Time, y = Estimate, color = Level)) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 0) +
  theme_fivethirtyeight() +
  theme(axis.title = element_text(),
        legend.position = 'None') +
  ggtitle('Augmented Synthetic Controls for State Collective Bargaining') +
  xlab('Time') +
  ylab('Expenditure on Pupil Estimate') +
  facet_wrap(~Level)
```

```
## Warning: Removed 506 rows containing missing values (geom_point).
```

```
## Warning: Removed 506 row(s) containing missing values (geom_path).
```

## Augmented Synthetic Controls for State Collection

